

Racial and ethnic differences in response to treatment for Marfan syndrome – Corrigendum

Rachel Ayers, Michael Kelleman, Glen Iannucci, Courtney McCracken and Matthew E. Oster

Corrigendum

Cite this article: Ayers R, Kelleman M, Iannucci G, McCracken C, and Oster ME (2021) Racial and ethnic differences in response to treatment for Marfan syndrome – Corrigendum. *Cardiology in the Young* 31: 1999. doi: [10.1017/S1047951121002341](https://doi.org/10.1017/S1047951121002341)

First published online: 10 June 2021

DOI: <https://doi.org/10.1017/S104795112100130X> Published online by Cambridge University Press: 13 April 2021

The authors apologise that this article was published with a typographical error within the abstract. The ‘N’ was missing from the word *non-Hispanic* in the first line of results. The correct abstract section should read as follows:

Results:

For aortic root enlargement by z score amongst non-Hispanic White patients, patients on losartan exhibited an annual z score change of -0.090 ± 0.016 , compared to -0.146 ± 0.015 for those on atenolol ($p = 0.01$), favouring atenolol. For Hispanic and non-Hispanic Black patients, there was no difference in primary or secondary outcomes between treatment groups.

The published article has been updated to correct this mistake.

Reference

Caliskan, S., & Bilgin, M. (2020). Non-linear analysis of heart rate variability for evaluating the acute effects of caffeinated beverages in young adults. *Cardiology in the Young*, 30(7), 1018–1023. doi: [10.1017/S1047951120001481](https://doi.org/10.1017/S1047951120001481)