

times tend to have higher health anxiety levels in comparison to healthy participants.

Disclosure of Interest: None Declared

EPV0317

TIME PERSPECTIVE OF THE PERSONALITY OF PATIENTS WHO HAD SEVERE AND MEDIUM COVID-19

E. R. Semenova^{1*}, E. Pervichko¹, A. Kulikova¹, I. Shishkova² and J. Konyukhovskaya¹

¹Moscow State University named after 'M. V. Lomonosov', Moscow and ²Ryazan State Medical University, Ryazan, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1666

Introduction: In connection with the COVID-19 pandemic and its consequences, the question of how personal time constructs are modified becomes highly relevant. In particular, it is important to understand the specifics of the personality's time perspective in patients who have undergone COVID-19 in varying degrees of severity.

Objectives: To study the time perspective of personality in patients who have undergone COVID-19 in medium and severe forms.

Methods: The study was conducted from February to April 2022. The first group of the sample (the medium form with hospitalization) consisted of 52 respondents (46.2% - men, 53.8% - women, M age = 31.2 years; S=6.7). The second group (severe form with hospitalization) consisted of 48 patients (60% - men and 40% - women, M age = 33.0 years; S=7.8). We used: "The questionnaire of the time perspective of the personality of F. Zimbardo (Short version)"; (Zimbardo, Boyd, 1997; Syrtsova, 2008), the "Scale of Time experience" questionnaire (Golovakha, Kronik, 2008) and the descriptive analysis.

Results: Respondents with a medium form of the disease have more developed hedonistic present (3.2±0.6), future (3.8±0.4) and positive past (3.7±0.6). The negative past (2.6±0.7) and fatalistic present (2.4±0.6) are the least represented in their lives. Respondents with a severe form have a more developed negative past (3.4±0.7), hedonistic present (3.4±0.4), future (3.7±0.5) and positive past (3.6±0.7). The fatalistic present is the least represented in their life (3.0±0.5). When analyzing the factors of time experiencing, it was revealed that respondents with a medium form perceive time as moderately continuous (3.4± 0.8), moderately tense (3.4± 0.8) due to pronounced emptiness and compactness, and also treat time on average not very positively (3.2± 1.0). Respondents with a severe form also perceive time as moderately continuous (3.4± 0.7), rather tense (4.1±1.1) due to pronounced saturation, compactness and rapidity, while they treat time on average moderately positively (2.5± 0.9).

Conclusions: The time perspective in patients with medium form is characterized by planning and achieving future goals, and these respondents also show a fairly high degree of acceptance of their own past, in which any experience is an experience that contributes to development and led to today's state. In respondents with a severe form, along with normative scores on the positive past scale, there is also an increase in negative perception of the past, which is

reflected in an increased degree of rejection of their own past, causing disgust, full of pain and frustration, as well as a hedonistic, risky attitude to time and life, while an orientation towards pleasure, excitement, excitement, enjoyment in the present and lack of concern for future consequences or sacrifices in favor of future rewards. Assistance and help to such respondents should be in the focus of specialists of the relevant profile.

Disclosure of Interest: None Declared

EPV0318

The Role of COVID-19 Pandemic Anxiety and Perceptions in COVID-19 Vaccination

E. R. Semenova^{1*}, J. Konyukhovskaya¹, S. Sviridova², E. Pervichko¹, O. Stepanova¹, O. Mitina¹ and I. Shishkova³

¹Moscow State University named after 'M. V. Lomonosov'; ²Pirogov Russian National Research Medical University, Moscow and ³Ryazan State Medical University, Ryazan, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1667

Introduction: Since stress and anxiety are significant manifestations of psychological distress during the COVID-19 pandemic, we studied their role in making a decision about vaccination.

Objectives: To study the relationship between the intention to be vaccinated against COVID-19 with health anxiety and stress levels.

Methods: The methodological complex includes the author's socio-demographic questionnaire (Pervichko, 2020, 2021, 2022); the questionnaire "Scale of perceived stress-10" (Ababkov, 2016); the questionnaire "Perceptions of the COVID-19 pandemic" (Pervichko et al., 2020), developed on the basis of the Russian-language version of the E. Broadbent's short questionnaire about the perception of disease (Broadbent, 2006); the State-Trait Anxiety Inventory (STAI) (Spielberger, 2002) and the "Short Health Anxiety Inventory" (Pervichko et al., 2020).

The study involved 232 respondents who did not have COVID-19 (average age – 29.1 ± 13.7 years). Among the respondents, 68.5% have already been vaccinated, 23.3% do not plan to be vaccinated and 8.2% plan to perform the procedure.

Results: The methodological complex includes the author's socio-demographic questionnaire (Pervichko, 2020, 2021, 2022); the questionnaire "Scale of perceived stress-10" (Ababkov, 2016); the questionnaire "Perceptions of the COVID-19 pandemic" (Pervichko et al., 2020), developed on the basis of the Russian-language version of the E. Broadbent's short questionnaire about the perception of disease (Broadbent, 2006); the State-Trait Anxiety Inventory (STAI) (Spielberger, 2002) and the "Short Health Anxiety Inventory" (Pervichko et al., 2020).

The study involved 232 respondents who did not have COVID-19 (average age – 29.1 ± 13.7 years). Among the respondents, 68.5% have already been vaccinated, 23.3% do not plan to be vaccinated and 8.2% plan to perform the procedure.

Conclusions: Higher health anxiety, situational anxiety, perceived stress, and greater perceived life threat due to coronavirus contribute to COVID-19 immunization procedures, which is accompanied by perceptions of greater control of the pandemic.

Disclosure of Interest: None Declared