

Acute malnutrition associated with MUAC among under-five children in tribal areas, India - A cross-sectional study

Shraboni Patra¹, Shashikant Sambharkar², Sheetal Harode³, Kalpana Barde⁴, Amta Pillewan⁵

¹Monitoring and Evaluation Expert, State Monitoring and Resource Cell, Bureau of Nutrition, Nagpur, Maharashtra, Email: shrageo@gmail.com

²Senior Scientific Officer, State Nutrition Bureau, Nagpur, Public Health Department, Maharashtra, Email: shashishambharkar@gmail.com

³Medical Officer, Nutrition Bureau, Nagpur, Email: girjamohane@yahoo.com

⁴Nutrition officer, Nutrition Bureau, Nagpur, Email: kalpanabarde@gmail.com

⁵Nutrition officer, Nutrition Bureau, Nagpur, Email: amita_pillewan@yahoo.com

Corresponding Author: Shraboni Patra; Email: shrageo@gmail.com

Short title: Acute Malnutrition among U5 Children in Tribal Areas



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980024002465

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Acknowledgements

The authors acknowledge the support of the Bureau of Nutrition, Nagpur, Government of Maharashtra for conducting the survey and UNICEF for providing technical support.

Financial support

The survey was carried out by the Nutrition Bureau, Public Health Department (PHD), Nagpur, Government of Maharashtra.

However, the funders had no role in the study design, analysis, decision to publish, or preparation of the manuscript.

Conflict of Interest

The authors have declared that there is no conflict of interest.

Authorship

SP conceptualised the study, analysed the data and prepared the primary draft and revised the manuscript; SS guided and supported in the survey, helped in manuscript preparation and its revision; SH, KB and AP helped in study design and data collection.

Ethical Standard Disclosure

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the Institutional Ethics Committee (IEC), State Institute of Health & Family Welfare, Nagpur. Verbal informed consent was obtained from all subjects/patients, and verbal consent was witnessed and formally recorded.

Abstract

Objective:

For the past three decades, India has implemented several nutrition programs to address malnutrition in the under-fives. To understand the program's impact, this study assesses the prevalence of acute malnutrition, MAM, and SAM, using MUAC among tribal children.

Design:

The survey was conducted in two tribal blocks (Desaiganj and Bhamragad) of the Gadchiroli district in Maharashtra to identify children registered in the 'Anganwadi' program.

Setting:

A community-based cross-sectional survey was carried out.

Subjects:

The total sample size was 1055 children (aged 0-59 months).

Results:

The overall prevalence of SAM and MAM was 1.4% (n=15) and 9.8% (n=103). A higher prevalence of MAM was found in males (38.5%, n= 40) and females (27.1%, n= 28) in below 6 months. Additionally, a higher prevalence of MAM was observed in females (10.7%, n=113) compared to males (9.0%, n=95). The prevalence of SAM was significantly ($p<0.001$) higher in females (1.7%, n=18) than in males (1.0%, n=11). Children aged between 12 and 17 months were sixteen times more likely (OR=16.9, $p<0.001$, CIs=4.8-59.6) to have MAM (MUAC<12.5cm) than children aged between 6 and 11 months. Children from the Desaiganj block were significantly less likely (OR=0.4, $p<0.001$, CIs= 0.2-0.7) to have MAM compared to children from Bhamragad. Approximately 4% (n=42) of children were classified as critically malnourished.

Conclusion:

There is an urgent need for block-level monitoring of MAM and SAM, as well as evaluation of existing nutrition programs, to address the disparity in the sex-specific prevalence of MAM and SAM in tribal areas.

Introduction

Child malnutrition is a persistent issue and a major contributor to disease burden in India⁽¹⁾. An estimated 7.3% (50 million) of all children under five suffer from wasting at any given time⁽²⁾. Wasting (low weight for height) is one of the basic indicators for assessing the severity of the health status of children. Acute malnutrition is a form of undernutrition caused by decreased food consumption and illness that results in sudden weight loss⁽³⁾. The Joint Child Malnutrition Estimates (JME) reveal insufficient progress to reach the 2025 World Health Assembly (WHA) global nutrition targets and Sustainable Development Goal (SDG) target 2.2. An assessment of progress towards the wasting target is not possible for nearly half of countries and a very high prevalence of under-five child wasting is considered a serious threat to child mortality in India⁽⁴⁾.

According to new data, child malnutrition might be worsening. Though fewer children in India are dying, and those who survive are more malnourished and anaemic in many states⁽⁵⁾. The crisis of child malnutrition in India has often been attributed to historical antecedents such as poverty, inequality, and food shortage. However, countries with similar historical and societal makeup and comparable per capita income have fared much better. Further, the immediate impacts of COVID-19 left several children acutely malnourished, as key health, Nutrition and other life-saving services became less accessible⁽³⁾.

An easy, accurate, and low-cost indicator is helpful for early identification of children with acute malnutrition. Therefore, weight-for-height z scores (WHZ)^(3, 6) and mid-upper arm circumference or MUAC (as subcutaneous fat and muscle mass decrease in undernourished children) have been used to screen for acute malnutrition^(7, 8). 'Acute malnutrition'⁽⁹⁾ can be divided into moderate acute malnutrition (MAM) with a WHZ between -2 and -3 Z-scores (SD) or a MUAC between 115mm and 125 mm., and severe acute malnutrition (SAM), being defined as a weight-for-height Z-score (WHZ) <-3SD or a MUAC of <115 mm^(10, 11). WHZ has been used for years in clinical settings for diagnosing SAM⁽¹²⁾. The use of MUAC was introduced with the development of community-based management of SAM⁽¹³⁾. The WHO standards for MUAC for age show that in a well-nourished population, there are very few children aged 6–69 months with a MUAC of less than 115 mm, but in an underprivileged population the percentage of SAM (with a MUAC less

than 115 mm) in children aged 6-59 months climb up ⁽¹⁰⁾. Though MUAC or WAZ cut-offs vary a lot in different populations, even within the countries a study ^(14, 15,16) shows that the combined case definition (WAZ<-3SD and MUAC<115mm) is found more effective in detecting the severity of malnutrition and preventing child mortality than considering WAZ<-3SD or MUAC<115mm alone. MUAC is closely related to the risk of dying and is easy to implement at the community level after minimum training by health workers or even by volunteers ⁽¹⁷⁾. MUAC and WHZ, however, do not identify the same set of children as having malnutrition and using only one of the diagnostic criteria proposed by WHO may potentially leave some high-risk children untreated ⁽¹⁸⁾. As they are additive and not complementary, it would be pragmatic to retain both criteria for admission to treatment programs ⁽¹⁴⁾.

Though the prevalence of undernutrition (i.e. stunting, wasting and underweight) among tribal children under five years in India has reduced from 43.8%, 27.4% and 45.3% in NFHS-4 to 40.9%, 23.2% and 39.5% in NFHS-5 respectively, the prevalence of malnutrition in under-five children is a grievous concern. ^(19, 20). As per the 2011 Census, 10.5 million tribal people are living in Maharashtra and Gadchiroli district has a 38.71% Scheduled Tribe (ST) population which is the highest in the Nagpur division ⁽²¹⁾. Within Maharashtra, all parameters of undernutrition i.e., stunting (41.4%), wasting (32%) and underweight (46.5%), even in their severity (18.3%, 13.5% and 20.2% respectively), are highest among the tribal children ^(19, 20).

Tribal people are the most underprivileged in terms of their hardship to thrive for better education and skill enhancement and the adversity they experience in income opportunities available and accessible to them, which hinders their overall development. Uncertain modes of employment and economic instability lead to poverty, which makes them victims of morbidity and malnutrition ^(21,22). There is wide acknowledgement that excess morbidity and poor nutritional status in childhood for tribes is partly due to poverty, and food insecurity and partly due to poor access to healthcare services ⁽²²⁾. The nutritional status of the children of the Indigenous tribe, in India as well as in the world, has not received much attention, and there is a lack of focused surveys to obtain data on their food habits and nutritional status. Hence, there is a large scope to explore the nutritional status and the underlying causes of malnutrition among tribal children. The present study contributes to understanding the

status of undernutrition among under-five children belonging to the indigenous tribal population in the Indian context.

The ‘APJ Abdul Kalam Amrut Yojana’ was launched in 2015 by the ‘Tribal Development Department (TDD)’ of the Government of Maharashtra (GOM) aims to provide at least one nutritious meal to pregnant and lactating women at the ‘Anganwadi Centre’ (courtyard shelter). Under the scheme, every tribal woman in her third trimester of pregnancy is entitled to a free nutritious hot meal daily continuing for three months post-delivery so that the newborn is healthy. However, to the best of our knowledge, there has not been a single study to check the nutritional scenario, whether it has improved or not in the tribal blocks of Gadchiroli district. Against this backdrop, the present study has been undertaken with a specific objective to assess the current scenario of under-nutrition among children below five years of age in Desaiganj (formerly known as Wadasa) and Bhamragad blocks, which are predominantly a tribal area in Gadchiroli district (21). Hence, the objective of the present study is to assess the current prevalence of acute malnutrition in terms of MAM and SAM using MUAC among children aged 0-5 years in the Gadchiroli district of Maharashtra.

Methods

Study area

Gadchiroli district, located in the north-eastern side of Maharashtra, shares borders with Telangana and Chhattisgarh. The district is classified as tribal and underdeveloped with most of the land covered by forest and hills, accounting for about 76% of its geographical area. The population of Scheduled Castes (SC) and Scheduled Tribes (ST) in the district is 11.25% and 38.7% respectively (21). The survey was conducted in two purposively selected tribal blocks, Desaiganj and Bhamragad blocks in the Gadchiroli district.

Study Design and Sampling

A community-based cross-sectional survey was conducted in six districts of Vidarbha region, Maharashtra by the Nutrition Bureau, Public Health Department, from December 2021 and March 2022. The aim was to assess the nutritional status and dietary intake of children under five, pregnant women and lactating mothers registered with the Integrated Child Development

Services (ICDS)¹ through Anganwadi centres. Detailed information on pregnant women, lactating mothers (within one year of delivery), and children (0-5 years) was collected from the respective Anganwadi centres through house-to-house visits. The present study focuses on the analysis of children's anthropometric measurements from this nutrition survey. A stratified simple random sampling method was used in the study. Gadchiroli, being a rural district with a 38.7% tribal population, was divided into two categories based on the 2011 Primary Census: urban population and 100% rural population. Blocks with 100% rural population were further categorized based on the number of tribal populations. From each group, two blocks were randomly selected: Desaiganj (<5000 ST households) and Bhamragad (>5000 ST households). Inhabited villages in the selected blocks were grouped by population size (below 500, 500 to 1000, above 1000) and randomly selected for inclusion in the survey. A total of 26 villages were covered, with 1055 children under five years of age registered at the Anganwadi centres included in the study. See Figure 1 for the sample exclusion criteria.

Sample

The total sample considered for analysis was 1055 children (0-59 months) from the Anganwadi centers of the selected villages in the survey period (figure 2).

Anthropometric Investigation

The procedure followed in taking anthropometric measurements as per the United Nations and using the World Health Organization (WHO) child growth charts/standards. The Z score values (WAZ, HAZ and WHZ) were classified by the WHO (6). Anthropometric measurements such as height and weight of mothers and their children (0-5 years) in addition to mid-upper arm circumference were measured using standardized equipment and procedures, i.e. MUACs were measured using the WHO method with two independent observers' methods, with both repeating if the measures were not within a pre-specified limit of agreement.

Height/length Measurement: The length of the children was measured using an 'infantometer', with a headboard and sliding foot piece, placed on a stable level. Diapers, shoes, socks, braids, and hair ornaments were removed. The length/height of the children

¹ Integrated Child Development Services (ICDS) is one of the most inclusive programmes for addressing child malnutrition and child well-being and development, implemented in 1975 by the Ministry of Women and Child Development, Government of India and supported by the UNICEF [4]. ICDS is operating through Anganwadi Centres (AWCs). An AWC is operated by an Anganwadi Worker (AWW) and an Anganwadi Helper (AWH).

under two years was measured by lying them and for the older children measured by standing. During measurement, hands were kept on the shoulder/knees and pressed gently without hurting the child. The soles of the feet were flat on the foot piece pointing up. Measurements were taken to the nearest 0.1 cm.

A *stadiometer* was used to measure standing height with a vertical backboard, a fixed baseboard, and a movable headboard placed on a level floor. During measurement socks, shoes, hair ornaments and braids were removed. One person kneeled near the child's foot and helped the child to stand with the back of the head, shoulder blades, buttocks, calves, and heels touching the vertical boards. At the same time with the other hand, the headboard was pulled firmly on the top of the head to compress hair. Measurements were taken to the nearest 0.1 cm.

Weight Measurement: The weighing scale, used to measure the child's weight, was a 'Spring Balance', which was solidly built, durable, electronic (Digital), able to measure with a precision of 0.01 kg(10g) and allowed taring (the act of zeroing the instrument). Hence, there was no need to subtract weight, reducing the risk of error. First, the child kept calm by holding him/her in the mother's arms. A beam scale or a hanging scale (Salter type) was used where a tared machine was not available. Before taking weight, the scale was adjusted to zero. Time was given for the child to settle, and the weight to stabilize.

Mid Upper Arm Circumference (MUAC) measurement was done with a tri-colored tape known as Shakir's tape (Annexure 1.1). MUAC was taken for children aged 6-59 months only. The arm circumference was measured on the upper left arm by flexing the child's elbow to 90 degrees. Then, the midpoint between the acromion process of the shoulder joint and the tip (olecranon) of the elbow was marked and then the arm was allowed to hang freely. At the same time, the measuring tape was placed snugly around the arm at the midpoint mark. The tape was not pulled too tightly during the measurement.

Data Collection

Data was collected using a pretested semi-structured interview schedule through personal interviews. A single interviewer collected the data by conducting face-to-face interviews with the mothers of eligible children. All those who fulfilled the inclusion criteria were eligible to participate in the study.

Ethical Consideration

Ethical approval to conduct the study was obtained from the Institutional Ethics Committee (IEC) of the State Institute of Health & Family Welfare (SIHFW), Nagpur. Mothers of the children selected for the study were informed about the study's purpose. Informed consent was obtained from the mothers before collecting information about their children, and confidentiality of the collected information was assured. Complete anonymity was maintained during data collection and analysis.

Variables

Independent variables include the *child's age* (categories below 6, 6-11, 12-17, 18-23, 24-35, 36-47 and 48-59 months), *child's sex* (male and female), *blocks* (Bhamragad and Desaignanj) and *wasting* (WHZ<-2SD, yes and no) used in logistic regression analysis.

Dependent variables include *Moderate Acute Malnutrition or MAM* (MUAC<12.5cm, yes and no), *Severe Acute Malnutrition or SAM* (MUAC<11.5cm, yes and no), and *Severe Wasting* (WHZ<-3SD, yes and no).

Statistical Analyses

Simple bivariate and multivariate analyses were carried out to analyze the data. Bivariate logistic regression analysis was done to assess the effect of independent variables (age, sex, block and wasting) on the prevalence of moderate acute malnutrition. The significance level of the association between the variables was shown by Pearson's correlation coefficient and Chi-square values.

The Z score value (i.e., Standard Deviation or SD) for the anthropometric measures in terms of height for age, weight for age and height for weight were calculated using the WHO Anthro, version 3.2.2. Computation of all other variables and all statistical analyses were carried out by IBMSPSS Statistics, version 29.

Findings

Characteristics of the study population

Table 1A represents the characteristics of the sample (0-59 months). Approximately 10% (n=103) of children are under six months old, while about 33% (n=347) are between 6 and 24 months old. The mean age of the sample is 28.6 months. The percentage of male children (50.5%) is almost equal to the percentage of female children (49.5%). Around 39%, (n=408) of the sample is from the Bhamragad block, while the majority (61%, n=647) is from the Desaiganj block. The mean MUAC, height and weight of the sample are 13.6 cm, 80.9cm and 9.9 kg respectively. The median age is 28 months.

The proportion of children with SAM is 1.4% (n=15), whereas the proportion of children with MAM is 9.8% (n=103) (Table 1B).

Prevalence of malnutrition among children between 0-59 months

Table 2 shows the prevalence of moderate acute malnutrition (MAM) and severe acute malnutrition (SAM) among children aged 0-59 months based on their age group and sex in the selected blocks of the Gadchiroli district.

The results reveal that the malnutrition prevalence is 11.2% in the study population, with MAM at 9.8% and SAM at 1.4%. MAM is highest (33.0%) in children under six months and decreases with age. The lowest MAM prevalence is in the age group above 47 months. Sex and age-specific differences in MAM prevalence are noted with significant differences across all age groups. Higher MAM prevalence is found in males (38.5%) and females (27.1%) in the youngest age groups. Overall, girls (12.3%) have a higher MAM prevalence than boys (10.1%). Severe acute malnutrition (SAM) prevalence is 1.4%, higher in females (1.7%) than males (1.0%). Children under six months show the highest SAM prevalence (8%), decreasing with age group. The overall acute malnutrition (MAM+SAM) is 11.2%, the highest in children under six months (41%). A significant difference ($p<0.001$) in MAM and SAM prevalence by sex is observed.

Figure 3 illustrates age and sex-specific MAM prevalence among children aged 0 to 5 years. While MAM prevalence decreases with age, girls show higher moderate malnutrition rates than boys in subsequent age groups.

Prevalence of MAM and SAM in Selected blocks

Figure 4 shows moderate acute malnutrition prevalence among children aged 0-5 years with MUAC<12.5 cm in selected blocks (Bhamragad and Desaiganj) in Gadchiroli district. MAM prevalence decreases with age in Bhamragad compared to Desaiganj. Both blocks have the highest prevalence below one year (40.7% in Bhamragad, 19.6% in Desaiganj) and lowest above 4 years (2.5% in Bhamragad, 0.9% in Desaiganj). Girls have higher MAM prevalence, especially in the age group of two to four years.

The figure shows that the prevalence of MAM has steadily decreased with the increase in the age group of the children in Bhamragad compared to the prevalence of MAM in children of Desaiganj block. Both blocks have observed the highest prevalence in the age group below one year (40.7% and 19.6% in Bhamragad and Desaiganj respectively) and lowest in the children aged above 4 years (2.5% and 0.9%). In both blocks, girls are showing a higher prevalence of MAM particularly in children aged between two to four years.

Factors associated with MAM among children (6-59 months)

Table 3 represents the result from bivariate logistic regression analysis in terms of the effect of age, sex, block and wasted (WHZ<-2SD) status of children 6-59 months on the prevalence of moderate acute malnutrition or MAM (MUAC<12.5cm). Results show that there is a significant decrease in the prevalence of MAM in the higher age group. Children in the age group 12 to 17 months are sixteen times more likely (OR=16.885, $p<0.001$) to have MAM (MUAC<12.5cm) than the reference category children (i.e. children aged 6-11 months).

Female children (OR=1.45) are more likely to have MAM than male children. Children from the Desaiganj block (OR=0.423, $p<0.001$) are significantly less likely to have MAM than the children from the Bhamragad block. Children who are moderately wasted (WHZ<-2SD) are three times (OR=3.510, $p<0.001$) more likely to have MUAC<12.5cm than the children who are not wasted.

Prevalence of MAM and SAM concerning MUAC, and WHZ in children (6-59 months)

Figure 5 represents the percentage of children aged between 6 to 59 months who have moderate acute malnutrition (MAM) in terms of having both MUAC less than 12.5cm and WHZ less than -2SD. About 17% and 4.2% of children are found moderately malnourished for MUAC(<12.5cm) and wasting (WHZ<2SD). It is also found that about 3.9% of children are critically malnourished as having both MUAC<12.5cm and WHZ<2SD.

Figure 6 represents the percentage of children who are severely acutely malnourished (SAM) for both parameters, i.e., MUAC<11.5cm and WHZ<3SD. About 6.2 percent (WHZ<- 3SD) and 0.5 per cent (MUAC<11.5cm) of children are found to be severely acutely malnourished. The percentage of children having both WHZ<-3SD and MUAC<11.5cm is 0.2 per cent, who are considered critically ill and need special treatment.

Discussion

The study found that acute malnutrition was more prevalent among female children aged 0 to 5 years compared to male children. Disparities in child malnutrition prevalence were observed at the block level, even with similar nutrition programs in place. Children in the Bhamragad block had a higher prevalence of acute malnutrition compared to those in the Desaiganj block with female children more affected than male children (23). Regression analysis indicated significant differences in malnutrition prevalence based on age and sex (24) in tribal blocks. A higher percentage of children with MUAC<12.5 cm was wasted compared to non-wasted children. The present research observed about 4% of children in the study area with both WHZ<2SD and MUAC<12.5cm who need to be considered critically ill. Further, about 7% of children who are severely malnourished have a higher risk of morbidity and mortality (22). The study highlighted the higher prevalence of acute malnutrition in children under two years old (7, 25).

The prevalence of acute malnutrition among children under 5 years was 11.2%, with 1.4% experiencing severe acute malnutrition. Children with severe acute malnutrition are at significantly higher risk of morbidity and mortality. The study emphasized the need for urgent

policy attention in districts like Gadchiroli to address the higher prevalence of severe acute malnutrition. Despite some improvements, child malnutrition rates in Maharashtra remained largely unchanged with a higher prevalence of wasting among children under two years in Gadchiroli compared to the state average. According to NFHS-5 the percentage of children (below 2 years) who are underweight (36%) or wasted (26%) has not changed since NFHS-4^(19, 20). Though the prevalence of stunting (35.7%) and underweight (35.4%) among children (below 2 years) in Gadchiroli is like that of the state average, NFHS-5 has found a higher prevalence (30.0%) of wasting (WHZ<2SD) recorded among children below two years for Gadchiroli than the state average (25.6%).

The study identified gender-based disparities in child malnutrition prevalence and the attributing factors be as breastfeeding duration and intra-household food allocation to the differences^(26, 27). Girls were found to have less dietary diversity⁽²⁸⁾ and access to food compared to boys, reflecting gender preferences in families⁽²⁹⁾. Maternal education, poverty and parental attitudes also influence the nutritional status of female children⁽²⁶⁾.

Government Programs like POSHAN Abhiyan (National Nutrition Mission), Anganwadi Services, Scheme for Adolescent Girls, under Mission Poshan 2.0 and Pradhan Mantri Matru Vandana Yojana (PMMVY) under Mission Shakti, aim to address malnutrition issues, especially in tribal areas. Though the Integrated Child Development Services-Common Application Software (ICDS-CAS) was created to digitise the records and create a real-time monitoring system for the beneficiaries under the POSHAN Abhiyaan, however, until 2019, only 611369 Anganwadi workers (AWW) were equipped with ICDS-CAS⁽³⁰⁾. Further, the COVID-19 period experienced severe disruption in the transportation and food supply chain, huge shortages and curtailing of employment opportunities which resulted in worsening the situation of economic crisis among the marginalized and poor population. Thus, the pandemic has resulted in reduced food expenditure and household food insecurity among daily wage earners and farmers^(31, 32).

Though malnourishment in tribal children showed a declining trend as the prevalence of stunting, wasting and underweight was reduced from 43.8%, 27.4% and 45.3% respectively in NFHS-4 to 40.9%, 23.2% and 39.5% respectively under NFHS-5, manifold increase in the prevalence of SAM in several Indian districts including in the tribal belts is a public health emergency that requires urgent policy response⁽³¹⁾. A study indicated that MUAC is better

than WHZ at identifying high-risk children in the community (33). It is also evident that in community-based surveys, both the parameters, i.e., WHZ and MUAC don't cover the same set of population samples (34, 35). Hence, there is a need to develop stand-alone parameters, unlike WHZ or MUAC which ascertain a subset of a population, that can identify all malnourished children (36). The Government of India's 'Poshan Tracker', the largest mobile phone-based nutrition surveillance system in the world, provides transparent data on anthropometric outcomes, the functioning of Anganwadi Centres (AWCs) and the receipt of care services (37). 'Protocol for Management of Malnourished Children' has been drafted by MWCD with inputs from MH&FW (38).

Most interventions suggest the treatment of acute malnutrition (SAM) and indicate preventive strategies for chronic malnutrition, as facility-based treatment of acute malnutrition has very limited coverage and incurs more cost (39). At the community level, to manage acute malnutrition, understanding the new guidelines (40) for the therapy of chronic malnutrition and its execution by frontline healthcare providers is necessary. The approach requires many trained staff and a substantial inpatient bed capacity. Besides available facilities, sufficient attention needs to be paid to the quality of care (41).

In India, there is a pressing need for enhanced data collection and monitoring systems to effectively address malnutrition. Additionally, increased investment in infrastructure development is necessary to ensure the timely delivery of vaccines, medicines, and hospital facilities (42). The importance of community-level interventions and trained healthcare providers to combat acute malnutrition is also underscored (43). The present study highlights the importance of closer collaboration between various ministries and departments to achieve this goal. The study highlighted gaps in programmatic data for identifying and treating children with severe acute malnutrition among the Indigenous population in India. Further, the present research contributes to understanding the gender gap in child malnutrition in the tribal area. Hence, future studies may address this lacuna in public health research targeting the health and nutritional status of children in tribal populations around the world.

Strengths and Limitations

India's economy shrank for the first time during the NFHS 5 data collection period due to the nationwide lockdowns from the first wave of the COVID-19 pandemic. This led to disruptions in many nutrition-sensitive programs, potentially impacting the prevalence of undernutrition.

To address this the current study advocates computing the prevalence of acute malnutrition using both $WHZ < 2SD$ and $MUAC < 12.5cm$ parameters to capture all forms of malnourishment. Furthermore, the survey did not collect information on edema in children under five years which is a limitation of the data.

However, the study focused on the health and nutrition status of children in Anganwadi centres in the selected villages. Household characteristics and dietary information were collected separately, limiting the ability to establish connections between women, children, and their household backgrounds. Long-term morbidity or mortality data for children under five years were not collected in the survey.

Conclusions and Recommendation

The block-level disparity in the prevalence of malnutrition among children under five years in the Gadchiroli district indicates the need for targeted interventions. Monitoring and evaluation of nutrition programs at the block level are crucial to addressing these disparities. Training of health workers to identify and manage malnourished children is essential. Children with critical conditions require specialized nutrition plans and prompt intervention. Besides, AWWs and health workers need to ensure the use of available growth tracker devices and to sensitize the mothers about the health risks associated with malnutrition. Based on the clinical assessment, therapeutic food needs to be provided to the targeted children. Therefore, feeding practices for infants and young children play a crucial role in preventing malnutrition-related illness.

Regular assessment and monitoring of nutrition programs at the block level, with data reporting is essential. Qualitative studies can provide valuable insights into nutritional requirements in program areas. Ensuring proper guidelines, followed by frontline health workers and officials, is crucial to address under-nutrition effectively. Identifying and treating malnourished children promptly and providing appropriate interventions are key recommendations for improving child nutrition outcomes.

References

1. Swaminathan S, Rasaily S, Bansal PG, *et al.* (2019) The burden of child and maternal malnutrition and trend in its indicators in the states of India: The Global Burden of Disease Study 1990-2017. *India State-Level Disease Burden Initiative Malnutrition Collaborators. The Lancet, Child & Adolescent Health*, 3(12) 855-870. [https://doi.org/10.1016/S2352-4642\(19\)30273-1](https://doi.org/10.1016/S2352-4642(19)30273-1)
2. World Health Organisation (2020) *Global action plan on child wasting: a framework for action to accelerate progress in preventing and managing child wasting and the achievement of the Sustainable Development Goals*. WHO, Geneva.
3. World Health Organization (2023) *Urgent action is needed as acute malnutrition threatens the lives of millions of vulnerable children*. WHO, Geneva.
4. United Nations Children's Fund (UNICEF), World Health Organization (WHO), International Bank for Reconstruction and Development/The World Bank. (2023) *Levels and trends in child malnutrition: UNICEF / WHO / World Bank Group Joint Child Malnutrition Estimates: Key findings of the 2023 edition*. New York: UNICEF and WHO. CC BY-NC-SA 3.0 IGO.
5. Chatterjee P. (2021) India's child malnutrition story worsens. *The Lancet. Child & Adolescent Health*. 2021 5(5): 319-320. Doi [https://doi.org/10.1016/S2352-4642\(21\)00064-X](https://doi.org/10.1016/S2352-4642(21)00064-X)
6. World Health Organization (2006) *Child Growth Standards: Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index-for-Age: Methods and Development*. WHO, Geneva. Available at: http://www.who.int/childgrowth/standards/technical_report/en/index.html
7. Taneja S, Rongsen-Chandola T, Mohan SB, *et al.* (2018) Mid-upper arm circumference as a predictor of risk of mortality in children in a low-resource setting in India. *PLoS ONE*13(6):e0197832. <https://doi.org/10.1371/journal.pone.0197832>
8. Myatt M, Khara T and Collins S. (2006) A Review of Methods to Detect Cases of Severely Malnourished Children in the Community for Their Admission into Community-Based Therapeutic Care Programs. *Food and Nutrition Bulletin*. 2006;27(3_suppl3): S7-S23. doi:10.1177/15648265060273S302
9. Note on Terminology of “Wasting” and “Acute malnutrition” (2023) World Health Organization.

https://fscluster.org/sites/default/files/documents/note_on_terminology_acute_malnutrition_and_wasting_who_feb_2023.pdf

10. WHO , UNICEF. (2009) *Child growth standards and the identification of severe acute malnutrition in infants and children. A joint Statement by the World Health Organization and the United Nations Children's Fund.*
11. Aguayo VM, Aneja S, Badgaiyan N, Singh K. (2015) Mid upper-arm circumference is an effective tool to identify infants and young children with severe acute malnutrition in India. *Public Health Nutr.* Dec;18(17):3244-8. doi: 10.1017/S1368980015000543. Epub 2015 Mar 11. PMID: 25757562; PMCID: PMC10271291.
12. World Health Organization (1999). *Management of severe malnutrition: a manual for physicians and other senior health workers.* WHO, Geneva. Available at: <https://www.who.int/publications/i/item/9241545119>
13. WHO, UNICEF, WFP, SCN. (2007). *Joint statement. Community-based management of severe acute malnutrition.* Geneva, New York, Rome, 2007. Available at: http://www.who.int/child_adolescent_health/documents/pdfs/severe_acute_malnutrition_en.pdf
14. Grellety E, Golden MH. Weight-for-height and mid-upper-arm circumference should be used independently to diagnose acute malnutrition: policy implications. *BMC Nutr* **2**, 10 (2016). <https://doi.org/10.1186/s40795-016-0049-7>
15. Briend A, Alvarez JL, Avril N. *et al.* Low mid-upper arm circumference identifies children with a high risk of death who should be the priority target for treatment. *BMC Nutr* **2**, 63 (2016). <https://doi.org/10.1186/s40795-016-0101-7>
16. Myatt M, Khara T, Collins S. (2006) A Review of Methods to Detect Cases of Severely Malnourished Children in the Community for Their Admission into Community-Based Therapeutic Care Programs. *Food and Nutrition Bulletin.* 27(3_suppl3): S7-S23. doi:10.1177/15648265060273S302
17. Shukla N, Toppo NA, Thakur A, *et al.* (2020) Mid-upper arm circumference versus weight for height Z-score for detecting malnutrition in children aged 06-59 months in rural and urban areas of Jabalpur district, Madhya Pradesh, India. *Int J Community Med Public Health* 2020; **7:1794-8.**
18. Berkley J, Mwangi I, Griffiths K, *et al.* (2005) Assessment of severe malnutrition among hospitalized children in rural Kenya: comparison of weight for

- height and mid-upper arm circumference. *JAMA*. 3;294(5):591-7. doi: 10.1001/jama.294.5.591. PMID: 16077053.
19. NFHS-4. National family health survey. Government of India: Ministry of Health and Family Welfare, 2015.
 20. NFHS-5. National family health survey. Government of India: Ministry of Health and Family Welfare, 2020.
 21. 2011 Census of India. Registrar General and Census Commissioner of India.
 22. Dani V, Satav K, Pendharkar J *et al.* (2017) Community-based management of severe malnutrition: SAM and SUW in the tribal area of Melghat, Maharashtra, India. *Clinical Epidemiology and Global Health* 5,62–69
 23. Kumar S, Kawalia S, Thitame SN *et al.* (2018) Malnutrition among under-five tribal children with a special focus on dietary intake in Akole block of Western Ghat, Maharashtra, India. *Int J Med Sci Public Health*, 7(3):165-169.
 24. Raut PK, Chaturvedi RM (2016) A Cross-sectional Study to Assess the Nutritional Status of Under Five Children in Tribal Area of Western Maharashtra, India. *International Journal of Current Medical and Applied Sciences*, May, 10(3),119-123.
 25. Jima BR, Hassen HY, Getnet Y *et al.* (2021) Diagnostic performance of mid upper arm circumference for detecting severe wasting among infants aged 1–6 months in Ethiopia. *Am J Clin Nutr* 2021; 113:55–62.
 26. Bhutia, D T (2014) Protein Energy Malnutrition in India: The Plight of Our Under Five Children. *Journal of Family Medicine and Primary Care* 3(1): p 63-67, Jan–Mar 2014. DOI: 10.4103/2249-4863.130279
 27. Soni A, Fahey N, Bhutta ZA *et al.* (2021) Early childhood undernutrition, preadolescent physical growth, and cognitive achievement in India: A population-based Cohort study. *PLoS Med* 18(10): e1003838.
<https://doi.org/10.1371/journal.pmed.1003838>
 28. Aurino E. (2016) Do Boys Eat Better Than Girls in India? Longitudinal Evidence from Young Lives. Working paper 146, Young Lives. Oxford Department of International Development (ODID), *University of Oxford*. ISBN 978-1-9099403-60-4. https://www.younglives.org.uk/sites/default/files/migrated/YL-WP146-Aurino_Do-Boys-Eat-Better-than-Girls.pdf

29. Thakur R, Gautam RK (2015). Nutritional status among boys and girls of a central Indian Town (Sagar). *Anthropological Review*, 78(2),197–212. <https://doi.org/10.1515/anre-2015-0014>
30. Niti Aayog. (2020) Government of India. Accelerating progress on nutrition in India: what will it take? Third Progress Report, July 2020. (Accessed on October 9, 2023). <https://www.niti.gov.in/sites/default/files/2020-10/AbhiyaanMonitoringReport.pdf>
31. Alvi M, Gupta M (2020) Learning in times of lockdown: how COVID-19 is affecting education and food security in India. *Food Secur* 2020;12:793–6.
32. Varghese JS, Gupta A, Mehta R *et al.* (2022) Changes in child undernutrition and overweight in India from 2006 to 2019: an ecological analysis of 36 states. *GlobHealthSciPract*,10(5):e2100569. <https://doi.org/10.9745/GHSP-D-21-00569>
33. Briend A, Maire B, Fontaine O *et al.* (2012) Mid-upper arm circumference and weight-for-height to identify high-risk malnourished under-five children. *Blackwell PublishingLtd Maternal and Child Nutrition*, 8(130–133). DOI: 10.1111/j.1740-8709.2011.00340.x
34. Fiorentino M, Sophonneary P, Laillou A *et al.* (2016). Current MUAC Cut-Offs to Screen for Acute Malnutrition Need to Be Adapted to Gender and Age: The Example of Cambodia. *PLoS ONE* 11(2): e0146442. doi:10.1371/journal.pone.0146442
35. Zaba T, Nyawo M, Morán JLA (2020) Does weight-for-height and mid-upper arm circumference diagnose the same children as wasted? An analysis using survey data from 2017 to 2019 in Mozambique. *Archives of Public Health*, 78:94 <https://doi.org/10.1186/s13690-020-00462-7>
36. John C, Ocheke IE, Diala U *et al.* (2017) Does mid upper arm circumference identify all acute malnourished 6–59 month old children, in field and clinical settings in Nigeria?, *South African Journal of Clinical Nutrition*, 30:3, 55-59, DOI: [10.1080/16070658.2016.1255486](https://doi.org/10.1080/16070658.2016.1255486)
37. Jaacks LM, Awasthi A, Kalra A. (2024) India’s Poshan Tracker: data-driven tool for maternal and child nutrition. Comment, March 10, 2024. (Accessed on March 13, 2024) DOI: <https://doi.org/10.1016/j.lansea.2024.100381>
38. National event for Launch of ‘Protocol for Management of Malnutrition in Children’. (2023). Press Release, GOI. (accessed on February 12, 2024) <https://pib.gov.in/PressReleaseIframePage.aspx?PRID=1965940>

39. World Health Organization. (2013) Guideline update: technical aspects of the management of severe acute malnutrition in infants and children. WHO, Geneva. https://scholar.google.com/scholar_lookup?title=Guideline%20update%3A%20Technical%20aspects%20of%20the%20management%20of%20severe%20acute%20malnutrition%20in%20infants%20and%20children&publication_year=2013 [Accessed 13 October 2023].
40. WHO guideline on the prevention and management of wasting and nutritional oedema (acute malnutrition) in infants and children under 5 year. (2024) March. <https://app.magicapp.org/#/guideline/noPQkE/section/n32P8W>
41. Collins S, Dent N, Binns P *et al.* (2006) Management of severe acute malnutrition in children. *Lancet*. Dec 2;368 (9551):1992-2000. doi: 10.1016/S0140-6736(06)69443-9. PMID: 17141707
42. Saha S, Singh R. (2021) Child malnutrition in India: A systemic failure. Down to Earth. Published: Thursday, 15 April, 2021. <https://www.downtoearth.org.in/blog/health/child-malnutrition-in-india-a-systemic-failure-76507>
43. Protocol for management of malnutrition in Children. 2023. POSHAN Abhiyaan. Ministry of Health and Family Welfare & Ministry of Women and Child Health Department, Government of India. <https://wcd.nic.in/sites/default/files/CMAM-Book-10-10-2023%20%281%29.pdf>

Table 1A Characteristics of children (0-59months), Gadchiroli, Maharashtra, India, 2022

Characteristics	Percentage	Children(N)
Age group (Months)		
<6	9.8	103
6 to11	11.2	118
12 to17	12.5	132
18 to 23	9.2	97
24 to 35	19	200
36 to 47	19.4	205
48 to 59	19	200
Sex		
Male	50.5	533
Female	49.5	522
Block		
Bhamragad	38.7	408
Desaiganj	61.3	647
Total	100	1055
Mean Age (months)		
SD		17.04
Median		28
IQR		30
Mean MUAC (cm)		
SD		1.18
Mean Height (cm)		
SD		12.62
Mean Weight (kg)		
SD		2.78

Table 1B Nutritional Status of Children (0-59 months) Gadchiroli, Maharashtra, India

Malnutrition type	Percentage	Children
SAM	1.4	15
MAM	9.8	103
Normal	88.8	928
Total	100	1055

Table 2. Percentage of children (0-59 months) with moderate acute malnutrition (MAM, WHZ<2SD) and severe acute malnutrition (SAM <3SD) by age and sex in Gadchiroli, Maharashtra, India, 2022

Age Group (months)	MAM (%)			SAM (%)			Acute Malnutrition = MAM+SAM	χ^2 Value Sex	Children (N)
	Male	Female	Total	Male	Female	Total			
<6	38.5	27.1	33.0	3.8	12.5	8.0	41.0	150.302, p<0.001	103
6 to 11	17.2	11.9	14.5	3.4	1.7	2.6	17.1		118
12 to 17	8.6	16.4	12.2	1.4	1.6	1.5	13.7		132
18 to 23	12.2	22.9	17.5	0	0	0	17.5		97
24 to 35	4.3	11.3	8.0	0	0	0	8.0		200
36 to 47	0.9	2.1	1.5	0	0	0	1.5		205
48 to 59	1.0	0	0.5	1.0	1.0	1.0	0.5		200
Total	9.0	10.7	9.8	1.0	1.7	1.4	11.2	1055	

Table 3. Bivariate Logistic Regression Analysis showing Prevalence of MAM among Children (6-59 months) by their Background Characteristics, Gadchiroli, Maharashtra, India, 2022

Characteristics	MAM (MUAC<12.5cm) in percentage		OR (<i>Exp-β</i>)	<i>P-values</i>	95% C.I. for <i>Exp (β)</i>	
					Lower	Upper
Age Group (months)						
6 to 11®	16.9	1				
12 to 17	13.6	16.9	<0.001	4.8	59.6	
18 to 23	17.5	11.4	<0.001	3.2	40.7	
24 to 35	8.0	15.3	<0.001	4.2	54.7	
36 to 47	1.5	7.3	<0.01	2.0	25.9	
48 to 59	1.5	1.1	0.243	0.2	5.3	
Child's sex						
Male®	6.7	1				
Female	9.5	1.4	0.211	0.9	2.4	
Blocks						
Bhamragad®	12.0	1				
Desaiganj	4.9	0.4	<0.001	0.2	0.7	
Wasting (WHZ<- 2SD)						
No®	5.3	1				
Yes	18.5	3.5	<0.001	2.1	5.9	
Total	8.1					

Note: Reference category is

®.

Figure 1: Flowchart showing Sample Exclusion




Figure 2: Sample Distribution




Figure 3 Prevalence of acute malnutrition (MUAC<12.5cm) in children 0-59 months, Gadchiroli, Maharashtra, India, 2022




Figure 4 Prevalence of acute malnutrition (MUAC<12.5cm) in children 0-59 months, Gadchiroli, Maharashtra, India, 2022




Figure 5 Percentage of children of 6-59 months with MAM (MUAC<12.5cm and WHZ<- 2SD), Gadchiroli, Maharashtra, India, 2022




Figure 6 Percentage of children of 6-59 months with SAM (MUAC<11.5cm and WHZ<-3SD), Gadchiroli, Maharashtra, India, 2022

