

A UNIVERSALITY RESULT FOR ENDOMORPHISM MONOIDS OF SOME ULTRAHOMOGENEOUS STRUCTURES

IGOR DOLINKA AND DRAGAN MAŠULOVIĆ

*Department of Mathematics and Informatics, University of Novi Sad,
Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia
(dockie@dmi.uns.ac.rs; masul@dmi.uns.ac.rs)*

(Received 15 September 2010)

Abstract We devise a fairly general sufficient condition ensuring that the endomorphism monoid of a countably infinite ultrahomogeneous structure (i.e. a Fraïssé limit) embeds all countable semigroups. This approach not only provides us with a framework unifying the previous scattered results in this vein, but actually yields new applications for endomorphism monoids of the (rational) Urysohn space and the countable universal ultrahomogeneous semilattice.

Keywords: endomorphism monoid; Fraïssé limit; pushout

2010 Mathematics subject classification: Primary 20M20
Secondary 03C15; 08A35; 18A30; 20M50

1. Introduction

The principal source of motivation for this paper stems from fascinating objects of model theory called *Fraïssé limits*. Namely, by a well-known result of Fraïssé [12, 13] (see also [16]), if \mathcal{C} is a countable set of finitely generated first-order structures of a fixed countable signature which is (up to isomorphism) closed under taking finitely generated substructures and enjoys the *amalgamation property* (AP) and the *joint embedding property* (JEP), then there is a unique countable structure F with the following two properties:

1. the set of all finitely generated substructures of F coincides (up to isomorphism) with \mathcal{C} ;
2. F is *ultrahomogeneous*, meaning that any isomorphism between its finitely generated substructures extends to an automorphism of F .

Such an F is called the \mathcal{C} -*universal ultrahomogeneous structure* or, following [16], the *Fraïssé limit* of \mathcal{C} . The corresponding class \mathcal{C} with the above properties is called a *Fraïssé class*. Conversely, it is known that every countably infinite ultrahomogeneous structure arises in this way: it is simply the limit of the class of structures isomorphic to its finitely

generated substructures. Of course, if all of this happens within a locally finite class of structures (e.g. relational structures with finitely many constants), then it suffices to replace ‘finitely generated’ by ‘finite’ in the above definitions.

Historically, the two oldest examples of Fraïssé limits are the *rational Urysohn space* $\mathbb{U}_{\mathbb{Q}}$, the limit of the class of all finite metric spaces with rational distances, and the linear order of the rationals, \mathbb{Q} , which is the limit of the class of all finite linear orders. Other well-known classes admitting Fraïssé limits include finite simple graphs, finite posets and finite semilattices, resulting in the random graph R [3, 4], the random (generic) poset \mathbb{P} [9, 23] and the countable universal homogeneous semilattice Ω [10], respectively.

More recently, searching for a broader perspective on the notion of ultrahomogeneity, Cameron and Nešetřil [5] introduced the concept of *homomorphism homogeneity*. A structure A is homomorphism homogeneous if any homomorphism $B \rightarrow A$ defined on a finitely generated substructure B of A can be extended to an endomorphism of A . In particular, any endomorphism of B extends to an endomorphism of A . Now, the somewhat more sophisticated question arises of whether one can select these extensions in a ‘regular’ fashion, in the sense that it is possible to choose, for each $\varphi \in \text{End}(B)$, an extension $\hat{\varphi} \in \text{End}(A)$ such that the mapping $\varphi \mapsto \hat{\varphi}$ is a monoid embedding $\text{End}(B) \rightarrow \text{End}(A)$. The same question applies to an arbitrary substructure B of A .

Here we consider the case when A is a countably infinite ultrahomogeneous structure, that is, a Fraïssé limit. Since A is universal for a certain class of countable structures, it is to be expected that $\text{End}(A)$, the endomorphism monoid of A , will exhibit a very rich structure and to some extent ‘inherit’ the universal properties of A . In fact, if for some (countably) infinite substructure B of A there is an embedding $\text{End}(B) \rightarrow \text{End}(A)$ and B admits any self-map of the set B as an endomorphism, then $\text{Self}(\aleph_0)$, the monoid of all self-maps of a countably infinite set, embeds into $\text{End}(A)$. Since any countable semigroup embeds into $\text{Self}(\aleph_0)$ (by the semigroup analogue of the Cayley Theorem,) it would then follow that the monoid $\text{End}(A)$ is *countably universal*, i.e. that it contains a copy of each countable semigroup.

For example, it was shown in [2] that this is true in the case of the random graph R , while the same conclusion is reached in [6] for the generic poset \mathbb{P} . Furthermore, Uspenskij [25] proved that every topological group with a countable base is isomorphic to a topological subgroup of the isometry (i.e. automorphism) group of the *Urysohn space* \mathbb{U} [18, 24], which implies that every countable group embeds into $\text{Aut}(\mathbb{U}_{\mathbb{Q}})$. Also, for any countably infinite ultrahomogeneous simple graph G (a complete list of these is given in [19]) it is true that the symmetric group $\text{Sym}(\aleph_0)$, and thus any countable group, embeds into $\text{Aut}(G)$ [15]. This is, however, no longer true if we move into the realm of monoids and semigroups: it follows from [21] that R is actually the *only* countable ultrahomogeneous graph whose endomorphism monoid embeds all countable semigroups. Moreover, an easy exercise shows that no non-trivial finite group embeds into $\text{End}(\mathbb{Q})$, the monoid of all order-preserving mappings of \mathbb{Q} .

In this paper, we develop a system of conditions on a Fraïssé class \mathcal{C} under which its limit $F = \text{Flim}(\mathcal{C})$ has the property that $\text{End}(F)$ embeds the endomorphism monoid of any of its substructures, with universality consequences as sketched above. After intro-

ducing the required preliminary notions and constructions, in §3 we state our general method. The concrete applications are presented in §4. First, we review the results of [2,6] in the light of the presented general approach. We also derive some new consequences by proving that both $\text{End}(\mathbb{U}_{\mathbb{Q}})$ and $\text{End}(\Omega)$ contain copies of $\text{Self}(\aleph_0)$. We also discuss some open problems.


2. Preliminaries: pushouts, amalgams, amalgamated sums

For a Fraïssé class \mathcal{C} , let $\bar{\mathcal{C}}$ denote the class of all countable structures, all of whose finitely generated substructures belong to \mathcal{C} . We shall be concerned with Fraïssé classes \mathcal{C} for which $\bar{\mathcal{C}}$, considered as a category of first-order structures, admits certain constructions and properties (see [20] for the basics of category theory). The most fundamental such construction will be that of a *pushout*. Let X, Y, Z be objects of a category \mathcal{C} , along with morphisms $f: X \rightarrow Y$ and $g: X \rightarrow Z$; such a configuration

$$Y \xleftarrow{f} X \xrightarrow{g} Z$$


is referred to as a *span*. The pushout of this span consists of an object P and two morphisms $i_1: Y \rightarrow P$ and $i_2: Z \rightarrow P$ such that:

- (a) the diagram



commutes;

- (b) for any object Q and morphisms $j_1: Y \rightarrow Q$ and $j_2: Z \rightarrow Q$ for which the part of the following diagram involving X, Y, Z, Q is commutative, there exists a unique morphism $u: P \rightarrow Q$ making the whole diagram



commutative.

A standard argument shows that the pushout is unique up to isomorphism. The diagram from (a) will be called a *pushout square* (in \mathcal{C}). Throughout the paper, the composition of morphisms will follow the right-to-left convention, so that if $\alpha: A \rightarrow B$ and $\beta: B \rightarrow C$,

then their composition is $\beta\alpha: A \rightarrow C$. Thus, condition (b) asserts the existence of a unique morphism u such that $ui_1 = j_1$ and $ui_2 = j_2$, where $j_1f = j_2g$.

In the rest of the paper we will work with *concrete* categories, whose objects are first-order structures (of a particular similarity type), and morphisms are homomorphisms of such structures. It is crucial to stress at this point that in the definition of a pushout, the homomorphism u need not be an embedding (or injective) even if all the other morphisms occurring in the previous diagram (that is, f, g, i_1, i_2, j_1, j_2) are embeddings.

Recall that an *amalgam* (in a concrete category) is a span (A, B, C, f, g) such that $f: B \rightarrow A$ and $g: B \rightarrow C$ are embeddings. If $A, B, C \in \mathcal{C}$ for some class \mathcal{C} , then we have an amalgam in \mathcal{C} . The *amalgamation property* for \mathcal{C} , mentioned earlier, asserts that any amalgam in \mathcal{C} can be embedded into a structure $D \in \mathcal{C}$, i.e. that there are embeddings $j_1: A \rightarrow D$ and $j_2: C \rightarrow D$ such that $j_1f = j_2g$. If \mathcal{C} is a class of finitely generated structures with the AP (for example, a Fraïssé class), then it is a part of folklore in model theory that the statement of the AP extends to non-finitely generated members of $\bar{\mathcal{C}}$ in the following sense.

Lemma 2.1. *Let \mathcal{C} be a class of finitely generated structures enjoying the amalgamation property, and let (A, B, C, f, g) be an amalgam such that $B \in \mathcal{C}$ and $A, C \in \bar{\mathcal{C}}$. Then (A, B, C, f, g) can be embedded into some structure $D \in \bar{\mathcal{C}}$.*

Now suppose that (A, B, C, f, g) is an amalgam in a concrete category \mathcal{C} . Then the \mathcal{C} -pushout P of (A, B, C, f, g) , if it exists, is usually called the *amalgamated free \mathcal{C} -sum* of this amalgam and denoted by $A *_B C$. If, in addition, A, B, C belong to a class \mathcal{C} (contained in \mathcal{C}) which has the AP, then i_1, i_2 must be embeddings. Indeed, by the AP, there exist a structure $D \in \mathcal{C}$ and embeddings $j_1: A \rightarrow D$ and $j_2: C \rightarrow D$, but then there is a homomorphism $u: A *_B C \rightarrow D$ such that $ui_1 = j_1$ and $ui_2 = j_2$, forcing i_1 and i_2 also to be embeddings.

Here we shall consider Fraïssé classes \mathcal{C} satisfying the following *strict amalgamation property*: we require that for any amalgam (A, B, C, f, g) in \mathcal{C} there exist a structure $P \in \mathcal{C}$ and embeddings $i_1: A \rightarrow P$ and $i_2: C \rightarrow P$ such that

$$\begin{array}{ccc} A & \xrightarrow{i_1} & P \\ f \uparrow & & \uparrow i_2 \\ B & \xrightarrow{g} & C \end{array} \quad (2.1)$$

is a pushout square in $\bar{\mathcal{C}}$, considered as a category of structures and homomorphisms. (Here we follow the convention that ‘hooked’ arrows always denote embeddings.) In other words, for any amalgam (A, B, C, f, g) in \mathcal{C} , their amalgamated free $\bar{\mathcal{C}}$ -sum exists and belongs to \mathcal{C} . The previous lemma can be extended to apply to the strict AP.

Lemma 2.2. *Let \mathcal{C} be a Fraïssé class enjoying the strict AP. Then for any amalgam (A, B, C, f, g) such that $B \in \mathcal{C}$ and $A, C \in \bar{\mathcal{C}}$ there exists a $P \in \bar{\mathcal{C}}$ such that (2.1) is a pushout square in $\bar{\mathcal{C}}$.*

Proof. First of all, consider the case when $B, C \in \mathcal{C}$ and $A \in \bar{\mathcal{C}}$. Without loss of generality we can assume that B is a substructure of both A and C and $A \cap C = B$. Let $A = \{a_0, a_1, \dots\}$ be an enumeration of the elements of A . Consider the sequence of finitely generated substructures of A defined by $A_0 = B$ and $A_{n+1} = \langle A_n \cup \{a_{m_n}\} \rangle$ for all $n \geq 0$, where $m_n = \min\{i: a_i \notin A_n\}$. Clearly, $A_i \in \bar{\mathcal{C}}$ for all i since $A \in \bar{\mathcal{C}}$ and $A = \bigcup_{j < \omega} A_j$.

Now let P_n be a sequence of structures and $h_n: A_n \rightarrow P_n$ a sequence of embeddings defined as follows. We set $P_0 = C$ and $h_0 = \mathbf{1}_B$ (where throughout the paper $\mathbf{1}_X$ denotes the identity mapping on the set X), that is, h_0 is just the inclusion map of B into C . For $n \geq 0$, P_{n+1} and h_{n+1} are defined by the requirement that the following diagram is a pushout square in $\bar{\mathcal{C}}$:

$$\begin{array}{ccc} P_n & \hookrightarrow & P_{n+1} \\ \uparrow h_n & & \uparrow h_{n+1} \\ A_n & \hookrightarrow \subseteq & A_{n+1} \end{array}$$

where the symbol ' \subseteq ' will always denote the corresponding inclusion map. Note that P_{n+1} exists and belongs to \mathcal{C} by the strict AP. Therefore, we have a sequence of pushout squares:

$$\begin{array}{ccccccc} C = P_0 & \hookrightarrow & P_1 & \hookrightarrow & P_2 & \hookrightarrow & \dots \\ \uparrow h_0 & & \uparrow h_1 & & \uparrow h_2 & & \\ B = A_0 & \hookrightarrow \subseteq & A_1 & \hookrightarrow \subseteq & A_2 & \hookrightarrow \subseteq & \dots \end{array}$$

Without loss of generality we can assume that P_n is a substructure of P_{n+1} for all $n \geq 0$, so let $P = \bigcup_{j < \omega} P_j$. Clearly, $P \in \bar{\mathcal{C}}$ is a union of a chain of structures from \mathcal{C} . Moreover, $h_0 \subseteq h_1 \subseteq h_2 \subseteq \dots$ and $i_1 = \bigcup_{j < \omega} h_j$ is an embedding $A \hookrightarrow P$. Let i_2 denote the obvious embedding $C = P_0 \hookrightarrow P$. Then the diagram (2.1) commutes and is easily seen to be a pushout square in $\bar{\mathcal{C}}$.

The statement of the lemma is now extended to the case when $C \in \bar{\mathcal{C}}$, essentially by repeating an argument analogous to that above. \square

Let us now quickly review the general method for constructing the Fraïssé limit of a Fraïssé class \mathcal{C} [16]. Recall that a structure C is a *one-point extension* of its substructure B if there is an element $x \in C \setminus B$ such that C is generated by $B \cup \{x\}$. Trivially, if B is finitely generated, so is C .

For a structure $A \in \bar{\mathcal{C}}$, we first construct its extension A^* . Let

$$\{(B_i, C_i): i < \omega\}$$

be the enumeration of all pairs of structures such that $B_i \in \mathcal{C}$ is a finitely generated substructure of A , while C_i is a one-point extension of B_i belonging to \mathcal{C} ; for each isomorphism type we take one such extension. (Here we consider extensions (B, C) and (B, C') to be isomorphic if there is an isomorphism $\iota: C \rightarrow C'$ such that $\iota|_B = \mathbf{1}_B$.) Now we construct a chain of structures A_i , $i \geq 0$, by successive amalgamations of these

extensions. More precisely, let $A_0 = A$ and assume that A_n has already been constructed for some $n \geq 0$ such that $A \subseteq A_n \in \bar{\mathcal{C}}$. Then B_n is a substructure of A and therefore of A_n , whence $(B_n, A_n, C_n, \mathbf{1}_{B_n}, \mathbf{1}_{B_n})$ is an amalgam such that $B_n, C_n \in \mathcal{C}$ and $A_n \in \bar{\mathcal{C}}$. By Lemma 2.1, there exists a structure $A_{n+1} \in \bar{\mathcal{C}}$ which embeds this amalgam. As we have just seen, in the case when \mathcal{C} satisfies the strict AP we can be more specific at this point and let $A_{n+1} = A_n *_{B_n} C_n$ (the free sum is taken with respect to $\bar{\mathcal{C}}$), by Lemma 2.2. Clearly, there is no loss of generality in assuming that $A_n \subseteq A_{n+1}$, so that A is a substructure of A_{n+1} . Finally, we let

$$A^* = \bigcup_{n < \omega} A_n.$$

Of course, this construction can be iterated, so that we apply \aleph_0 successive rounds of amalgamation. Namely, let $A^{(0)} = A$ and define $A^{(n+1)} = (A^{(n)})^*$ for all $n \geq 0$. We set

$$F_A = \bigcup_{n < \omega} A^{(n)},$$

which is an extension of A . Any finitely generated substructure of F_A must belong to some $A^{(m)}$, and, since $A^{(m)} \in \bar{\mathcal{C}}$, the finitely generated structure in question belongs to \mathcal{C} ; hence, $F_A \in \bar{\mathcal{C}}$. A standard model-theoretic argument (which can be found in [7]) shows that F_A is precisely the Fraïssé limit of \mathcal{C} .

Proposition 2.3. *For any $A \in \bar{\mathcal{C}}$ we have $F_A \cong \text{Flim}(\mathcal{C})$.*

Now we wish to modify the above construction, so that, instead of amalgamating the extensions C_i one by one into A^* , we do it ‘all at once’. To this end, suppose we have a structure A and a family of its substructures $\{B_i : i \in I\}$, while $\{C_i : i \in I\}$ is another family of structures such that $A \cap C_i = B_i$ for each $i \in I$ (so that B_i is a substructure of C_i). Furthermore, for any $i, j \in I$, $i \neq j$, we assume that $(C_i \setminus B_i) \cap (C_j \setminus B_j) = \emptyset$. Then the tuple

$$(A, (B_i, C_i)_{i \in I})$$

is called a *rooted multi-amalgam*. Assume that \mathcal{C} is a class of finitely generated structures of a fixed signature. If we are concerned with a rooted multi-amalgam in which $A \in \bar{\mathcal{C}}$ is a countable structure, each B_i is a finitely generated substructure of A , $B_i \in \mathcal{C}$ and each C_i is a finitely generated extension of B_i lying in \mathcal{C} , then we call $(A, (B_i, C_i)_{i \in I})$ a rooted multi-amalgam *over* \mathcal{C} , and the structure A is its *root*.

The *free \mathcal{C} -sum* of the rooted amalgam $(A, (B_i, C_i)_{i \in I})$ over \mathcal{C} is a structure $D \in \bar{\mathcal{C}}$ with the following properties (see Figure 1):

- (a) there are embeddings $f: A \rightarrow D$ and $g_i: C_i \rightarrow D$, $i \in I$, such that $f|_{B_i} = g_i|_{B_i}$ for any $i \in I$;
- (b) for any structure $D' \in \bar{\mathcal{C}}$ and any homomorphisms $\varphi: A \rightarrow D'$, $\psi_i: C_i \rightarrow D'$, $i \in I$, such that for any $i \in I$ we have $\varphi|_{B_i} = \psi_i|_{B_i}$, there exists a unique homomorphism $\delta: D \rightarrow D'$ extending all the given homomorphisms, i.e. such that we have $\delta f = \varphi$ and $\delta g_i = \psi_i$ for all $i \in I$.




Figure 1. A rooted multi-amalgam.

Speaking just a bit more loosely, the free sum is freely generated within $\bar{\mathcal{C}}$ by its partial substructure $A \cup \bigcup_{i \in I} C_i$. Yet another way of saying this is that the free \mathcal{C} -sum is the colimit of the following diagram:

$$\begin{array}{ccccc}
 & & A & & \\
 & \subseteq & \uparrow & \subseteq & \\
 B_i & & B_j & & B_k \quad \dots \\
 \downarrow \subseteq & & \downarrow \subseteq & & \downarrow \subseteq \\
 C_i & & C_j & & C_k \quad \dots
 \end{array} \tag{2.2}$$

The free \mathcal{C} -sum D of $(A, (B_i, C_i)_{i \in I})$, if it exists, is unique up to isomorphism and generated by $f(A) \cup \bigcup_{i \in I} g_i(C_i)$. We denote it by $\coprod^*(A, (B_i, C_i)_{i \in I})$.

Lemma 2.4. *Let \mathcal{C} be a Fraïssé class enjoying the strict AP. Then, for every rooted multi-amalgam $(A, (B_i, C_i)_{i \in I})$ over \mathcal{C} , the free \mathcal{C} -sum $\coprod^*(A, (B_i, C_i)_{i \in I})$ exists and belongs to $\bar{\mathcal{C}}$.*

Proof. Since each B_i, C_i is finitely generated, the index set I is countable, so there is no loss of generality if we assume that I is in fact the ordinal ω . Let us inductively define a sequence of structures $P_i, i \geq 0$, as follows. First of all, let $P_0 = A$. Given P_n , let P_{n+1} be the ($\bar{\mathcal{C}}$ -)pushout of the amalgam $(C_n, B_n, P_n, \mathbf{1}_{B_n}, \mathbf{1}_{B_n})$ (note that B_n is a substructure of $A = P_0$ and so a substructure of each P_n). This pushout exists and belongs to $\bar{\mathcal{C}}$ by Lemma 2.2. Also, there will be no loss of generality in assuming that P_n is actually a substructure of P_{n+1} for each $n \geq 0$.

Let $P = \bigcup_{j < \omega} P_j$. Clearly, $P \in \bar{\mathcal{C}}$, since P is the union of a chain of structures from $\bar{\mathcal{C}}$. Let us show that P satisfies the properties required by the free \mathcal{C} -sum of $(A, (B_i, C_i)_{i \in I})$, i.e. that it is the colimit of the diagram (2.2) in $\bar{\mathcal{C}}$. Obviously, the condition (a) from the definition of a free \mathcal{C} -sum is satisfied, as both A and all C_i (in fact, the whole rooted multi-amalgam $(A, (B_i, C_i)_{i < \omega})$) are contained in P , so the corresponding inclusion mappings will take the role of f and the g_i .




Figure 2. Three commuting diagrams.




Figure 3. The mediating homomorphism.

Now let $Q \in \bar{\mathcal{C}}$ and let $\varphi: A \rightarrow Q$, $\psi_i: C_i \rightarrow Q$, $i \geq 0$, be homomorphisms such that $\varphi|_{B_i} = \psi_i|_{B_i}$ for all i . By the construction of P_1 , there is a unique homomorphism $m_1: P_1 \rightarrow Q$ such that the diagram in Figure 2 (a) commutes. The diagram in Figure 2 (b) also commutes: the triangle commutes by Figure 2 (a), while the square commutes by the initial assumption on Q . Hence, the outer square in Figure 2 (c) commutes for $n = 1$, so there is a unique homomorphism $m_2: P_2 \rightarrow Q$ that makes the entire diagram commutative for $n = 1$. By induction, for every $n \geq 1$, there is a unique homomorphism $m_{n+1}: P_{n+1} \rightarrow Q$ such that the diagram in Figure 2 (c) commutes. Thus, we have a sequence of homomorphisms $\varphi \subseteq m_1 \subseteq m_2 \subseteq \dots$, so define $m = \bigcup_{j < \omega} m_j$. Clearly, $m: P \rightarrow Q$ and the diagram in Figure 3 (a) commutes for all $i \geq 0$.

Let $m': P \rightarrow Q$ be another homomorphism that makes the diagram in Figure 3 (a) commutative for all $i \geq 0$, and let us show that $m' = m$. Let $m'_n = m'|_{P_n}: P_n \rightarrow Q$ for $n \geq 1$ and let $m'_0 = m'|_A$. Clearly, $m'_0 = \varphi$ (see Figure 3 (a)). Specializing to $i = 0$ and using the fact that C_0 and A are substructures of P_1 , we get the diagram in Figure 3 (b). Therefore, $m'_1 = m_1$, since m_1 is the unique homomorphism $P_1 \rightarrow Q$ that makes the diagram commutative. On the other hand, the diagram in Figure 3 (c) commutes (the two triangles commute because of the assumptions on m' and the fact that m'_1 is a restriction of m'_2 and m'_2 is a restriction of m'). Therefore, $m'_2 = m_2$, since m_2 is the unique homomorphism $P_2 \rightarrow Q$ that makes the diagram commutative. Proceeding by induction,

we obtain $m'_n = m_n$ for every $n \geq 1$, so $m' = m$. This completes the proof that P is the colimit of the diagram (2.2) in $\bar{\mathcal{C}}$, i.e. that P is the free \mathcal{C} -sum of $(A, (B_i, C_i)_{i \in I})$. \square

Example 2.5.

- (i) Finite simple graphs have the strict AP, since the free sum of an amalgam of finite graphs is basically the amalgam itself. So, the free sum of a rooted multi-amalgam $(A, (B_i, C_i)_{i \in I})$ over the class of finite simple graphs is the graph with the vertex set $A \cup \bigcup_{i < \omega} C_i$ in which the set of edges is just the union of edges of A and the C_i .
- (ii) Finite posets have the strict AP: consequently, the free sum $\coprod^*(A, (B_i, C_i)_{i \in I})$ of a rooted multi-amalgam over the class of finite posets is just the transitive closure of the reflexive and antisymmetric relation on $A \cup \bigcup_{i < \omega} C_i$ obtained as the union of order relations on A and the C_i .
- (iii) As is well known [14], if \mathbf{V} is a variety of algebras, then $\coprod^*(A, (B_i, C_i)_{i \in I})$ is just the free algebra freely generated in \mathbf{V} by the partial algebra $(A, (B_i, C_i)_{i \in I})$, which exists if and only if the rooted multi-amalgam $(A, (B_i, C_i)_{i \in I})$ can be embedded into a member of \mathbf{V} ; essentially, for algebraic structures the ordinary and the strict AP for the class of finitely generated members of \mathbf{V} are equivalent (bearing in mind Lemmas 2.1 and 2.2). Therefore, finite semilattices, finite distributive lattices and finite Boolean algebras, respectively, have the strict AP and thus all free sums of their rooted multi-amalgams exist in the corresponding varieties.

Given the notion of a rooted multi-amalgam and the conclusion of Lemma 2.4, we slightly change the construction of the extension A^* of a structure $A \in \bar{\mathcal{C}}$. As before, let $\{(B_i, C_i) : i < \omega\}$ be an arbitrary enumeration of all pairs, where B_i is a finitely generated substructure of A and C_i is a one-point extension of B_i , such that each isomorphism type of such pairs is represented exactly once. Then, clearly, $(A, (B_i, C_i)_{i \in I})$ is a rooted multi-amalgam over \mathcal{C} , so we redefine

$$A^* = \coprod^*(A, (B_i, C_i)_{i \in I}).$$

It is not difficult to show that if we iterate this construction and define

$$F_A^* = \bigcup_{n < \omega} A^{(n)},$$

then the result is again isomorphic to the Fraïssé limit of \mathcal{C} . In other words, we have the following variant of the previous proposition.

Proposition 2.6. *For any $A \in \bar{\mathcal{C}}$ we have $F_A^* \cong \text{Flim}(\mathcal{C})$.*

Proof. By [16, Lemma 6.1.3], since F_A^* is countably infinite, it suffices to show that it is *weakly homogeneous* (or that it *realizes all one-point extensions* in the terminology of [9]), i.e. that for each finitely generated substructure B of F_A^* and its one-point extension $C \in \mathcal{C}$ there should be an embedding $f: C \rightarrow F_A^*$ such that $f|_B = \mathbf{1}_B$.

Indeed, since B is finitely generated, there exists an index $k < \omega$ such that $B \subseteq A^{(k)}$. Furthermore, if $\{(B_i^{(k)}, C_i^{(k)}): i < \omega\}$ is the enumeration of all isomorphism types of finitely generated substructures of $A^{(k)}$ and its one-point extensions, yielding the components of the rooted multi-amalgam, the free sum of which is $A^{(k+1)}$, then for some $p < \omega$ we have $B = B_p^{(k)}$ and $C \cong C_p^{(k)}$. By the properties of the free \mathcal{C} -sum, the latter isomorphism gives rise to an embedding of C into $A^{(k+1)}$, and thus into F_A^* . \square

3. A general embedding theorem

A fundamental property of the free \mathcal{C} -sum is that, in order to extend an endomorphism φ of a structure A to an endomorphism $\hat{\varphi}$ of A^* , one only needs to define homomorphisms $\psi_i: C_i \rightarrow A^*$, $i < \omega$, which agree with φ on each B_i . This underlines the importance of the following configuration: we have a finitely generated structure $B \in \mathcal{C}$ and a surjective homomorphism $f: B \rightarrow B'$, where $B' \in \mathcal{C}$; at the same time, we have a one-point extension C of B . In fact, we have a span

$$C \xleftarrow{1_B} B \xrightarrow{f} B'$$

where we follow the convention that ‘double-headed’ arrows stand for surjective homomorphisms (recall that ‘hooked’ ones denote embeddings).

We say that a class \mathcal{C} enjoys the *one-point homomorphism extension property* (1PHEP) if for any $B, B', C \in \mathcal{C}$ forming a configuration as above there exist an extension C' of B' and a surjective homomorphism $f^*: C \rightarrow C'$ such that $f^*|_B = f$; in other words, the following diagram commutes:

$$\begin{array}{ccc} C & \xrightarrow{f^*} & C' \\ \uparrow 1_B & & \uparrow 1_{B'} \\ B & \xrightarrow{f} & B' \end{array}$$

As it easily turns out, either C' is a one-point extension of B' , or it coincides with B' . As noted in [7, Remark 3.1], this property of a Fraïssé class \mathcal{C} is equivalent to the *homomorphism amalgamation property* (HAP), intimately related to homomorphism homogeneity. In fact, by [7, Proposition 3.8], the ultrahomogeneous structure $\text{Flim}(\mathcal{C})$ is homomorphism homogeneous if and only if \mathcal{C} has the 1PHEP. Results of [7] show that 1PHEP is enjoyed by the classes of finite simple graphs, finite posets and finite metric spaces (with rational distances), as well as by each class of finitely generated algebras with the *congruence extension property* (CEP) [14] closed under taking homomorphic images, such as finite semilattices.

We need a stronger form of the 1PHEP for our purpose, related to a construction that in a way represents the analogue of the amalgamated free sum for the spans considered here. We say that a class of structures \mathcal{C} has the *strict 1PHEP* if any span of the form

$$C \xleftarrow{i} B \xrightarrow{f} B',$$

where $B, B', C \in \mathcal{C}$ and C is a one-point extension of B , has a pushout $P \in \mathcal{C}$ with respect to $\bar{\mathcal{C}}$ as a category (of structures and homomorphisms), and if

$$\begin{array}{ccc} C & \xrightarrow{f'} & P \\ \uparrow i & & \uparrow i' \\ B & \xrightarrow{f} & B' \end{array}$$

is a pushout square in $\bar{\mathcal{C}}$, then i' is an embedding and the homomorphism f' is surjective. The following is the main result of this paper.

Theorem 3.1. *Let \mathcal{C} be a Fraïssé class satisfying the following three properties:*

- (i) \mathcal{C} enjoys the strict AP;
- (ii) \mathcal{C} enjoys the strict 1PHEP;
- (iii) for any $B, C \in \mathcal{C}$ such that C is a one-point extension of B , the pointwise stabilizer of B in $\text{Aut}(C)$ is trivial.

Then for any $A \in \bar{\mathcal{C}}$ there is an embedding of $\text{End}(A)$ into $\text{End}(A^*)$. Consequently, if $F = \text{Flim}(\mathcal{C})$, then $\text{End}(A)$ embeds into $\text{End}(F)$.

Proof. We start by considering an arbitrary $\varphi \in \text{End}(A)$. Assume that A^* is obtained as a free \mathcal{C} -sum of the rooted multi-amalgam $(A, (B_i, C_i)_{i \in I})$. Our first task here is to specify the homomorphisms $\psi_i^{(\varphi)}: C_i \rightarrow A^*$, $i < \omega$, agreeing with φ on each B_i . For convenience, write $\varphi_i = \varphi|_{B_i}$. Then there is an index $j < \omega$ such that $\varphi(B_i) = B_j$. By (ii) we have the pushout square

$$\begin{array}{ccc} C_i & \xrightarrow{\xi_i^{(\varphi)}} & C'_i \\ \uparrow \subseteq & & \uparrow \subseteq \\ B_i & \xrightarrow{\varphi_i} & B_j \end{array}$$

However, as a consequence of (iii), the extension (B_j, C_j) is of the same isomorphism type as (B_j, C'_i) . Therefore, there is an isomorphism $\iota_i^{(\varphi)}: C'_i \rightarrow C_j$ such that

$$\begin{array}{ccccc} C_i & \xrightarrow{\xi_i^{(\varphi)}} & C'_i & \xrightarrow{\iota_i^{(\varphi)}} & C_j \\ \uparrow \subseteq & & & \swarrow \subseteq & \uparrow \subseteq \\ B_i & \xrightarrow{\varphi_i} & B_j & & B_j \end{array}$$


Since $B_i B_j C_i C'_i$ is a pushout square and C_j is isomorphic to C'_i , it follows that $B_i B_j C_i C_j$ is also a pushout square. Now we define $\psi_i^{(\varphi)} = \iota_i^{(\varphi)} \xi_i^{(\varphi)}$; therefore, we have a family of homomorphisms

$$(\varphi, (\psi_i^{(\varphi)})_{i < \omega})$$


defined on the components of the rooted multi-amalgam in \mathcal{C} whose sum is A^* . By the properties of the free \mathcal{C} -sum, this family extends to a unique endomorphism $\hat{\varphi}$ of A^* . We claim that the assignment $\varphi \mapsto \hat{\varphi}$ is a monoid embedding $\text{End}(A) \rightarrow \text{End}(A^*)$. Since $\hat{\varphi}|_A = \varphi$, the latter assignment is injective, and since $(\mathbf{1}_A, (\mathbf{1}_{C_i})_{i < \omega})$ yields $\widehat{\mathbf{1}}_A = \mathbf{1}_{A^*}$, we need to prove that for any two $\varphi, \varphi' \in \text{End}(A)$ we have

$$\widehat{\varphi'\varphi} = \widehat{\varphi'}\widehat{\varphi}.$$


To this end, let φ' be another endomorphism of A , $\Phi = \varphi'\varphi$ and $\varphi'_i = \varphi'|_{B_i}$, $\Phi_i = \Phi|_{B_i}$ for any $i < \omega$. We still assume that the index j is selected for a given arbitrary i such that $\varphi(B_i) = B_j$. So, let $\phi'_j(B_j) = B_k$. By the definition of $\widehat{\varphi'}$, there exist a surjective homomorphism $\xi_j^{(\varphi')} : C_j \rightarrow C'_j$ and an isomorphism $\iota_j^{(\varphi')} : C'_j \rightarrow C_k$, where (B_k, C_k) is the representative of the isomorphism type of (B_k, C'_j) , such that $B_j B_k C_j C_k$ in the following diagram



is a pushout square. Then, by the properties of pushouts, $B_i B_k C_i C_k$ is a pushout square. On the other hand, we can repeat the procedure in a slightly different setting to get the following pushout square:



Therefore, there exist homomorphisms $m, n : C_k \rightarrow C_k$ such that the following diagram commutes (both inner and the outer square are pushout squares):



In particular,

$$m\psi_i^{(\varphi'\varphi)} = \psi_j^{(\varphi')}\psi_i^{(\varphi)},$$

$$n(\psi_j^{(\varphi')}\psi_i^{(\varphi)}) = \psi_i^{(\varphi'\varphi)}.$$

Then

$$nm\psi_i^{(\varphi'\varphi)} = n(\psi_j^{(\varphi')}\psi_i^{(\varphi)}) = \psi_i^{(\varphi'\varphi)}, \quad mn(\psi_j^{(\varphi')}\psi_i^{(\varphi)}) = m\psi_i^{(\varphi'\varphi)} = \psi_j^{(\varphi')}\psi_i^{(\varphi)}.$$

Since both $\psi_j^{(\varphi')}\psi_i^{(\varphi)}$ and $\psi_i^{(\varphi'\varphi)}$ are surjective, it follows that $mn = nm = \mathbf{1}_{C_k}$, so m and n are mutually inverse automorphisms of C_k fixing B_k pointwise. Requirement (iii) now yields that $m = n = \mathbf{1}_{C_k}$, whence we obtain

$$\psi_j^{(\varphi')}\psi_i^{(\varphi)} = \psi_i^{(\varphi'\varphi)}.$$

This completes the proof that $\widehat{\varphi'\varphi} = \widehat{\varphi'}\widehat{\varphi}$. □

We say that a one-point extension C of B is *uniquely generated* if for any $x, x' \in C \setminus B$ such that $C = \langle B \cup \{x\} \rangle = \langle B \cup \{x'\} \rangle$ we have $x = x'$. We focus our interest on Fraïssé classes with this property, since it is always present in (but, as we shall see, not confined to) the case when the signature of \mathcal{C} contains no operation symbols (in particular, in all relational structures): the unique generator is the single element in $C \setminus B$. Notice that the uniquely generated one-point extensions instantly satisfy condition (iii) from Theorem 3.1, so that it may be dropped altogether, e.g. for classes of relational structures.

Corollary 3.2. *Let \mathcal{C} be a Fraïssé class satisfying the condition of uniquely generated one-point extensions. If \mathcal{C} satisfies the strict AP and the strict 1PHEP, then for any $A \in \widehat{\mathcal{C}}$, $\text{End}(A)$ embeds into $\text{End}(A^*)$ and thus into the endomorphism monoid of $\text{Flim}(\mathcal{C})$.*

4. Applications

4.1. The cases of the random graph and the generic poset

As noted in items (i) and (ii) of Example 2.5, the classes of finite simple graphs and finite posets enjoy the strict AP and thus admit free sums of their rooted multi-amalgams. Of course, these classes have the property of uniquely generated one-point extensions; thus, we may use Corollary 3.2. Our aim in this section is to briefly re-prove the main results of [2, 6] in order to show that they are transparent instances of our general scheme.

Lemma 4.1. *The class of finite simple graphs satisfies the strict 1PHEP.*

Proof. Let G be a finite simple graph, let $f: G \rightarrow G'$ be a surjective graph homomorphism and let $H = G \cup \{\mathbf{v}\}$ be a one-point extension of G (for convenience, we assume that the embedding i in the definition of the strict 1PHEP is an inclusion map). Let $N_G(\mathbf{v}) \subseteq V(G)$ be the set of all vertices adjacent to \mathbf{v} . Define H' to be the graph obtained from G' by adjoining a new vertex, \mathbf{v}' , which is adjacent to all vertices from $\{f(\mathbf{w}): \mathbf{w} \in N_G(\mathbf{v})\}$ and no other vertex from $V(G')$. Furthermore, let f' be the homomorphism extending f which sends \mathbf{v} to \mathbf{v}' .

We claim that H' , together with $\mathbf{1}_{G'}$ and f' , is a pushout for the span $(G, G', H, f, \mathbf{1}_G)$. Indeed, let Γ be a graph such that there are homomorphisms $g: G' \rightarrow \Gamma$ and $h: H \rightarrow \Gamma$ with $gf = h|_G$. Now there is only one function $u: V(H') \rightarrow V(\Gamma)$ such that $u|_{V(G')} = g$ and $uf' = h$: it is defined by $u(\mathbf{x}) = g(\mathbf{x})$ for $\mathbf{x} \in V(G')$ and $u(\mathbf{v}') = h(\mathbf{v})$. Since g is a

graph homomorphism, and since for any $w \in N_G(v)$ we have that $u(f'(w)) = uf(w) = h(w)$ is adjacent to $h(v) = u(v')$ (as h is also a homomorphism), it follows that u is also a graph homomorphism. Since f' is surjective, the lemma is proved. \square

Corollary 4.2 (Bonato et al. [2]). Self(\aleph_0) (and so any countable semigroup) embeds into $\text{End}(R)$, the endomorphism monoid of the countably infinite random graph.

Proof. In Corollary 3.2 choose A to be the countably infinite graph with no edges, since then $\text{End}(A) \cong \text{Self}(\aleph_0)$. \square


Lemma 4.3. The class of finite posets satisfies the strict 1PHEP.

Proof. Let P be a finite poset, let $f: P \rightarrow P'$ be a surjective poset homomorphism (order-preserving map) and let $Q = P \cup \{x\}$ be a one-point extension of P . In order to define the poset Q' , let

$$L = \{p \in P: p < x\} \quad \text{and} \quad U = \{p \in P: x < p\}.$$

Clearly, $\ell < u$ for any $\ell \in L$ and $u \in U$; therefore, $f(\ell) \leq f(u)$ holds in P' . We distinguish two cases. If $f(L) \cap f(U) = \emptyset$, then we define Q' to be the poset obtained from P' by inserting a new element x' 'between' the down-set $f(L)$ and up-set $f(U)$: we have $y < x'$ if and only if $y \in f(L)$ and $x' < y$ if and only if $y \in f(U)$ (the other elements of P' are set to be incomparable to x'). Then, the homomorphism $f': Q \rightarrow Q'$ is defined as the extension of f obtained by sending $x \mapsto x'$. However, if $f(L) \cap f(U) \neq \emptyset$, then for any $y \in f(L) \cap f(U)$, $\ell' \in f(L)$ and $u' \in f(U)$ we have $\ell' \leq y \leq u'$, implying that such y must be unique: we have $f(L) \cap f(U) = \{y_0\}$ for some $y_0 \in P'$. Then we define $Q' = P'$ and $f': Q \rightarrow P'$ is an extension of f determined by $f(x) = y_0$. In both cases f' is surjective.

We claim that Q' is the pushout of the span $(P, P'Q, f, \mathbf{1}_P)$, together with the poset homomorphisms (order-preserving maps) $\mathbf{1}_{P'}$ and f' ; that is, we claim that there is a unique order-preserving map $m: Q' \rightarrow X$ making the following diagram commutative:



where X is a poset, while $g: P' \rightarrow X$ and $h: Q \rightarrow X$ are order-preserving functions such that $gf = h|_P$. As in the case of graphs, there is a single function $m: Q' \rightarrow X$ that makes the above diagram commutative, and it is defined by $m(y) = g(y)$ for all $y \in P'$ and, if x' exists, $m(x') = h(x)$. Thus, it remains to verify that m is order preserving on Q' . This is clear if $Q' = P'$, since then $m = g$; hence, we may assume that $f(L) \cap f(U) = \emptyset$ and $Q' = P' \cup \{x'\}$. If $y \in P'$ is such that $y < x'$, then $y \in f(L)$, so $m(y) = g(y) = gf(\ell) = h(\ell)$ for some $\ell \in L$. Since $\ell < x$ and h is order preserving on Q ,

we have $m(y) = h(\ell) \leq h(x) = m(x')$. The conclusion that $x' < z$ for $z \in P'$ implies that $m(x') \leq m(z)$ follows in a dual fashion. Finally, m preserves the order on the remaining pairs of distinct elements of Q' , as it agrees with g on such pairs; therefore, m is a poset homomorphism. \square

Corollary 4.4 (Dolinka [6]). *Self(\aleph_0) (and so any countable semigroup) embeds into $\text{End}(\mathbb{P})$, the endomorphism monoid of the generic poset.*

Proof. In Corollary 3.2 choose A to be the countably infinite anti-chain. \square

Actually, in [6] strict endomorphisms of \mathbb{P} were used in order to obtain a more specific version of the above result, thus avoiding the case when $Q' = P'$ in the above lemma. Still, even the strict endomorphisms were sufficient to accomplish the desired embedding of $\text{Self}(\aleph_0)$ into $\text{End}(\mathbb{P})$.

4.2. The case of the (rational) Urysohn space

The traditional way to work with metric spaces is to consider them as pairs (X, d_X) consisting of a non-empty set of points X and a distance function (a *metric*) $d_X: X \times X \rightarrow \mathbb{R}_0^+$ satisfying the triangle inequality. However, there is an easy trick to convert them into first-order relational structures by defining, for each $\alpha \in \mathbb{R}$, $\alpha > 0$, a binary relation R_α on X so that $(x, y) \in R_\alpha$ if and only if $d(x, y) < \alpha$. Furthermore, we may restrict the set of possible distances between points to some countable subset of \mathbb{R}_0^+ , thus obtaining structures over a countable signature; for example, the set \mathbb{Q}_0^+ of non-negative rationals will serve this purpose.

In a posthumous paper [24] (probably the last contribution before his utterly tragic and untimely death), Urysohn showed that up to isometry (a distance-preserving bijection) there is a unique complete separable metric space \mathbb{U} which is ultrahomogeneous and universal, i.e. which embeds all separable metric spaces. Today, \mathbb{U} is called the *Urysohn space* (see, for example, [17, 18, 22, 26, 27] for further background). There is one construction of \mathbb{U} which is of particular interest to us. Namely, let \mathcal{C} be the class of all finite metric spaces with rational distances; it is easily verified that \mathcal{C} is a Fraïssé class. Therefore, this class has a Fraïssé limit, which is denoted by $\mathbb{U}_{\mathbb{Q}}$ and called the *rational Urysohn space*. This space is not complete, but its completion $\bar{\mathbb{U}}_{\mathbb{Q}}$ has all the defining properties of the Urysohn space, so $\bar{\mathbb{U}}_{\mathbb{Q}}$ must be isometric to \mathbb{U} by Urysohn's result.

It was proved in [7, Lemma 3.5] that the class of finite metric spaces has the 1PHEP (and the same is true if we confine ourselves only to spaces with rational distances), implying that both $\mathbb{U}_{\mathbb{Q}}$ and \mathbb{U} are homomorphism homogeneous. Here the notion of a *homomorphism* of metric spaces follows from its first-order-structure representation: for metric spaces X, Y , a function $f: X \rightarrow Y$ is a homomorphism if it is *non-expanding*, that is,

$$d_Y(f(x_1), f(x_2)) \leq d_X(x_1, x_2)$$

holds for all $x_1, x_2 \in X$. In other words, we have the 1-Lipschitz mappings.

Here we prove a somewhat stronger result.

Lemma 4.5. *Let Σ be an additive submonoid of \mathbb{R}_0^+ . The class \mathcal{M}_Σ of all finite metric spaces with distances in Σ enjoys the strict 1PHEP.*

Proof. Let X be a finite metric space and $Y = X \cup \{x^*\}$ its one-point extension. Let $f: X \rightarrow X'$ be a surjective non-expanding map. Let $P = X' \cup \{y^*\}$, where $y^* \notin X'$ is a new point. Let us define a metric on P as follows: for $x, x' \in X'$ let $d_P(x, x') = d_{X'}(x, x')$, while

$$d_P(y^*, x) = \min\{d_Y(x^*, w) + d_{X'}(f(w), x) : w \in X\}.$$

It is easy to verify that d_P is indeed a metric on P . Note that since all the distances in X' and Y are from Σ , so are all distances in P . The mapping $g: Y \rightarrow P$ defined by $g(x) = f(x)$ for $x \in X$ and $g(x^*) = y^*$ is clearly a surjective homomorphism and the diagram

$$\begin{array}{ccc} Y & \xrightarrow{g} & P \\ \uparrow \subseteq & & \uparrow \subseteq \\ X & \xrightarrow{f} & X' \end{array}$$

commutes. Let us show that this is a pushout square (in \mathcal{M}_Σ , but actually it will be a pushout square in the category of all metric spaces). Let Z be a finite metric space (with distances from Σ), and let $\mu: Y \rightarrow Z$ and $\nu: X' \rightarrow Z$ be homomorphisms such that $\mu|_X = \nu f$:

$$\begin{array}{ccc} & & Z \\ & \nearrow \mu & \uparrow \\ Y & \xrightarrow{g} & P \\ \uparrow \subseteq & & \uparrow \subseteq \\ X & \xrightarrow{f} & X' \\ & \searrow \nu & \uparrow \\ & & Z \end{array}$$

Then there exists exactly one mapping $u: P \rightarrow Z$ such that the diagram above commutes; namely, that defined by

$$\begin{aligned} u(x) &= \nu(x), & x \in X', \\ u(y^*) &= \mu(x^*). \end{aligned}$$

Let us show that u is non-expanding, i.e. a homomorphism of metric spaces. Let $x, x' \in X'$. Then $d_Z(u(x), u(x')) = d_Z(\nu(x), \nu(x')) \leq d_{X'}(x, x') = d_P(x, x')$, since ν is a homomorphism. To show $d_Z(u(x), u(y^*)) \leq d_P(x, y^*)$ note that there exists a $w \in X$

such that $d_P(y^*, x) = d_Y(x^*, w) + d_{X'}(f(w), x)$. Hence,

$$\begin{aligned} d_P(y^*, x) &= d_Y(x^*, w) + d_{X'}(f(w), x) \\ &= d_Z(\mu(x^*), \mu(w)) + d_Z(\nu(f(w)), \nu(x)) \quad (\mu \text{ and } \nu \text{ are homomorphisms}) \\ &= d_Z(\mu(x^*), \mu(w)) + d_Z(\mu(w), \nu(x)) \quad (\nu f = \mu|_X) \\ &\leq d_Z(\mu(x^*), \nu(x)) \quad (\text{triangle inequality}) \\ &= d_Z(u(y^*), u(x)) \quad (\text{definition of } u), \end{aligned}$$

as required. □

Metric spaces do not admit coproducts: for example, the existence of a hypothetical coproduct of a metric space X and a trivial space would result in a one-point extension of X by a ‘farthest’ possible point, whereas the values appearing in the vector of distances from the new point to the old ones are obviously not bounded. However, the free *amalgamated* sum of an amalgam of two spaces exists, provided that their non-empty intersection is finite.

Lemma 4.6. *For any additive submonoid Σ of \mathbb{R}_0^+ , the class \mathcal{M}_Σ has the strict AP.*

Proof. Let $(X, Y, Z, \mathbf{1}_Y, \mathbf{1}_Y)$ be an amalgam of finite metric spaces with distances from Σ , so that $Y = X \cap Z$. We define a metric space M on $X \cup Z$ with distances from Σ and prove that M has the properties required by the definition of the free amalgamated sum $X *_Y Z$ with respect to $\bar{\mathcal{M}}_\Sigma$. If either $u, v \in X$ or $u, v \in Z$, then the distance $d_M(u, v)$ coincides with the distance in the corresponding metric space. If, however, $x \in X$ and $z \in Z$, then we define

$$d_M(x, z) = \min\{d_X(x, y) + d_Z(y, z) : y \in Y\}.$$

Notice that the indicated minimum exists because Y is finite. A straightforward verification shows that all triangle inequalities are satisfied, so that we indeed obtain a metric space. Moreover, $M \in \mathcal{M}_\Sigma$.

Assume now that for a (countable) metric space M' (with distances from Σ) we have given metric space homomorphisms $f: X \rightarrow M'$ and $g: Z \rightarrow M'$ such that f and g agree on Y . Then the binary relation $h = f \cup g$ is a function, and we know that it is non-expanding on pairs of points that both belong either to X or to Z . Now let $x \in X \setminus Z$ and $z \in Z \setminus X$. Then $d_M(x, z) = d_X(x, y_0) + d_Z(y_0, z)$ for a particular $y_0 \in Y$, so

$$\begin{aligned} d_{M'}(h(x), h(z)) &= d_{M'}(f(x), g(z)) \\ &\leq d_{M'}(f(x), f(y_0)) + d_{M'}(g(y_0), g(z)) \\ &\leq d_X(x, y_0) + d_Z(y_0, z) = d_M(x, z), \end{aligned}$$

bearing in mind that $f(y_0) = g(y_0)$ and that f and g are homomorphisms (i.e. non-expanding functions). Therefore, $h: M \rightarrow M'$ is a homomorphism of metric spaces, and the proof is complete, since h is obviously the unique function extending both f and g to M . □

We can now turn to the main results of this section.

Corollary 4.7. $\text{Self}(\aleph_0)$ (and so any countable semigroup) embeds into $\text{End}(\mathbb{U}_{\mathbb{Q}})$.

Proof. First of all, note that, by Lemmas 2.4 and 4.6, if all the distances in X and Z_i , $i \in I$, are rational, so are the distances in $\coprod^*(X, (Y_i, Z_i)_{i \in I})$. Now apply Corollary 3.2 while selecting A to be the countably infinite metric space in which the distance between any two different points is 1 (the unit \aleph_0 -simplex) and noticing that any self-map of A is a metric space endomorphism of A . \square

Proposition 4.8. $\text{End}(\mathbb{U}_{\mathbb{Q}})$ embeds into $\text{End}(\mathbb{U})$, so the assertion of Corollary 4.7 also holds for \mathbb{U} .

Proof. Any homomorphism of metric spaces is a uniformly continuous mapping: recall that we are concerned with particular Lipschitz functions. Therefore, any homomorphism $f: X \rightarrow Y$ of metric spaces, where Y is complete, extends uniquely to the completion \bar{X} of X , yielding a (uniformly continuous) function $\bar{f}: \bar{X} \rightarrow Y$. Thus, if $x, y \in \bar{X}$ and if $\{x_n\}_{n < \omega}$ and $\{y_n\}_{n < \omega}$ are any Cauchy sequences converging to x and y , respectively, then

$$\begin{aligned} d(\bar{f}(x), \bar{f}(y)) &= d\left(\bar{f}\left(\lim_{n \rightarrow \infty} x_n\right), \bar{f}\left(\lim_{n \rightarrow \infty} y_n\right)\right) \\ &= d\left(\lim_{n \rightarrow \infty} f(x_n), \lim_{n \rightarrow \infty} f(y_n)\right) \\ &= \lim_{n \rightarrow \infty} d(f(x_n), f(y_n)) \\ &\leq \lim_{n \rightarrow \infty} d(x_n, y_n) \\ &= d(x, y), \end{aligned}$$

implying that \bar{f} is a homomorphism.

So, for any $f, g \in \text{End}(\mathbb{U}_{\mathbb{Q}})$, we have $\bar{f}, \bar{g} \in \text{End}(\mathbb{U})$, and both \overline{gf} and $\bar{g}\bar{f}$ are endomorphisms (and thus uniformly continuous functions) of \mathbb{U} extending gf . Since $\mathbb{U} = \overline{\mathbb{U}_{\mathbb{Q}}}$, we must have $\overline{gf} = \bar{g}\bar{f}$, and the proposition follows. \square

Remark 4.9. Since metric space endomorphisms are just a particular type of Lipschitz function, the monoid of all Lipschitz functions of both $\mathbb{U}_{\mathbb{Q}}$ and \mathbb{U} also embeds $\text{Self}(\aleph_0)$.

Problem 4.10. Is every topological semigroup with a countable base isomorphic to a topological subsemigroup of $\text{End}(\mathbb{U})$?

4.3. The case of the countable universal ultrahomogeneous semilattice

In previous applications of our general results we confined ourselves solely to relational structures; here we consider an example in algebraic structures. However, there is a sense in which semilattices are a unique type of algebra with respect to the method used in this paper (in particular, in Corollary 3.2), making them similar to relational structures.

Lemma 4.11. Any semilattice (S, \wedge) that is a one-point extension of its subsemilattice T is uniquely generated over T .

Proof. Assume $S = \langle T \cup \{x_1\} \rangle = \langle T \cup \{x_2\} \rangle$ for some $x_1, x_2 \in S \setminus T$. Bearing in mind that \wedge is a commutative and idempotent operation, there exist $t, t' \in T$ such that $x_2 = x_1 \wedge t$ and $x_1 = x_2 \wedge t'$. Hence, $x_1 \wedge t' = x_1$ and thus

$$x_1 = x_1 \wedge t \wedge t' = x_1 \wedge t = x_2,$$

as required. □

This behaviour is actually quite atypical for algebraic structures: in general, a one-point extension of an algebra can have many relative generators. Basically, the whole of Galois theory is built around the simple idea of subfield-fixing automorphisms of a given field. Similarly, any $n \times n$ regular diagonal matrix having 1 as all but one of its diagonal entries gives rise to a nonidentical automorphism of an n -dimensional vector space fixing pointwise one of its $(n - 1)$ -dimensional subspaces.

Recall (e.g. from [14]) that an algebra A has the CEP if, for any subalgebra B of A and any congruence ρ of B , there exists a congruence θ of A whose restriction to B is precisely ρ , that is, $\theta \cap (B \times B) = \rho$. Semilattices represent a classical example of algebras with the CEP, so the following lemma applies in particular to them.

Lemma 4.12. *Let A be an algebra with the CEP, and let B be a subalgebra of A . Then for every congruence ρ of B the congruence θ of A generated by ρ (i.e. the smallest $\theta \in \text{Con } A$ containing ρ) has the property that $\theta \cap (B \times B) = \rho$.*

Proof. Since A has the CEP, there exists a congruence θ' of A such that $\theta' \cap (B \times B) = \rho$; but then $\rho \subseteq \theta$, implying $\theta \subseteq \theta'$. Therefore,

$$\rho \subseteq \theta \cap (B \times B) \subseteq \theta' \cap (B \times B) = \rho;$$

thus, $\theta \cap (B \times B) = \rho$ also holds. □

The next assertion is a stronger form of [7, Lemma 3.6].

Lemma 4.13. *Let \mathcal{C} be a class of finitely generated algebras with the CEP closed under taking homomorphic images. Then \mathcal{C} (and, in particular, the class of all finite semilattices) has the strict 1PHEP.*

Proof. Let $A, B, B' \in \mathcal{C}$ be such that $f: B \rightarrow B'$ is a surjective homomorphism and B is a subalgebra of A (the restriction that A is a one-point extension of B will turn out to be immaterial). Then $\ker f$ is a congruence of B . By the CEP and Lemma 4.12, if θ is the congruence of A generated by the relation $\ker f$, then $\theta \cap (B \times B) = \ker f$. Therefore, $B' \cong B/\ker f$ embeds into A/θ by the (well-defined) mapping $\iota: f(b) \mapsto b/\theta, b \in B$. On the other hand, the natural homomorphism $\nu_\theta: A \rightarrow A/\theta$ is obviously surjective. Summing up, the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{\nu_\theta} & A/\theta \\ \uparrow \subseteq & & \uparrow \iota \\ B & \xrightarrow{f} & B' \end{array}$$

By the given conditions, $A/\theta \in \mathcal{C}$. We claim that the above diagram is a pushout square in \mathcal{C} .

To this end, assume that $C \in \mathcal{C}$, while $g: B' \rightarrow C$ and $f: A \rightarrow C$ are two homomorphisms such that $gf = h|_B$:

$$\begin{array}{ccc}
 & & C \\
 & \nearrow h & \\
 A & \xrightarrow{\nu_\theta} & A/\theta \\
 \uparrow \subseteq & & \uparrow \iota \\
 B & \xrightarrow{f} & B' \\
 & \searrow g & \\
 & & C
 \end{array}$$

Clearly, a function $u: A/\theta \rightarrow C$ makes the above diagram commutative if and only if it satisfies the condition $u(a/\theta) = h(a)$ for all $a \in A$. It is immediate that there is at most one such function; we argue that such a function exists, i.e. that the considered condition consistently defines a function u . For this it suffices to show that $\theta \subseteq \ker h$. However, note that $\ker f \subseteq \ker gf$; thus if η is the smallest congruence of A containing the relation $\ker gf$, then $\theta \subseteq \eta$. But $gf = h|_B$, which implies $\ker gf = \ker h \cap (B \times B)$ and so $\ker gf \subseteq \ker h$. Since $\ker h$ is a congruence of A , it follows that $\eta \subseteq \ker h$, yielding the desired conclusion $\theta \subseteq \ker h$. Finally, it is routinely verified that the mapping u is a homomorphism, which completes the proof. \square

Remark 4.14. We have already recorded in Example 2.5 (iii) that the Fraïssé class of finite semilattices has the strict AP. It is possible, however, to be more specific than this. Namely, it is not difficult to prove the following handy representation: we can take $\coprod^*(A, (B_i, C_i)_{i \in I})$ to consist of all finite subsets X of $A \cup \bigcup_{i < \omega} C_i$ such that $|X \cap A| \leq 1$ and $|X \cap C_i| \leq 1$ for all $i < \omega$. Furthermore, $X \wedge Y$ is defined so that for $C \in \{A\} \cup \{C_i: i < \omega\}$ if any of $X \cap C, Y \cap C$ is empty, then $(X \wedge Y) \cap C = (X \cup Y) \cap C$, while otherwise if $X \cap C = \{x\}$ and $Y \cap C = \{y\}$, then $(X \wedge Y) \cap C = \{x \wedge y\}$.

By the following main result of this section we supply an affirmative solution to [8, Problem 4.1].

Corollary 4.15. *Self(\aleph_0) (and so any countable semigroup) embeds into End(Ω), the endomorphism monoid of the countable universal ultrahomogeneous semilattice.*

Proof. In Corollary 3.2 (which applies due to Lemma 4.11), choose the initial semilattice A to be $F(\aleph_0)$, the free semilattice on a countably infinite set of generators. Namely, any self-map of the free generating set of $F(\aleph_0)$ induces an endomorphism of $F(\aleph_0)$. Such endomorphisms form a submonoid of $\text{End}(F(\aleph_0))$ that is isomorphic to $\text{Self}(\aleph_0)$. \square

Previously, it was only known that $\text{Self}(\aleph_0)$ is a homomorphic image of a subsemigroup of $\text{End}(\Omega)$ [7, Corollary 4.4] and that $\text{End}(\Omega)$ embeds all *finite* semigroups [8, Corollary 4.4]. We refer the reader to [1, 8–10] for further properties of Ω , its automorphism group and its endomorphism monoid.

Problem 4.16. Is it possible to embed $\text{Self}(\aleph_0)$ into $\text{End}(\mathbb{D})$ and $\text{End}(\mathbb{A})$, the endomorphism monoids of the countable universal ultrahomogeneous distributive lattice [11] and the countable atomless Boolean algebra, respectively?

The main obstacle here is the lack of the property of uniquely generated one-point extensions, i.e. the existence of non-trivial automorphisms of a one-point extension fixing pointwise the initial structure. Of course, other methods might be developed for approaching this problem; as a very simple example, we mention that $\text{Self}(\aleph_0)$ embeds into the endomorphism monoid of the Fraïssé limit of the class of all finite-dimensional vector spaces over a given field \mathbb{F} . This is because the limit in question is the \aleph_0 -dimensional vector space over \mathbb{F} , which is at the same time a free algebra in the variety of vector spaces over \mathbb{F} , so any self-map of its basis extends uniquely to a linear map of this space into itself.

Acknowledgements. The research of both authors is supported by Grant 174019 of the Ministry of Education and Science of the Republic of Serbia. The authors are grateful to the anonymous referee, whose comments helped to improve the clarity of the presentation.

References

1. M. H. ALBERT AND S. N. BURRIS, Finite axiomatizations for existentially closed posets and semilattices, *Order* **3** (1986), 169–178.
2. A. BONATO, D. DELIĆ AND I. DOLINKA, All countable monoids embed into the monoid of the infinite random graph, *Discrete Math.* **310** (2010), 373–375.
3. P. J. CAMERON, The random graph, in *Algorithms and combinatorics*, Volume 14 (ed. R. L. Graham and J. Nešetřil), pp. 333–351 (Springer, 1997).
4. P. J. CAMERON, The random graph revisited, in *European Congress of Mathematics, Barcelona, 10–14 July 2000, Volume I*, Progress in Mathematics, Volume 201, pp. 267–274 (Birkhäuser, 2001).
5. P. J. CAMERON AND J. NEŠETŘIL, Homomorphism-homogeneous relational structures, *Combinator. Prob. Comput.* **15** (2006), 91–103.
6. I. DOLINKA, The endomorphism monoid of the random poset contains all countable semigroups, *Alg. Univers.* **56** (2007), 469–474.
7. I. DOLINKA, The Bergman property for endomorphism monoids of some Fraïssé limits, *Forum Math.*, doi:10.1515/FORM.2011.153 (in press).
8. I. DOLINKA AND D. MAŠULOVIĆ, Remarks on homomorphism-homogeneous lattices and semilattices, *Monatsh. Math.* **164** (2011), 23–37.
9. M. DROSTE AND D. KUSKE, On random relational structures, *J. Combin. Theory A* **102** (2003), 241–254.
10. M. DROSTE, D. KUSKE AND J. K. TRUSS, On homogeneous semilattices and their automorphism groups, *Order* **16** (1999), 31–56.
11. M. DROSTE AND H. D. MACPHERSON, The automorphism group of the universal distributive lattice, *Alg. Univers.* **43** (2000), 295–306.
12. R. FRAÏSSÉ, Sur certains relations qui généralisent l'ordre des nombres rationnels, *C. R. Acad. Sci. Paris* **237** (1953), 540–542.
13. R. FRAÏSSÉ, Sur l'extension aux relations de quelques propriétés des ordres, *Annales Sci. École Norm. Sup.* **71** (1954), 363–388.
14. G. GRÄTZER, *Universal algebra*, 2nd edn (Springer, 2008).

15. C. W. HENSON, A family of countable homogeneous graphs, *Pac. J. Math.* **38** (1971), 69–83.
16. W. HODGES, *A shorter model theory* (Cambridge University Press, 1997).
17. J. HUBIČKA AND J. NEŠETŘIL, A finite presentation of the rational Urysohn space, *Topol. Applic.* **155** (2008), 1483–1492.
18. M. KATĚTOV, On universal metric spaces, in *Proc. 6th Prague Topological Symp., Prague, 1986* (ed. Z. Frolík), pp. 323–330 (Heldermann, Berlin, 1988).
19. A. H. LACHLAN AND R. E. WOODROW, Countable ultrahomogeneous undirected graphs, *Trans. Am. Math. Soc.* **262** (1980), 51–94.
20. S. MAC LANE, *Categories for the working mathematician*, 2nd edn (Springer, 1998).
21. N. MUDRINSKI, Notes on endomorphisms of Henson graphs and their complements, *Ars Combin.* **96** (2010), 173–183.
22. V. PESTOV, The isometry group of the Urysohn space as a Lévy group, *Topol. Applic.* **154** (2007), 2173–2184.
23. J. H. SCHMERL, Countable homogeneous partially ordered sets, *Alg. Univers.* **9** (1979), 317–321.
24. P. URYSOHN, Sur un espace métrique universel, *Bull. Math. Sci.* **51** (1927), 43–64, 74–96.
25. V. V. USPENSKIJ, On the group of isometries of the Urysohn universal metric space, *Commentat. Math. Univ. Carolinae* **31** (1990), 181–182.
26. V. V. USPENSKIJ, The Urysohn universal metric space is homeomorphic to a Hilbert space, *Topol. Applic.* **139** (2004), 145–149.
27. A. M. VERSHIK, A random metric space is a Urysohn space, *Dokl. Akad. Nauk SSSR* **387** (2002), 733–736 (in Russian).