

HCO⁺ emission possibly related with a shielding mechanism that protects water molecules in the young PN K 3-35

Y. Gómez¹, D. Tafoya^{1,2}, G. Anglada³, L. Loinard¹, J. M. Torrelles⁴,
L. F. Miranda³, M. Osorio³, R. Franco-Hernández^{1,2}, L. Nyman⁵,
J. Nakashima⁶, and S. Deguchi⁷

¹Centro de Radioastronomía y astrofísica, UNAM, México
email: y.gomez@astrosmo.unam.mx

²Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

³Instituto de Astrofísica de Andalucía, CSIC, Granada, Spain

⁴Instituto de Ciencias del Espacio, and IEE de Catalunya, Barcelona, Spain

⁵European Southern Observatory, Santiago, Chile

⁶Department of Physics, University of Hong Kong, Hong Kong

⁷Nobeyama Radio Observatory, NAO, Nagano, Japan

Abstract. Water maser emission has been detected only toward three planetary nebulae (PNe). In particular, in K3-35, the first PN where water vapor maser emission was detected, the components are located in a torus-like structure with a radius of 85 AU and also at the surprisingly large distance of 5000 AU from the star, in the tips of the bipolar lobes. The existence of these water molecules in PNe is puzzling, probably related to some unknown mechanism shielding them against the ionizing radiation. We report the detection of HCO⁺ ($J = 1 - 0$) emission toward K 3-35, that not only suggests that dense molecular gas ($\sim 10^5 \text{ cm}^{-3}$) is present in this PN, but also that this kind of PN can enrich their surroundings with organic molecules.

Keywords. Planetary Nebula, stars: individual (K 3-35), radio lines: molecules, stars: circumstellar matter

1. Introduction

The presence of H₂O maser emission in planetary nebulae (PNe) is opening a new field in the evolutionary study of intermediate mass stars (Gómez 2007, Gómez *et al.* 2008). Even when water maser emission is typically found in the envelopes of asymptotic giant branch (AGB) stars, it is not expected to persist in the PNe phase, where the envelope not only begins to be rarefied but also becomes ionized. Water maser emission has been detected toward three PNe (K 3-35, Miranda *et al.* 2001; IRAS 17347-3139, de Gregorio-Monsalvo *et al.* 2004; IRAS 18061-2505, Gómez *et al.* 2008). In the particular case of K 3-35, the radio continuum emission exhibits a point-symmetric morphology that has been modelled with a precessing jet evolving in a dense AGB circumstellar medium (Velázquez *et al.* 2007). OH and H₂O maser emission has been reported toward this PN (Miranda *et al.* 2001), with the H₂O masers located at the tips of the jets and toward the core in a torus-like structure (Gómez 2007, Uscanga *et al.* 2008 in preparation). In order to understand which is the mechanism that is maintaining the presence of water molecules, we made a search for molecular gas toward K 3-35 under the hypothesis that dense material can protect the water molecules from the UV radiation of the central star.




Figure 1. (a): VLA (3.6 cm) radio continuum image of K 3-35, white dots mark the position of the water masers detected by Miranda *et al.* (2001). (b): HCO^+ ($J = 1 - 0$) spectra detected with the Onsala telescope (Tafuya *et al.* 2007).

2. Summary

The survey for molecular lines toward the young planetary nebula K 3-35 was made using the 20-m telescope of Onsala Space Observatory, the 45-m radio telescope of Nobeyama and the IRAM 30-m telescope. This survey included the molecules SiO, HCO^+ , H^{13}CO^+ , HNC, HCN, HC_5N , HC_3N , CS, CN, CH_3OH , H_2O , CO and ^{13}CO . The details of these observations are published in Tafuya *et al.* (2007).

We discuss here the detection of HCO^+ ($J = 1 - 0$) emission toward K 3-35 (see Figure 1). The HCO^+ peak emission is centered at $\sim 28 \text{ km s}^{-1}$ and the line has a FWHM of $\sim 20 \text{ km s}^{-1}$. We have found that the HCO^+ abundance in K 3-35 is $\sim 6 \times 10^{-7}$, which is similar to values found in other young PNe. Broad CO (2-1) and (1-0) emission were also detected toward K 3-35. Using the CO (2-1;1-0) emission we have estimated an excitation temperature $T_{ex} \sim 20 \text{ K}$. Assuming LTE and that the CO emission is optically thin, a molecular mass for the envelope of $\sim 0.017 M_{\odot}$ is derived. The ratio of molecular to ionized mass is ~ 1.9 . All these results support the presence of a massive molecular envelope in K 3-35 that could be responsible for the shielding mechanism that protects water molecules from being destroyed by the stellar radiation. High angular observations are needed to image the dense molecular gas envelope in this PN.

Acknowledgements

YG acknowledges financial support from PAPIIT-UNAM (IN100407) and CONACyT (49947), México.

References

- de Gregorio-Monsalvo, I., Gómez, Y., Anglada, G., Cesaroni, R., Miranda, L. F., Gómez, J. F., & Torrelles, J. M. 2004, *ApJ* 601, 921
- Gómez, Y. 2007, *Astrophysical masers and their environments*, *IAU Proceedings* 242, 292
- Gómez, J. F., Suárez, O., Gómez, Y., Miranda, L. F., Torrelles, J. M., Anglada, G., & Morata, O. 2008, *AJ* in press.
- Miranda, L. F., Gómez, Y., Anglada, G., & Torrelles, J. M. 2001, *Nature* 414, 284
- Tafuya, D., *et al.* 2007, *ApJ* 133, 364
- Velázquez, P. F., Gómez, Y., Esquivel, A., & Raga, A. C. 2007, *MNRAS* 382, 1965