



Discrete Curvature and Abelian Groups

Bo'az Klartag, Gady Kozma, Peter Ralli, and Prasad Tetali

Abstract. We study a natural discrete Bochner-type inequality on graphs, and explore its merit as a notion of “curvature” in discrete spaces. An appealing feature of this discrete version of the so-called Γ_2 -calculus (of Bakry-Émery) seems to be that it is fairly straightforward to compute this notion of curvature parameter for several specific graphs of interest, particularly, abelian groups, slices of the hypercube, and the symmetric group under various sets of generators. We further develop this notion by deriving Buser-type inequalities (à la Ledoux), relating functional and isoperimetric constants associated with a graph. Our derivations provide a tight bound on the Cheeger constant (*i.e.*, the *edge-isoperimetric constant*) in terms of the spectral gap, for graphs with nonnegative curvature, particularly, the class of abelian Cayley graphs, a result of independent interest.

1 Introduction

For several decades it has been a fruitful endeavour to translate notions from Riemannian geometry to graph theory. It is now clear what are the graph analogs of the Laplacian, Poincaré inequality, Harnack inequality, and many related notions. The graph point of view led to generalizations that would have been less natural in Riemannian geometry, such as β -parabolic Harnack inequalities (see, *e.g.*, [5]) and to some counterexamples [4, 13, 22].

Despite this progress, the graph analog of the notion of curvature remains elusive. In their 1985 paper, Bakry and Émery [2] suggested a notion analogous to curvature that would work in the very general framework of a Markov semigroup (which, of course, incorporates both continuous diffusions and random walks on graphs). The condition was based on the Bochner formula and was denoted by $CD(K, \infty)$ (for curvature-dimension), where K is a curvature parameter. A semigroup satisfying $CD(K, \infty)$ is a generalization of Brownian motion on a manifold with Ricci curvature $\geq K$, and hence the condition $CD(K, \infty)$ is often called simply “ $\text{Ric} \geq K$ ”, and we will stick to this convention here. This notion as a possible definition of Ricci curvature in Markov chains was in fact considered and discussed in [32] in 1999, but seems to have largely been neglected ever since. For additional and more recent approaches to discrete Ricci curvature and related inequalities, see [6, 16, 19, 26, 30, 33, 34]. The fact that one can conclude from positive (or negative) curvature, a local property, global facts about the manifold, has inspired similar “local-to-global” principles in group theory; see *e.g.*, [18, 31].

Received by the editors April 8, 2015; revised September 23, 2015.

Published electronically January 26, 2016.

Research of B. K. was supported in part by the European Research Council. Research of G. K. was supported in part by the Israel Science Foundation and the Jesselson Foundation. Research of P. R. and P. T. was supported in part by the NSF grant DMS-1407657.

AMS subject classification: 53C21, 57M15.

Keywords: Ricci curvature, graph theory, abelian groups.

Beyond lower bounds on curvature, the proofs in [2] (and in the recent book [3]) rely on two additional assumptions on the semigroup. The first was the existence of an appropriate algebra of smooth functions. The second was a chain-rule formula for the generator of the semigroup. A generator satisfying the latter assumption is called a *diffusion operator*; see [3, Definition 1.11.1, p. 43]. In a continuous setting it is actually the existence of the required algebra of smooth functions that is the most difficult condition to verify, but in graph settings, this condition holds immediately. Nevertheless, the diffusion condition can never hold in the discrete setting.

However, the diffusion condition is not always necessary. Denote the Cheeger constant (sometimes known as the isoperimetric constant) by h , the spectral gap by λ and recall the inequality of Buser [8] that states that for a manifold with non-negative Ricci curvature $\lambda \leq 9h^2$ (exact definitions will be given in the next section). In 2006, the first two authors noted that the arguments of Ledoux [23] allow us to derive a discrete Buser-type inequality just by assuming non-negative Ricci curvature.

Theorem 1.1 *A graph satisfying $\text{Ric} \geq 0$ satisfies that $\lambda \leq 16h^2$.*

The graph version of Cheeger's inequality (e.g., [1,10]), which does not require positive curvature, states that $\lambda \geq h^2/(2d)$, where d is the maximum degree of the graph. Thus, for graphs with non-negative Ricci and bounded degree, we get that $\lambda \approx h^2$. As the results from 2006 were never published, we include them in Section 4. A preprint of these results did circulate and a number of papers built on it [6,24]. Particularly relevant for us is the paper [24], which shows that the *eigenvalues* of the Laplacian on a graph with positive curvature satisfy $\lambda_k \leq Ck^2\lambda_1$. In a similar spirit, we use the techniques of [23] to show a Gaussian type isoperimetric inequality for graphs satisfying $\text{Ric} \geq 0$ (see Section 4.3).

In light of Theorem 1.1, an intriguing and challenging open problem is to characterize the class of graphs with non-negative Ricci curvature. The main new results of this paper are examples of such graphs that satisfy $\text{Ric} \geq 0$. These include Cayley graphs of abelian groups, the complete graph, the group S_n with all transpositions, and slices of the hypercube.

In particular, we get Buser's inequality for any Cayley graph of a finite abelian group. We remark that this is not true for a general group. For example, the Cayley graph of the group S_n with the generators being $\{(12), (12 \cdots n)^{\pm 1}\}$ has h of order $1/n^2$ and $\lambda \geq 1/n^3$, up to an absolute constant (we give some details about these well-known facts in Subsection 2.3). This should be compared against the fact that any compact Lie group has positive Ricci curvature; see [9, Corollary 3.19, page 65].

Note that our results above translate to $\lambda(M) \leq 16d h^2(M)$ for a simple random walk M on an abelian Cayley graph, regular of degree d , with $h(M)$ and $\lambda(M)$ being defined for the Markov chain version.

A result of the above type was also recently derived independently by Erbar and by Oveis-Gharan and Trevisan (private communications). An earlier, weaker result, $\lambda(M) = O(d^2 h^2(M))$ follows from the work in [6], which uses a different notion of curvature (and a different argument of Ledoux), starting from a finite-dimensional curvature-dimension $CD(K, n)$ inequality for graphs.

Recently there have been several attempts to modify the $CD(K, n)$ criterion in order to allow certain results involving the heat equation [6, 21, 29]. A recent result of Münch [29] is that the $CDE'(K, n)$ criterion of [21] implies the $CD(K, n)$ criterion of Bakry-Émery. These criteria are often useful; for example, it is known that Ricci-flat graphs satisfy both the $CDE(0, \infty)$ criterion of [6] and the $CDE'(0, \infty)$ criterion.

In the remainder of this section, we introduce Bochner’s Γ_2 -type curvature for graphs along with various notations and definitions. In Section 2, we bound the curvature for several examples, including slices of the discrete cube, symmetric group with adjacent as well as all transpositions as the generating sets; and nonnegativity of curvature for Cayley graphs of abelian groups. In Section 3, we show that the spectral gap can be bounded from below by curvature. In Section 4, we derive the above-mentioned Buser-type inequalities.

1.1 Preliminaries

We first recall some basic definitions and fairly standard notions. Let $G = (V, E)$ be an undirected and locally finite graph. Throughout, we will assume that G has no isolated vertices. We have that the graph Laplacian $\Delta = \Delta(G) = -(D(G) - A(G))$, where $D(G)$ is the diagonal matrix of the degrees of the vertices, and $A(G)$ is the adjacency matrix of G . As an operator, its action on an $f: V \rightarrow \mathbb{R}$ can be described as:

$$\Delta f(x) = \sum_{y \sim x} (f(y) - f(x)),$$

where here and below the notation $y \sim x$ means that y is a neighbour of x in the graph. The sum is, of course, only over the y . Note that Δ is a negative semi-definite matrix.

The spectral gap $\lambda(G)$ is the least non-zero eigenvalue of $-\Delta$. We define the Cheeger constant

$$h(G) = \min_{0 < |A| \leq |V|/2} \frac{|\partial A|}{|A|},$$

where $|\partial A|$ denotes the number of edges from A to $V - A$.

Given functions $f, g: V \rightarrow \mathbb{R}$, we also define:

$$\Gamma(f, g)(x) = \frac{1}{2} \sum_{y \sim x} (f(x) - f(y))(g(x) - g(y)).$$

When $f = g$, the above becomes the more commonly denoted (square of the l_2 -type) discrete gradient. For each $x \in V$,

$$\Gamma(f)(x) := \Gamma(f, f)(x) = \frac{1}{2} \sum_{y \sim x} (f(x) - f(y))^2 =: |\nabla f(x)|^2.$$

It becomes useful to define the iterated gradient

$$2\Gamma_2(f, g) = \Delta\Gamma(f, g) - \Gamma(f, \Delta g) - \Gamma(\Delta f, g).$$

By convention,

$$\Gamma_2(f) := \Gamma_2(f, f) = \frac{1}{2} \Delta \Gamma(f) - \Gamma(f, \Delta f).$$

Note that, given a measure $\pi: V \rightarrow [0, \infty)$, one can consider the expectation (with respect to π) of the above quantity, which gives us the more familiar Dirichlet form associated with a graph:

$$\mathcal{E}(f, g) := \frac{1}{2} \sum_x \sum_{y \sim x} (f(x) - f(y))(g(x) - g(y)) \pi(x).$$

It is useful to note an identity:

$$(1.1) \quad \sum_{x \in V} \Gamma(f, g)(x) = - \sum_{x \in V} f(x) \Delta g(x) = - \sum_{x \in V} g(x) \Delta f(x).$$

An additional useful local identity is

$$(1.2) \quad \Delta(fg) = f \Delta g + 2\Gamma(f, g) + g \Delta f.$$

Definition 1.1 The (Bochner) curvature $\text{Ric}(G)$ of a graph G is defined as the maximum value K such that for any function f and vertex x , we have

$$(1.3) \quad \Gamma_2(f)(x) \geq K\Gamma(f)(x).$$

Let $x \in V$, and let $f: V \rightarrow \mathbb{R}$ be a function. Observe that (1.3) is unchanged on adding a constant to f , so we may assume that $f(x) = 0$. We expand $\Gamma_2(f)(x)$:

$$\begin{aligned} (1.4) \quad 2\Gamma_2(f)(x) &= \Delta \Gamma(f)(x) - 2\Gamma(f, \Delta f)(x) \\ &= \sum_{v \sim x} \Gamma(f)(v) - d(x)\Gamma(f)(x) - \sum_{v \sim x} f(v)(\Delta f(v) - \Delta f(x)) \\ &= \frac{1}{2} \sum_{u \sim v \sim x} (f(u) - f(v))^2 - \frac{d(x)}{2} \sum_{v \sim x} f^2(v) \\ &\quad + \sum_{v \sim x} f(v) \sum_{u \sim x} f(u) - \sum_{u \sim v \sim x} f(v)(f(u) - f(v)) \\ &= \left(\sum_{v \sim x} f(v) \right)^2 - \frac{d(x)}{2} \sum_{v \sim x} f^2(v) + \sum_{u \sim v \sim x} \frac{f^2(u) - 4f(u)f(v) + 3f^2(v)}{2} \\ &= \left(\sum_{v \sim x} f(v) \right)^2 - \sum_{v \sim x} \frac{d(x) + d(v)}{2} f^2(v) + \frac{1}{2} \sum_{u \sim v \sim x} (f(u) - 2f(v))^2. \end{aligned}$$

Now, we break the latter term into the cases where $u = x$, $u \sim x$ and $d(x, u) = 2$. In the second case, we denote by $\Delta(x, v, u)$ the set of all unordered pairs (u, v) satisfying

$x \sim u \sim v \sim x$. The above is equal to

$$\begin{aligned}
 (1.5) \quad 2\Gamma_2(f) &= \frac{1}{2} \sum_{\substack{u \sim v \sim x \\ d(x,u)=2}} (f(u) - 2f(v))^2 + \left(\sum_{v \sim x} f(v) \right)^2 + \sum_{v \sim x} \left(2 - \frac{d(x) + d(v)}{2} \right) f^2(v) \\
 &\quad + \sum_{\Delta(x,v,u)} \frac{(f(v) - 2f(u))^2 + (f(u) - 2f(v))^2}{2} \\
 &= \frac{1}{2} \sum_{\substack{u \sim v \sim x \\ d(x,u)=2}} (f(u) - 2f(v))^2 + \left(\sum_{v \sim x} f(v) \right)^2 + \sum_{v \sim x} \frac{4 - d(x) - d(v)}{2} f^2(v) \\
 &\quad + \sum_{\Delta(x,v,u)} \left[2(f(v) - f(u))^2 + \frac{1}{2}(f^2(v) + f^2(u)) \right].
 \end{aligned}$$

Fixing $f(v)$ for all vertices $v \sim x$, we may ask what choice of $f(u)$ (for $d(x, u) = 2$) minimizes the above expression. We wish to minimize

$$\frac{1}{2} \sum_{\substack{v: \\ x \sim v \sim u}} (f(u) - 2f(v))^2;$$

it is simple to see that the minimizer is

$$(1.6) \quad f(u) = 2 \cdot \frac{1}{r(u)} \sum_{x \sim v \sim u} f(v),$$

where $r(u)$ is the number of common neighbors of u and x .

We first prove a general upper bound on the above notion of curvature, which will be used in the next section to show tightness of our bounds on curvature for several example graphs.

Theorem 1.2 *Let $G = (V, E)$ be a graph. If $e \in E$, let $t(e)$ denote the number of triangles containing e . Define $T := \max_e t(e)$. Then $\text{Ric}(G) \leq 2 + \frac{T}{2}$.*

Proof Let $x \in V$ be any vertex with the minimum degree d , and consider the distance (to x) function $f(v) = \text{dist}(v, x)$. It is simple to calculate that

$$2\Gamma_2(f)(x) \stackrel{(1.5)}{=} d^2 + \sum_{v \sim x} \left(2 - \frac{d + \text{deg}(v)}{2} \right) + \sum_{\Delta(x,v,u)} 1 \leq 2d + \frac{dT}{2},$$

observing that

$$|\Delta(x, v, u)| = \frac{1}{2} \sum_{v \sim x} t(x, v) \leq \frac{dT}{2}$$

and that $\Gamma(f)(x) = \frac{1}{2}d$. Any value of $K > 2 + \frac{T}{2}$ will not satisfy (1.3) for the function f at vertex x , thus $\text{Ric}(G) \leq 2 + \frac{T}{2}$. ■

2 Examples

In this section we provide bounds on the curvature for several graphs of general interest.

2.1 The Hypercube H_n

Let H_n represent the n -dimensional hypercube, where vertices are adjacent if their Hamming distance is one. While the following result also follows from the tensorization result of [24], we provide a direct proof.

Theorem 2.1 $\text{Ric}(H_n) = 2$ if $n \geq 1$.

Proof For any vertex $x \in H_n$, and for any f with $f(x) = 0$, we get from (1.5)

$$2\Gamma_2(f)(x) = \frac{1}{2} \sum_{\substack{u: \\ d(x,u)=2}} \sum_{\substack{v: \\ x \sim v \sim u}} (f(u) - 2f(v))^2 + \left(\sum_{v \sim x} f(v) \right)^2 + (2 - n) \sum_{v \sim x} f^2(v).$$

Let u be a vertex of distance 2 from x , and let v and w be the two distinct vertices such that $u \sim v \sim x \sim w \sim u$. Then for fixed values of $f(v)$ where $v \sim x$, according to (1.6), $\Gamma_2(f)(x)$ is minimized by $f(u) = f(v) + f(w)$. With this value,

$$\sum_{v: u \sim v \sim x} (f(u) - 2f(v))^2 = 2(f(v) - f(w))^2.$$

As for every pair $v, w \sim x$, there is a unique vertex u with $u \sim v, w$ and $d(x, u) = 2$,

$$2\Gamma_2(f)(x) \geq \sum_{\substack{v \neq w \\ v, w \sim x}} (f(v) - f(w))^2 + \left(\sum_{v \sim x} f(v) \right)^2 + (2 - n) \sum_{v \sim x} f^2(v),$$

where the first sum is over all unordered pairs (v, w) of distinct neighbors of x . We use this convention throughout the paper. Expanding the above gives

$$\begin{aligned} & \sum_{\substack{v \neq w \\ v, w \sim x}} (f^2(v) + f^2(w)) - \sum_{\substack{v \neq w \\ v, w \sim x}} 2f(v)f(w) + \sum_{v \sim x} f^2(v) \\ & + \sum_{\substack{v \neq w \\ v, w \sim x}} 2f(v)f(w) + (2 - n) \sum_{v \sim x} f^2(v) = 2 \sum_{v \sim x} f^2(v) = 4\Gamma(f)(x). \end{aligned}$$

So $\text{Ric} \geq 2$, and by Theorem 1.2, we may conclude that $\text{Ric} = 2$. ■

In the sequel, we compute the curvature of the complete graph. With the tensorization result of [24], this provides another proof of the fact that the hypercube has curvature 2.

2.2 The Complete Graph K_n

Theorem 2.2 $\text{Ric}(K_n) = 1 + \frac{n}{2}$ if $n \geq 2$.

Proof For the complete graph on n vertices, we have from (1.5), for every $x \in V$ and every $f: V \rightarrow \mathbb{R}$ such that $f(x) = 0$,

$$\begin{aligned} 2\Gamma_2(f)(x) &= \left(\sum_{v \sim x} f(v) \right)^2 \\ &+ (3 - n) \sum_{v \sim x} f^2(v) + \sum_{\substack{u, v \sim x \\ u \neq v}} \left(2(f(v) - f(u))^2 + \frac{1}{2}(f(u)^2 + f(v)^2) \right). \end{aligned}$$

Expanding the above gives

$$\begin{aligned} & \sum_{v \sim x} f^2(v) + \sum_{\substack{u, v \sim x \\ u \neq v}} 2f(u)f(v) + (3 - n) \sum_{v \sim x} f^2(v) + \frac{5}{2} \sum_{\substack{u, v \sim x \\ u \neq v}} (f^2(v) + f^2(u)) \\ & - \sum_{\substack{u, v \sim x \\ u \neq v}} 4f(u)f(v) \\ & = (4 - n) \sum_{v \sim x} f^2(v) + \frac{5}{2}(n - 2) \sum_{v \sim x} f^2(v) - 2 \sum_{\substack{u, v \sim x \\ u \neq v}} f(u)f(v) \\ & = \left(\frac{3n}{2} - 1\right) \sum_{v \sim x} f^2(v) - 2 \sum_{\substack{u, v \sim x \\ u \neq v}} f(u)f(v) = \frac{3n}{2} \sum_{v \sim x} f^2(v) - \left(\sum_{v \sim x} f(v)\right)^2. \end{aligned}$$

By the Cauchy–Schwarz inequality,

$$\left(\sum_{v \sim x} f(v)\right)^2 \leq |\{v : v \sim x\}| \sum_{v \sim x} f^2(v) = (n - 1) \sum_{v \sim x} f^2(v),$$

so

$$\frac{3n}{2} \sum_{v \sim x} f^2(v) - \left(\sum_{v \sim x} f(v)\right)^2 \geq \left(1 + \frac{n}{2}\right) \sum_{v \sim x} f^2(v).$$

Thus, $\text{Ric} \geq 1 + \frac{n}{2}$, and once again by Theorem 1.2, we conclude that $\text{Ric} = 1 + \frac{n}{2}$. ■

2.3 Finite Abelian Cayley Graphs

A finite abelian group is, of course, a product of cyclic groups, and hence one might think that the curvature of the graph can be deduced from the tensorization result of [24]. However, a Cayley graph is determined by an underlying group and a generating set for that group. Here we show that a finitely generated abelian group with any set of generators has positive Ricci curvature, not only with the generating set inherited from a decomposition into cyclic groups. This result was implicit in the literature, since abelian Cayley graphs are “Ricci flat” [12], and this property, in turn, gives $\text{Ric} \geq 0$ [27]. We give here a direct proof.

Let us remark that the problem of graphs locally identical to an abelian group has also been attacked successfully using combinatorial tools. See [7] and references within.

Theorem 2.3 *Let X be a finitely generated abelian group, and let S be a finite set of generators for X . Let G be the Cayley graph corresponding to X and S . Then $\text{Ric}(G) \geq 0$.*

Recall that the Cayley graph of a group G with respect to a given set S that generates G is the graph whose vertices are the elements of G and whose edges are $\{(g, gs)\}_{g \in G, s \in S}$. Since we are interested in undirected graphs, S should be symmetric, i.e., $s \in S \Rightarrow s^{-1} \in S$.

Proof Without loss of generality, we may set x to be the identity element of X . Denote the degree of every vertex by d . As usual, let $f: G \rightarrow \mathbb{R}$ with $f(x) = 0$.

For this calculation, we prefer not to distinguish between u according to their distance from x , so we start the calculation from (1.4), and using the constant degree we

get

$$(2.1) \quad 2\Gamma_2(f)(x) = d \sum_{v \sim x} f^2(v) + \left(\sum_{v \sim x} f(v) \right)^2 + \sum_{v \sim x} \sum_{u \sim v} \left(\frac{f^2(u)}{2} - 2f(u)f(v) \right).$$

Because x is the identity, we observe that if $u \sim v \sim x$, there is a unique $w \sim x$ so that $u = vw$. We can express the last term of (2.1) as

$$\begin{aligned} & \sum_{v \sim x} \sum_{u \sim v} \left(\frac{f^2(u)}{2} - 2f(u)f(v) \right) \\ &= \sum_{v \sim x} \sum_{w \sim x} \left(\frac{f^2(vw)}{2} - 2f(vw)f(v) \right) \\ &= \sum_{v \sim x} \left(\frac{f^2(v^2)}{2} - 2f(v^2)f(v) \right) + \sum_{\substack{v, w \sim x \\ v \neq w}} \left(f^2(vw) - 2f(vw)(f(v) + f(w)) \right) \\ &\geq -2 \sum_{v \sim x} f^2(v) - \sum_{\substack{v, w \sim x \\ v \neq w}} (f(v) + f(w))^2 = (-d - 1) \sum_{v \sim x} f^2(v) - 2 \sum_{\substack{v, w \sim x \\ v \neq w}} f(v)f(w). \end{aligned}$$

In the last passage we used the elementary inequalities $a^2/2 - 2ab \geq -2b^2$ and $a^2 - 2ab \geq -b^2$

Plugging this bound into (2.1), we find that

$$2\Gamma_2(f)(x) \geq \left(\sum_{v \sim x} f(v) \right)^2 - \sum_{v \sim x} f^2(v) - 2 \sum_{\substack{v, w \sim x \\ v \neq w}} f(v)f(w) = 0.$$

This completes the proof. ■

Now, the assumption that the group is abelian is necessary. An infinite example demonstrating this is the d -ary tree, which is the Cayley graph of the group

$$\langle s_1, \dots, s_d : s_i^2 = \text{id for } i = 1, \dots, d \rangle$$

with the generating set s_1, \dots, s_d . This graph has $\text{Ric} = 2 - d$, which is achieved whenever $\sum_{y \sim x} f(y) = 0$ and $f(z) = 2f(y)$ whenever $z \sim y \sim x$. This is optimal; it is not difficult to see that no d -regular graph has $\text{Ric}(G) < 2 - d$.

A little more surprising, perhaps, is that the Heisenberg group also has negative curvature. We mean here the group of upper triangular matrices with 1 on the diagonal and integer entries, equipped with the set of generators

$$\left\{ \begin{pmatrix} 1 & \pm 1 & 0 \\ & 1 & 0 \\ & & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ & 1 & \pm 1 \\ & & 1 \end{pmatrix} \right\}.$$

It is straightforward to check that these generators do not satisfy any relation of length 4, so the environment within distance 2 (which is the only relevant distance for calculation of the curvature) is tree-like, and the curvature would be -2 .

Switching to finite Cayley graphs, it is well known that there exist finite Cayley graphs that are locally tree-like, and hence would have negative curvature. What is perhaps more interesting is that even Buser’s inequality (the conclusion of Theorem 4.2) may fail.

Theorem 2.4 For the group S_n and the (left) Cayley graph generated by $\{(12), (12 \dots n)^{\pm 1}\}$, the Cheeger constant is $\leq c_1 n^{-2}$, while the spectral gap is $\geq c_2 n^{-3}$, with $c_1, c_2 > 0$, independent of n .

Proof sketch To show an upper bound on the Cheeger constant, we consider the set

$$A = \{ \phi \in S_n : \text{dist}(\phi(1), \phi(2)) \leq \frac{1}{4}n \}$$

(there is no connection between the 1 and 2 in the definition of A and the fact that we took (12) as a generator). Here dist is the cyclic distance between two numbers in $\{1, \dots, n\}$, i.e., $\min(|x - y|, n - |x - y|)$. Clearly, $|A| = (\frac{1}{2} + o(1))n!$. To calculate the size of the boundary we first note that the generators $(12 \dots n)^{\pm 1}$ keep A invariant, so the boundary of A is composed of edges between $\phi \in A$ and $(12)\phi \notin A$. This makes two requirements on ϕ : first it must satisfy that $\text{dist}(\phi(1), \phi(2)) = \lfloor \frac{1}{4}n \rfloor$, and second it must satisfy that one of $\phi(1), \phi(2)$ is in the set $\{1, 2\}$; otherwise, the application of (12) does nothing to $\phi(1)$ and $\phi(2)$ and $(12)\phi$ would still be in A . Thus, $\partial A \approx n!/n^2$ and $h \geq c/n^2$ (this argument gives $c = 2 + o(1)$).

The estimate of the spectral gap (from below) for the random walk on this Cayley graph was done by Diaconis and Saloff-Coste (see [15, Section 5.3]), as an example of the comparison argument; comparing with the random transposition chain, which has a spectral gap of order $1/n$, gives a lower bound of $(1/10)n^{-3}$ for this chain. Since the graph has a bounded degree, the spectral gap of the graph Laplacian is only a constant factor off that of the random walk on the graph.

For the convenience of the reader, and for completeness, we now sketch a proof of a lower bound of $1/(n^3 \log n)$, which serves to justify the point of the theorem. We construct a coupling between two lazy random walkers on our group S_n that succeeds by time $n^3 \log n$. It is well known (see e.g., [25]) that this bounds the mixing time, and hence the relaxation time, which is the inverse of the spectral gap. The coupling is as follows: assume ϕ_n and ψ_n are our two walkers. We apply exactly the same random walks steps to ϕ_n except in one case: when for some i $\phi_n(i) = 1$ and $\psi_n(i) = 2$. In this case when we apply a (12) step for ϕ_n , we apply a lazy step to ψ_n , and vice versa (the $(12 \dots n)^{\pm 1}$ are still applied together). It is easy to check that for each i , $\phi_n(i) - \psi_n(i)$ is doing a random walk on $\{1, \dots, n\}$, slowed down by a factor of n , with gluing at 0. Therefore, it glues with positive probability by time n^3 and with probability $> 1 - 1/2n$ by time $Cn^3 \log n$. Thus, by this time, with probability $> \frac{1}{2}$ we have $\phi(i) = \psi(i)$ for all i , or in other words, the coupling succeeded. This shows that the mixing time is $\leq Cn^3 \log n$ and in turn gives a lower bound on the spectral gap. ■

2.4 Cycles and Infinite Path

We consider the cycle C_n for $n \geq 3$. We extend the notation by letting C_∞ denote the infinite path. From previous results, it is simple to observe that $\text{Ric}(C_3) = \frac{5}{2}$, as $C_3 = K_3$, and that $\text{Ric}(C_4) = 2$, because $C_4 = H_2$.

Theorem 2.5 If $n \geq 5$, then $\text{Ric}(C_n) = 0$.

Proof We note that the calculation of $\text{Ric}(G)$ at x requires us to consider only the subgraph consisting of those vertices v with $d(x, v) \leq 2$ and those edges incident to at least one neighbor of x .

If $n \geq 5$, this subgraph will always be a path of length 4 centered at x , so we only need calculate the curvature for this graph. Since C_n is an abelian Cayley graph, $\text{Ric} \geq 0$.

Then $\text{Ric} = 0$ is achieved by the function f that takes values $-2, -1, 0, 1, 2$ in order along the path. ■

Corollary 2.6 Let \mathbb{Z}^d represent the infinite d -dimensional lattice. $\text{Ric}(\mathbb{Z}^d) = 0$.

We simply note that \mathbb{Z}^d is the product of d copies of C_∞ .

2.5 Slices of the Hypercube

2.5.1 k -slice with Transpositions

For some fixed value k with $1 \leq k < n$, let $G = (V, E)$ be the graph with

$$V = \{x \in \{0, 1\}^n : \sum_i x_i = k\},$$

and $x \sim y$ whenever $|\text{supp}(x - y)| = 2$.

Theorem 2.7 This graph has curvature $\text{Ric} = 1 + \frac{n}{2}$.

Proof Let $x \in V$. Define $s_{ij}x$ to be the vertex obtained by exchanging coordinates i and j in x . A vertex u with $d(x, u) = 2$ will be $u = s_{ij}s_{lm}x$ for some distinct coordinates i, j, l, m with $x_i = x_l = 1, x_j = x_m = 0$. Vertices v with $x \sim v \sim u$ are $s_{ij}x, s_{im}x, s_{lj}x, s_{lm}x$. Observe that

$$\sum_{v: x \sim v \sim u} (f(u) - 2f(v))^2 \geq 2(f(s_{ij}x) - f(s_{lm}x))^2 + 2(f(s_{im}x) - f(s_{lj}x))^2.$$

Summing over all vertices u with $d(x, u) = 2$ gives

$$\frac{1}{2} \sum_{\substack{x \sim v \sim u \\ d(x, u) = 2}} (f(u) - 2f(v))^2 \geq \sum_{\substack{v, w \sim x \\ \nexists(x, v, w)}} (f(v) - f(w))^2,$$

as for each pair $v, w \sim x$ with $v \not\sim w$, there is exactly one u with $v, w \sim u$ and $d(x, u) = 2$. (Here we use the notation $\nexists(x, v, w)$ to denote the set of unordered pairs (v, w) of distinct neighbors of x for which $v \not\sim w$.)

Also notice that any $v \sim x$ has $t(\{x, v\}) = n - 2$: if $v = s_{ij}x$, the vertices that make a triangle with x and v are $s_{lj}x$ when $l \neq i$ and $x_l = x_i$, and $s_{im}x$ when $m \neq j$ and $x_m = x_j$.

Now we can compute

$$\begin{aligned}
 & 2\Gamma_2(f)(x) \\
 & \geq \sum_{\substack{v, w \sim x \\ \Delta(xvw)}} (f(v) - f(w))^2 + \left(\sum_{v \sim x} f(v)\right)^2 + \left(2 - d + \frac{n-2}{2}\right) \sum_{v \sim x} f(v)^2 \\
 & \quad + 2 \sum_{\Delta(vwx)} (f(v) - f(w))^2 \\
 & \geq \sum_{v, w \sim x} (f(v) - f(w))^2 + \left(\sum_{v \sim x} f(v)\right)^2 + \left(1 - d + \frac{n}{2}\right) \sum_{v \sim x} f(v)^2 \\
 & = (d-1) \sum_{v \sim x} f(v)^2 - 2 \sum_{v, w \sim x} f(v)f(w) + \sum_{v \sim x} f(v)^2 + 2 \sum_{v, w \sim x} f(v)f(w) \\
 & \quad + \left(1 - d + \frac{n}{2}\right) \sum_{v \sim x} f(v)^2 \\
 & = \left(1 + \frac{n}{2}\right) \sum_{v \sim x} f(v)^2.
 \end{aligned}$$

So $\text{Ric}(G) \geq 1 + \frac{n}{2}$. Together with Theorem 1.2, we get that $\text{Ric} = 1 + \frac{n}{2}$. ■

2.5.2 Middle Slice with Adjacent Transpositions

We now consider G with $V = \{x \in \{-1, 1\}^{2n} : \sum_i x_i = 0\}$, where $x \sim y \Leftrightarrow \text{supp}(x - y)$ consists of 2 consecutive elements. Alternately, V is the set of paths in \mathbb{Z}^2 that move from $(0, 0)$ to $(2n, 0)$ with steps of $(+1, +1)$ and $(+1, -1)$, and paths x and y are neighbors if y can be achieved by transposing an adjacent $(+1, +1)$ and $(+1, -1)$ in x .

Theorem 2.8 $\text{Ric}(G) \geq -1$. Further, $\lim_{n \rightarrow \infty} \text{Ric}(G) = -1$.

Proof Let $x \in V$. Let $I(x) = \{i \in \{1, \dots, 2n-1\} : x_i \neq x_{i+1}\}$, so $i \in I$ if and only if we are allowed to switch segments i and $i + 1$. If $i \in I(x)$, denote by $a_i x$ the vertex obtained by making this switch. Observe $|I(x)| = \text{deg}(x)$.

The neighbors of $a_i x$ are: $a_i(a_i x) = x$, $a_j(a_i x)$ for any $j \in I(x)$ with $|i - j| > 1$, and $a_j(a_i x)$ for any $j \notin I(x)$ with $|i - j| = 1$ and $j \neq 0, 2n$. We calculate that

$$\text{deg}(a_i x) = \text{deg}(x) + 2 - 2\#\{j \in I(x) : |i - j| = 1\} - \mathbb{1}_{i=1} - \mathbb{1}_{i=2n-1}.$$

We observe that a neighbor of the form $a_j(a_i x)$ if $j \in I(x)$ and $|i - j| > 1$ will be identical to $a_i(a_j x)$ and have $d(x, a_j a_i x) = 2$.

Now, for any function f ,

$$\begin{aligned}
 & \frac{1}{2} \sum_{\substack{u \sim v \sim x \\ d(x,u)=2}} (f(u) - 2f(v))^2 \\
 & \geq \frac{1}{2} \sum_{\substack{i, j \in I \\ |i-j|>1}} (f(a_i a_j x) - 2f(a_i x))^2 + (f(a_i a_j x) - 2f(a_j x))^2
 \end{aligned}$$

$$\begin{aligned} &\geq \sum_{\substack{i,j \in I \\ |i-j| > 1}} (f(a_i x) - f(a_j x))^2 \\ &= \sum_{i \in I(x)} \#\{j \in I(x) : |j - i| > 1\} f^2(a_i x) - 2 \sum_{\substack{i,j \in I \\ |i-j| > 1}} f(a_i x) f(a_j x). \end{aligned}$$

Observe that G is triangle-free. We have that

$$\begin{aligned} 2\Gamma_2(f)(x) &\geq \sum_{i \in I(x)} \#\{j \in I(x) : |j - i| > 1\} f^2(a_i x) - 2 \sum_{\substack{i,j \in I \\ |i-j| > 1}} f(a_i x) f(a_j x) \\ &\quad + \sum_{i \in I(x)} f^2(a_i x) + 2 \sum_{i,j \in I} f(a_i x) f(a_j x) \\ &\quad + \sum_{i \in I(x)} \left(2 - \frac{2 \cdot \deg(x) + 2 - 2\#\{j \in I(x) : |i - j| = 1\} - \mathbb{1}_{i=1} - \mathbb{1}_{i=2n-1}}{2} \right) f^2(a_i x) \\ &\geq \sum_{i \in I(x)} (\#\{j \in I(x) : i \neq j\} + 2 - \deg(x)) f^2(a_i x) + 2 \sum_{\substack{i,j \in I \\ |i-j|=1}} f(a_i x) f(a_j x) \\ &= \sum_{i \in I(x)} f^2(a_i x) + 2 \sum_{\substack{i,j \in I \\ |i-j|=1}} f(a_i x) f(a_j x) > - \sum_{i \in I(x)} f^2(a_i x) \\ &\quad + \sum_{\substack{i,j \in I \\ |i-j|=1}} (f(a_i x) + f(a_j x))^2 \\ &\geq -2\Gamma(f)(x). \end{aligned}$$

So $\text{Ric}(G) > -1$, where we ignore a slight dependence on n in the lower order term.

Define a function with $f(+1, -1, +1, -1, \dots) = 0$ and $f(a_i x) = f(x) - x_i$; that is, if the switch lowers the path, f decreases by 1; a switch that raises the path will increase f by 1.

Using this f and $x = (+1, -1, +1, -1, \dots)$, we find that $\text{Ric} \rightarrow -1$ as $n \rightarrow \infty$. ■

We now calculate the curvature for the subgraph G_+ that is induced on the Dyck paths, *i.e.*, those paths that are always on or above the x -axis, alternately, sequences in $\{\pm 1\}^{2n}$ with $\sum_{i=1}^{2n} x_i = 0$ and $\sum_{i=1}^j x_i \geq 0$ for all $j = 0, \dots, 2n$. It is well known that the number of Dyck paths is the Catalan number C_n .

Corollary 2.9 *For this subgraph G_+ , $\text{Ric}(G_+) \geq -1$. Further, $\lim_{n \rightarrow \infty} \text{Ric}(G_+) = -1$.*

Proof sketch Let $x \in V$ and let

$$I(x) = \{i \in [2n - 1] : \text{a possible move is to transpose } x_i, x_{i+1}\}.$$

If $i \in I$, let $a_i x$ be the sequence obtained by transposing x_i, x_{i+1} .

Observe that $\deg(a_i x) \leq \deg(x) + 2 - 2\#\{j \in I(x) : |i - j| = 1\} - \mathbb{1}_{i=1} - \mathbb{1}_{i=2n-1}$. Using the same analysis as in the unrestricted problem, we can conclude that

$$2\Gamma_2(f)(x) \geq -2\Gamma(f)(x).$$

A similar test-function as above will prove that $\text{Ric} \leq -1 + o(1)$. We can use the same function f and take x identical to the above example but with the first -1 and last $+1$ transposed. This will give a similar upper bound on Ric . (Observe that the neighbors and second-neighbors of x in the unrestricted graph are all Dyck paths, so the curvature at x will be unchanged from the original.) ■

2.6 The Symmetric Group S_n with all Transpositions

Theorem 2.10 *Let G be the Cayley graph on the symmetric group S_n with all transpositions as generators. Then $\text{Ric}(G) = 2$.*

Let us remark that in recent work [17] the authors also provided a lower bound for the Ricci curvature of the (Cayley) graph on the symmetric group with the edge set given by transpositions, but with a different notion of Ricci curvature, one developed by Erbar and Maas [16]. It is easy to see that the Ricci curvature developed by Ollivier [30] gives a value of $\kappa = 2/\binom{n}{2}$ for this problem in the setting of a Markov chain. A simple coupling argument shows that this agrees with our result, modulo the normalizing factor between the graph setting and the Markov chain setting.

Proof Let $x \in S_n$. A vertex u with $d(u, x) = 2$ will either be $(ijk)x$ for some distinct $i, j, k \in [n]$ or $(ij)(kl)x$ for distinct $i, j, k, l \in [n]$.

In the first case, the vertices v s.t. $(ijk)x \sim v \sim x$ are $v = (ij)x, (ik)x, (jk)x$. For $u = (ijk)(x)$,

$$\begin{aligned} & \sum_{v:u \sim v \sim x} (f(u) - 2f(v))^2 \\ &= (f(u) - 2f((ij)x))^2 + (f(u) - 2f((ik)x))^2 + (f(u) - 2f((jk)x))^2 \\ &\geq \frac{4}{3} \left[(f((ij)x) - f((ik)x))^2 + (f((ij)x) - f((jk)x))^2 \right. \\ &\quad \left. + (f((ik)x) - f((jk)x))^2 \right]. \end{aligned}$$

In the second case, a v such that $(ij)(kl)x \sim v \sim x$ is either $v = (ij)x$ or $v = (kl)x$. If $u = (ij)(kl)(x)$,

$$\sum_{v:u \sim v \sim x} (f(u) - 2f(v))^2 \geq 2(f((ij)x) - f((kl)x))^2.$$

Taking a sum over all values of u gives

$$\frac{1}{2} \sum_{\substack{u \sim v \sim x \\ d(u,x)=2}} (f(u) - 2f(v))^2 \geq \sum_{v,w \sim x} (f(v) - f(w))^2.$$

Indeed, if v, w are $v = (ij)x$ and $w = (ik)x$ for some i, j, k , the term $(f(v) - f(w))$ is counted twice in the sum: for $u = (ijk)x$ and $u = (ikj)x$. If v, w are $v = (ij)x$ and $w = (kl)x$ for some i, j, k, l , the term $2(f(v) - f(w))$ is counted once: for $u = (ij)(kl)x$.

Observe that G is triangle-free and regular with degree $d = \binom{n}{2}$. Using this bound, we see that

$$\begin{aligned} 2\Gamma_2(f)(x) &\geq \sum_{v,w \sim x} (f(v) - f(w))^2 + \left(\sum_{v \sim x} f(v) \right)^2 + (2-d) \sum_{v \sim x} f^2(v) \\ &= 2 \sum_{v \sim x} f^2(v) = 4\Gamma(f)(x). \end{aligned}$$

Therefore, $\text{Ric} \geq 2$, as G is triangle-free, $\text{Ric} = 2$ by Theorem 1.2. \blacksquare

3 Spectral Gap and Curvature

Let $\lambda(G)$ denote the spectral gap of G , i.e., the least nonzero eigenvalue of $-\Delta$.

Theorem 3.1 *Let G be a graph with curvature $\text{Ric} \geq K \geq 0$. Then $\lambda \geq K$.*

A different proof of this result was given in [11].

Proof We may use the 2nd derivative versus the first derivative (of variance of the heat kernel) characterization of the spectral gap

$$\lambda = \min_f \frac{\mathcal{E}(-\Delta f, f)}{\mathcal{E}(f, f)},$$

(see e.g., [28]) so that $\alpha \leq \lambda$ if and only if, for any function f , we have $\alpha \cdot \mathcal{E}(f, f) \leq \mathcal{E}(-\Delta f, f)$.

By assumption, G satisfies (1.3) with parameter K , i.e., that

$$\Delta\Gamma(f)(x) - 2\Gamma(f, \Delta f)(x) - 2K\Gamma(f)(x) \geq 0,$$

for all functions $f: V \rightarrow \mathbb{R}$ and all $x \in V$. Summing the above inequality over all vertices gives

$$\begin{aligned} \sum_x \Delta\Gamma(f)(x) - 2 \sum_x \Gamma(f, \Delta f)(x) - 2K \sum_x \Gamma(f)(x) \\ &= 2 \sum_x (\Delta f(x))^2 - K \sum_x \sum_{y \sim x} (f(y) - f(x))^2 \\ &= 2 \sum_x (\Delta f(x))^2 - 2K \sum_{x \sim y} (f(y) - f(x))^2 \geq 0 \end{aligned}$$

where in the first equality, we used identity (1.1) and the fact that for any g , $\sum \Delta g = 0$.

Now let $|V| = n$, and recall the Dirichlet form (with respect to the measure $\pi \equiv 1$),

$$\mathcal{E}(f, f) = \sum_{x \sim y} (f(y) - f(x))^2$$

and that

$$\mathcal{E}(-\Delta f, f) = \sum_x -\Delta f(x) \left(\sum_{y \sim x} (f(x) - f(y)) \right) = \sum_x (\Delta f(x))^2.$$

Plugging into the above inequality gives

$$2\mathcal{E}(-\Delta f, f) - 2K\mathcal{E}(f, f) \geq 0,$$

and so $K\mathcal{E}(f, f) \leq \mathcal{E}(-\Delta f, f)$, resulting in $\lambda \geq K$. \blacksquare

4 Buser-type Inequalities

The proofs in this section are a straightforward discrete version of Ledoux’s paper [23, § 5]. First we derive a key gradient estimate on the heat kernel associated with a graph, which will then be used in to derive a Buser inequality for graphs, as mentioned in the introduction.

4.1 Gradient Estimates

For $t \geq 0$, we write $P_t = \exp(t\Delta)$ for the heat kernel associated with the graph G . Then P_t is a positive definite matrix on \mathbb{R}^V , with P_0 being the identity matrix. Note that P_t commutes with Δ and with P_s , and that $\partial P_t / \partial t = P_t \Delta = \Delta P_t$. Finally, the matrix P_t has non-negative entries. So if f has non-negative entries, then $P_t(f)$ also has non-negative entries. For a vector $f: V \rightarrow \mathbb{R}$, we write $\|f\|_p = (\sum_v |f(v)|^p)^{1/p}$.

Lemma 4.1 *Suppose G has $\text{Ric}(G) \geq K$ for some $K \in \mathbb{R}$. Then for any $f: V \rightarrow \mathbb{R}$ and any $0 \leq t \leq 1/|2K|$,*

$$\|f - P_t f\|_1 \leq 2\sqrt{t} \|\sqrt{\Gamma(f)}\|_1.$$

Note that the restriction on t applies only when K is negative. If $K > 0$, then $\text{Ric} \geq K$ implies $\text{Ric} \geq 0$, and the lemma holds with no restriction on t .

Proof The proof is in three steps.

Step 1. We first prove that

$$\Gamma(P_t f) \leq e^{-2Kt} P_t(\Gamma(f)),$$

where the inequality holds pointwise on V (recalling that these are real-valued functions on V). To that end, define the auxiliary function $g_s = e^{-2Ks} P_s(\Gamma(P_{t-s} f))$, a function on V . It is enough to show that $\partial g_s / \partial s$ is pointwise non-negative on $(0, t)$. We compute

$$\frac{\partial g_s}{\partial s} = e^{-2Ks} P_s [2\Gamma_2(P_{t-s} f) - 2K\Gamma(P_{t-s} f)].$$

Since P_s preserves non-negativity, it is enough to prove that

$$\Gamma_2(P_{t-s} f) - K\Gamma(P_{t-s} f) \geq 0,$$

which is true by our assumption that $\text{Ric}(G) \geq K$.

Step 2. Next we prove that

$$(4.1) \quad P_t(f^2) - (P_t f)^2 \geq \left(\int_0^t 2e^{2Ks} ds \right) \Gamma(P_t f).$$

To that end, define the auxiliary function $g_s = P_s[(P_{t-s} f)^2]$. It is enough to show that $\partial g_s / \partial s \geq 2e^{2Ks} \Gamma(P_t f)$, for any $0 \leq s \leq t$. Using the local identity (1.2) mentioned earlier, we compute

$$\frac{\partial g_s}{\partial s} = P_s [2P_{t-s} f \cdot \Delta P_{t-s} f + 2\Gamma(P_{t-s} f)] + P_s [2P_{t-s} f \cdot (-\Delta P_{t-s} f)].$$

Hence, by Step 1, for any $0 \leq s \leq t$,

$$\frac{\partial g_s}{\partial s} = 2P_s(\Gamma(P_{t-s}f)) \geq 2e^{2Ks}\Gamma(P_t f),$$

which gives (4.1).

Denote $c_K(t) = \int_0^t 2e^{2Ks} ds$. Then $c_K(t) = (e^{2Kt} - 1)/K$, for non-zero K , and $c_K(t) = 2t$ for $K = 0$. In both cases, $c_K(t) \approx 2t$ for small $t > 0$. For instance, $c_K(t) \geq t$ for $0 \leq t \leq 1/(2|K|)$. Hence, for $0 \leq t \leq 1/(2|K|)$, (4.1) gives,

$$(4.2) \quad \max \sqrt{\Gamma(P_t f)} \leq \frac{1}{\sqrt{t}} \max \sqrt{P_t(f^2)} \leq \frac{1}{\sqrt{t}} \max |f|.$$

Step 3. As can be guessed by now, we begin by writing

$$P_t f - f = \int_0^t \frac{\partial P_s f}{\partial s} ds = \int_0^t P_s \Delta f ds.$$

To prove the lemma, it suffices to show that $\|P_s(\Delta f)\|_1 \leq s^{-1/2} \|\sqrt{\Gamma(f)}\|_1$ (since we have $\int_0^t s^{-1/2} ds = 2\sqrt{t}$). Let $\psi = \text{sgn}(P_s(\Delta f))$. Then

$$\begin{aligned} \|P_s(\Delta f)\|_1 &= \sum_{x \in V} P_s(\Delta f)(x) \cdot \psi = \sum_{x \in V} \Delta f(x) \cdot P_s(\psi)(x) = \sum_{x \in V} -\Gamma(f, P_s(\psi))(x) \\ &\leq \sum_{x \in V} \sqrt{\Gamma(f)(x) \cdot \Gamma(P_s(\psi))(x)} \leq \|\sqrt{\Gamma(f)}\|_1 \cdot \max_{x \in V} \sqrt{\Gamma(P_s(\psi))(x)}, \end{aligned}$$

and the desired inequality follows from (4.2), as $\max |\psi| = 1$. ■

4.2 Spectral Gap and Isoperimetry

Theorem 4.2 *Suppose G has $\text{Ric}(G) \geq K$ for some $K \in \mathbb{R}$. Denote by $\lambda > 0$ the minimal non-zero eigenvalue of $-\Delta$. Then, for any subset $A \subset V$,*

$$|\partial A| \geq \frac{1}{2} \min \left\{ \sqrt{\lambda}, \frac{\lambda}{\sqrt{2|K|}} \right\} |A| \left(1 - \frac{|A|}{|V|} \right).$$

Here, by ∂A , we mean the collection of all edges connecting A to its complement.

As noted in the previous lemma, the term $\lambda/\sqrt{2|K|}$ is relevant only in the case $K < 0$.

Proof Apply the previous lemma to $f = \mathbb{1}_A$. Then $\Gamma(\mathbb{1}_A)$ is the function that associates with each $v \in V$, the number of edges in ∂A that are incident with v . Consequently, for any $0 < t < 1/(2|K|)$,

$$\|\mathbb{1}_A - P_t(\mathbb{1}_A)\|_1 \leq 2\sqrt{t} \cdot |\partial A|.$$

Note that $0 \leq P_t(\mathbb{1}_A) \leq 1$, hence the left-hand side can be written as follows:

$$\|\mathbb{1}_A - P_t(\mathbb{1}_A)\|_1 = |A| - \sum_A P_t(\mathbb{1}_A) + \sum_{A^c} P_t(\mathbb{1}_A) = 2 \left[|A| - \sum_V \mathbb{1}_A \cdot P_t(\mathbb{1}_A) \right].$$

Since P_t is self-adjoint and $P_{t/2}P_{t/2} = P_t$,

$$(1/2)\|\mathbb{1}_A - P_t(\mathbb{1}_A)\|_1 = |A| - \|P_{t/2}(\mathbb{1}_A)\|_2^2 = \|\mathbb{1}_A\|_2^2 - \|P_{t/2}(\mathbb{1}_A)\|_2^2.$$

Let $\phi_i : 1 \leq i \leq n$ be the orthonormal eigenvectors of Δ , and let λ_i be the corresponding eigenvalues. Let $\mathbb{1}_A = \sum a_i \phi_i$ be the spectral decomposition of A , with $\phi_0 \equiv 1/\sqrt{|V|}$ and $a_0 = |A|/\sqrt{|V|}$. Then $P_{t/2}(\mathbb{1}_A) = \sum_i a_i e^{-\lambda_i t/2} \phi_i$, and hence

$$(1/2)\|\mathbb{1}_A - P_t(\mathbb{1}_A)\|_1 = \sum_i (1 - e^{-\lambda_i t}) a_i^2 \geq (1 - e^{-\lambda t}) \sum_{i \geq 1} a_i^2 = (1 - e^{-\lambda t}) \left(|A| - \frac{|A|^2}{|V|} \right).$$

To summarize, for any $0 < t \leq 1/(2|K|)$,

$$|\partial A| \geq \frac{1 - e^{-\lambda t}}{\sqrt{t}} |A| \left(1 - \frac{|A|}{|V|} \right).$$

If $\lambda \geq 2|K|$, we select $t = 1/\lambda \leq 1/2|K|$ and deduce the theorem (use $(1 - 1/e) > 1/2$). If $\lambda \leq 2|K|$, we take the maximal possible value, $t = 1/(2|K|)$. Then $1 - e^{-\lambda/2|K|} \geq \lambda/(4|K|)$, and the theorem follows. ■

Corollary 4.3 Suppose a graph G has $\text{Ric}(G) \geq K$, for some $K \geq 0$. Then $h \geq \frac{1}{4}\sqrt{\lambda}$.

Proof As already explained, when $K \geq 0$, we can ignore the term $\lambda/\sqrt{2|K|}$ in the minimum in Theorem 3.1, and then the theorem gives

$$\frac{|\partial A| \cdot |V|}{|A| \cdot |\bar{A}|} \geq \frac{1}{2}\sqrt{\lambda},$$

and so we have $h \geq \frac{1}{4}\sqrt{\lambda}$. ■

4.3 Logarithmic Sobolev Constant and Isoperimetry

We now prove an analogue of [23, Theorem 5.3], relating the log-Sobolev constant ρ to an isoperimetric quantity. Consider the hypercontractive formulation of the log-Sobolev constant (see e.g., [20],[14]): namely, define ρ to be the greatest value so that whenever $1 < r < q < \infty$ and $\sqrt{(q-1)/(r-1)} \leq e^{\rho t}$, then

$$n^{-1/q} \|P_t f\|_q \leq n^{-1/r} \|f\|_r.$$

Theorem 4.4 Suppose G has $\text{Ric}(G) \geq K$ for some value $K \in \mathbb{R}$. Then for any subset $A \subset V$ with $|A| \leq |V|/2 = n/2$,

$$|\partial A| \geq \frac{1}{16} \min\left(\sqrt{\rho}, \frac{\rho}{\sqrt{2|K|}}\right) |A| \log \frac{n}{|A|}.$$

Proof As in the proof of Theorem 4.2, we can observe that

$$\sqrt{t} \frac{|\partial A|}{n} \geq \frac{|A|}{n} - \frac{\|P_{t/2}(\mathbb{1}_A)\|_2^2}{n}$$

if $0 < t < 1/(2|K|)$.

Using the hypercontractivity property with $q = 2$ and $r = 1 + e^{-2\rho t}$ gives that

$$\frac{\|P_{t/2}(\mathbb{1}_A)\|_2^2}{n} \leq \frac{\|\mathbb{1}_A\|_r^2}{n^{2/r}} = \left(\frac{|A|}{n}\right)^{2/r}.$$

Hence,

$$\sqrt{t} \frac{|\partial A|}{n} \geq \frac{|A|}{n} - \frac{\|P_{t/2}(\mathbb{1}_A)\|_2^2}{n} \geq \frac{|A|}{n} - \left(\frac{|A|}{n}\right)^{2/r}.$$

As $2/r \geq 1 + \rho t/4$, whenever $0 \leq \rho t \leq 1$ and $|A|/n \leq 1$,

$$(4.3) \quad \sqrt{t} \frac{|\partial A|}{n} \geq \frac{|A|}{n} - \left(\frac{|A|}{n}\right)^{1+\rho t/4} = \frac{|A|}{n} \left(1 - \left(\frac{|A|}{n}\right)^{\rho t/4}\right).$$

Let $t_0 = \min(1/2|K|, 1/\rho)$. If $|A|/n < e^{-4}$, set $t = \frac{4t_0}{\log(n/|A|)}$.

Using this value of t in (4.3), we find

$$\begin{aligned} \frac{|\partial A|}{n} &\geq \frac{1}{\sqrt{t}} \frac{|A|}{n} (1 - e^{-\rho t_0}) \geq \frac{1}{2\sqrt{t_0}} \frac{|A|}{n} (1 - e^{-\rho t_0}) \log\left(\frac{n}{|A|}\right)^{1/2} \\ &\geq \frac{1}{4} \rho \sqrt{t_0} \frac{|A|}{n} \left(\log \frac{n}{|A|}\right)^{1/2}. \end{aligned}$$

On the other hand, if $e^{-4} \leq |A|/n \leq \frac{1}{2}$, use $t = t_0$ in (4.3) to find

$$\frac{|\partial A|}{n} \geq \frac{1}{\sqrt{t_0}} \frac{|A|}{n} (1 - 2^{-\rho t_0/4}) \geq \frac{1}{8} \rho \sqrt{t_0} \frac{|A|}{n} \geq \frac{1}{16} \rho \sqrt{t_0} \frac{|A|}{n} \left(\log \frac{n}{|A|}\right)^{1/2},$$

where, for the second inequality, we use $1 - 2^{-x} \geq x/2$, if $0 \leq x \leq 1$. Hence,

$$\begin{aligned} \frac{|\partial A|}{n} &\geq \frac{1}{16} \rho \sqrt{\min\left(\frac{1}{2|K|}, \frac{1}{\rho}\right)} \frac{|A|}{n} \left(\log \frac{n}{|A|}\right)^{1/2} \\ &\geq \frac{1}{16} \min\left(\sqrt{\rho}, \frac{\rho}{\sqrt{2|K|}}\right) \frac{|A|}{n} \left(\log \frac{n}{|A|}\right)^{1/2}, \end{aligned}$$

proving the theorem. ■

The optimality of the above theorem (in terms of the dependence on the parameters involved) remains open at this time; in particular, we do not have tight examples. It is also natural to ask if the bound $\rho \geq K$ holds when $\text{Ric} \geq K \geq 0$, similar to the bound on λ in Theorem 3.1. In general, this is not true; consider the complete graph on n vertices. We have seen that $\text{Ric} = 1 + \frac{n}{2}$, and it is easy to see (by considering the characteristic function of a set as a test function), and it is also well known that $\rho = O\left(\frac{n}{\log n}\right)$ (see e.g., [28]).

It is true, however, that under a different notion of discrete curvature for reversible Markov chains, one developed by Erbar and Maas, the so-called *modified* logarithmic Sobolev constant, ρ_0 , can be lower bounded by the curvature; see [16]. Thus, it is certainly interesting to explore whether an analog of Theorem 3.1 is true with ρ_0 in place of λ . Recall here that ρ_0 captures the rate of decay of relative entropy of the Markov chain, relative to the equilibrium distribution, while ρ captures the hypercontractivity property of the Markov kernel (see [28] for additional information).

Acknowledgment The last author thanks Matthias Erbar, Shayan Oveis-Gharan, and Luca Trevisan for discussions on Buser inequality for graphs. We thank the referee for bringing a lot of relevant literature to our attention, especially [11, 12, 27, 29].

References

- [1] N. Alon, O. Schwartz, and A. Shapira, *An elementary construction of constant-degree expanders*. *Combin. Probab. Comput.* 17(2008), no. 3, 319–327. <http://dx.doi.org/10.1017/S0963548307008851>
- [2] D. Bakry and M. Émery, *Diffusions hypercontractives*. In: *Séminaire de probabilités, XIX, 1983/84*, Lecture Notes in Math., 1123, Springer, Berlin, 1985, pp. 177–206. <http://dx.doi.org/10.1007/BFb0075847>
- [3] D. Bakry, I. Gentil, and M. Ledoux, *Analysis and geometry of Markov diffusion operators*. *Grundlehren der Mathematischen Wissenschaften*, 348, Springer, 2014. <http://dx.doi.org/10.1007/978-3-319-00227-9>
- [4] M. Barlow, *Which values of the volume growth and escape time exponent are possible for a graph?* *Rev. Mat. Iberoamericana* 20(2004), no. 1, 1–31. <http://dx.doi.org/10.4171/RMI/378>
- [5] M. Barlow and R. Bass, *Stability of parabolic Harnack inequalities*. *Trans. Amer. Math. Soc.* 356(2004), no. 4, 1501–1533. <http://dx.doi.org/10.1090/S0002-9947-03-03414-7>
- [6] F. Bauer, P. Horn, Y. Lin, G. Lippner, D. Mangoubi, and S.-T. Yau, *Li-Yau inequality on graphs*. *J. Differential Geom.* 99(2015), no. 3, 359–405.
- [7] I. Benjamini and D. Ellis, *On the structure of graphs which are locally indistinguishable from a lattice*. *arxiv:1409.7587*
- [8] P. Buser, *A note on the isoperimetric constant*. *Annales scientifiques de l'École Normale Supérieure* 15(1982), no. 2, 213–230.
- [9] J. Cheeger and D. Ebin, *Comparison theorems in Riemannian geometry*. North-Holland Mathematical Library, 9, North-Holland Publishing Co., Amsterdam-Oxford; American Elsevier Publishing Co., Inc., New York, 1975.
- [10] F. Chung, *Four proofs for the Cheeger inequality and graph partition algorithms*. In: *Fourth International Congress of Chinese Mathematicians*, AMS/IP Stud. Adv. Math., 48, American Mathematical Society, Providence, RI, 2010, pp. 331–349.
- [11] F. Chung, Y. Lin, and S.-T. Yau, *Harnack inequalities for graphs with non-negative Ricci curvature*. *J. Math. Anal. Appl.* 415(2014), no. 1, 25–32. <http://dx.doi.org/10.1016/j.jmaa.2014.01.044>
- [12] F. R. K. Chung and S.-T. Yau, *Logarithmic Harnack inequalities*. *Math. Res. Lett.* 3(1996), no. 6, 793–812. <http://dx.doi.org/10.4310/MRL.1996.v3.n6.a8>
- [13] T. Delmotte, *Graphs between the elliptic and parabolic Harnack inequalities*. *Potential Anal.* 16(2002), no. 2, 151–168. <http://dx.doi.org/10.1023/A:1012632229879>
- [14] P. Diaconis and L. Saloff-Coste, *Logarithmic Sobolev inequalities for finite Markov chains*. *Ann. Appl. Probab.* 6(1996), no. 3, 695–750. <http://dx.doi.org/10.1214/aoap/1034968224>
- [15] ———, *Random walks on finite groups: a survey of analytic techniques*. In: *Probability measures on groups and related structures*, XI (Oberwolfach, 1994), pp. 44–75.
- [16] M. Erbar and J. Maas, *Ricci curvature of finite Markov chains via convexity of the entropy*. *Arch. Ration. Mech. Anal.* 206(2012), no. 3, 997–1038. <http://dx.doi.org/10.1007/s00205-012-0554-z>
- [17] M. Erbar, J. Maas, and P. Tetali, *Discrete Ricci curvature bounds for Bernoulli-Laplace and random transposition models*. *Annales Fac. Sci. Toulouse*, to appear. *arxiv:1409.8605*
- [18] H. Garland, *p-adic curvature and the cohomology of discrete subgroups of p-adic groups*. *Ann. of Math.* (2) 97(1973), 375–423. <http://dx.doi.org/10.2307/1970829>
- [19] N. Gozlan, C. Roberto, P.-M. Samson, and P. Tetali, *Displacement convexity of entropy and related inequalities on graphs*. *Probability Theory and Related Fields* 160(2014), 47–94. <http://dx.doi.org/10.1007/s00440-013-0523-y>
- [20] L. Gross, *Logarithmic Sobolev inequalities*. *Amer. J. Math.* 97(1975), no. 4, 1061–1083. <http://dx.doi.org/10.2307/2373688>
- [21] P. Horn, Y. Lin, S. Liu, and S.-T. Yau, *Volume doubling, Poincaré inequality and Gaussian heat kernel estimate for nonnegative curvature graphs*. *arxiv:1411.5087*
- [22] G. Kozma, *A graph counterexample to Davies' conjecture*. *Rev. Mat. Iberoam.* 30(2014), no. 1, 1–12. <http://dx.doi.org/10.4171/RMI/766>
- [23] M. Ledoux, *Spectral gap, logarithmic Sobolev constant, and geometric bounds*. *Surveys in Diff. Geom.*, 9(2004), 219–240.
- [24] S. Liu, and N. Peyerimhoff, *Eigenvalue ratios of nonnegatively curved graphs*. *arxiv:1406.6617*

- [25] D. Levin, Y. Peres, and E. Wilmer, E., *Markov chains and mixing times*. American Mathematical Society, Providence, RI, 2009.
- [26] J. Lott and C. Villani, *Ricci curvature for metric-measure spaces via optimal transport*. *Ann. Math.* (2) **169**(2009), no. 3, 903–991. <http://dx.doi.org/10.4007/annals.2009.169.903>
- [27] Y. Lin and S.-T. Yau, *Ricci curvature and eigenvalue estimate on locally finite graphs*. *Math. Res. Lett.* **17**(2010), no. 2, 343–356. <http://dx.doi.org/10.4310/MRL.2010.v17.n2.a13>
- [28] R. Montenegro and P. Tetali, *Mathematical aspects of mixing times in Markov chains*. *Found. Trends Theor. Comput. Sci.* **1**(2006), no. 3.
- [29] F. Münch, *Remarks on curvature dimension conditions on graphs*. [arxiv:1501.05839](https://arxiv.org/abs/1501.05839)
- [30] Y. Ollivier, *Ricci curvature of Markov chains on metric spaces*. *J. Funct. Anal.* **256**(2009), no. 3, 810–864. <http://dx.doi.org/10.1016/j.jfa.2008.11.001>
- [31] P. Papasoglu, *An algorithm detecting hyperbolicity*. In :Geometric and computational perspectives on infinite groups (Minneapolis, MN and New Brunswick, NJ, 1994), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 25, American Mathematical Society, Providence, RI, 1996, pp.193–200.
- [32] M. Schmuckenschläger, *Curvature of nonlocal Markov generators*. In: *Convex geometric analysis* (Berkeley, CA, 1996), Cambridge Univ. Press, Cambridge, 1999, pp. 189–197.
- [33] M. D. Sammer, *Aspects of mass transportation in discrete concentration inequalities*. PhD thesis, Georgia Institute of Technology, 2005. http://etd.gatech.edu/theses/available/etd-04112005-163457/unrestricted/sammer_marcus_d_200505_phd.pdf
- [34] K.-Th. Sturm, *On the geometry of metric measure spaces. I*. *Acta Math.* **196**(2006), no. 1, 65–131. <http://dx.doi.org/10.1007/s11511-006-0002-8>

School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel
e-mail: klartagb@tau.ac.il

Department of Mathematics, The Weizmann Institute of Science, Rehovot 76100, Israel
e-mail: gady.kozma@weizmann.ac.il

School of Mathematics and School of Computer Science, Georgia Institute of Technology, Atlanta, GA 30332, USA
e-mail: pralli3@math.gatech.edu tetali@math.gatech.edu