



Weighted nonlinear flag manifolds as coadjoint orbits

Stefan Haller  and Cornelia Vizman 

Abstract. A weighted nonlinear flag is a nested set of closed submanifolds, each submanifold endowed with a volume density. We study the geometry of Fréchet manifolds of weighted nonlinear flags, in this way generalizing the weighted nonlinear Grassmannians. When the ambient manifold is symplectic, we use these nonlinear flags to describe a class of coadjoint orbits of the group of Hamiltonian diffeomorphisms, orbits that consist of weighted isotropic nonlinear flags.

1 Introduction

The nonlinear Grassmannian $Gr_S(M)$, consisting of all smooth submanifolds in a manifold M which are diffeomorphic to a closed manifold S , has a natural Fréchet manifold structure. Nonlinear Grassmannians, a.k.a. differentiable Chow manifolds or shape spaces, play an important role in computer vision [1, 20] and continuum mechanics [21]. They have also been used to describe coadjoint orbits of diffeomorphism groups [8]. Further coadjoint orbits of diffeomorphism groups can be described using weighted nonlinear Grassmannians, i.e., spaces of submanifolds equipped with volume densities. For instance, weighted nonlinear Grassmannians of isotropic submanifolds in a symplectic manifold have been used to describe coadjoint orbits of the Hamiltonian group [7, 15, 26], and weighted nonlinear Grassmannians of isotropic submanifolds in a contact manifold have been identified with coadjoint orbits of the contact group [10].

Nonlinear Grassmannians have been generalized to spaces of nonlinear flags in [9]. Given a collection of closed manifolds $S = (S_1, \dots, S_r)$, a nonlinear flag of type S in M is a sequence of nested embedded submanifolds $N_1 \subseteq \dots \subseteq N_r$ in M , with N_i diffeomorphic to S_i , for all i . The space of all nonlinear flags of type S in M is a Fréchet manifold in a natural way, denoted by $Flag_S(M)$. Manifolds of low-dimensional nonlinear flags have appeared as shape spaces in [3, 12, 23]. Symplectic nonlinear flags have been used to describe coadjoint orbits of the Hamiltonian group [9].

Received by the editors December 31, 2022; revised August 6, 2023; accepted October 2, 2023.

Published online on Cambridge Core October 9, 2023.

The first author gratefully acknowledges the support of the Austrian Science Fund (FWF) (Grant No. P31663). The second author was partially supported by CNCS UEFISCDI, project number PN-III-P4-ID-PCE-2020-2888.

AMS subject classification: 58D10, 37K65, 53C30, 53D20, 58D05.

Keywords: Weighted nonlinear flag, coadjoint orbit, Hamiltonian group, moment map, manifold of mappings.



In this article, we study manifolds of weighted nonlinear flags, motivated by the fact that one can use them to describe new coadjoint orbits of the Hamiltonian group. Considering submanifolds N_i equipped with nowhere zero densities ν_i , one obtains the manifold $\text{Flag}_S^{\text{wt}}(M)$ of weighted nonlinear flags. We describe its Fréchet manifold structure in two ways: as a splitting smooth submanifold in the Cartesian product of weighted nonlinear Grassmannians of type S_i in M , and as a locally trivial smooth fiber bundle over $\text{Flag}_S(M)$ associated with the principal bundle of nonlinear frames of type S in M . To each weighted nonlinear flag, one associates a compactly supported distribution on M by the $\text{Diff}(M)$ equivariant inclusion

$$(1.1) \quad J : \text{Flag}_S^{\text{wt}}(M) \hookrightarrow C^\infty(M)^*, \quad \langle J((N_1, \nu_1), \dots, (N_r, \nu_r)), f \rangle := \sum_{i=1}^r \int_{N_i} f \nu_i.$$

We aim at describing the $\text{Diff}(M)$ orbits in $\text{Flag}_S^{\text{wt}}(M)$. Under the $\text{Diff}(M)$ action, the diffeomorphism type of the inclusions $N_1 \subseteq \dots \subseteq N_r$ remains invariant, and the total volumes of the connected components of $N_i \setminus N_{i-1}$ with respect to ν_i remain invariant as well, for all i . This suggests to fix embeddings $S_1 \xrightarrow{t_1} S_2 \xrightarrow{t_2} \dots \xrightarrow{t_{r-1}} S_r$ and volume densities μ_i on S_i , and to consider the $\text{Diff}(M)$ invariant subspace $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M)$ consisting of all weighted nonlinear flags diffeomorphic to the model (S, ι, μ) , where $\iota = (\iota_1, \dots, \iota_{r-1})$ and $\mu = (\mu_1, \dots, \mu_r)$. Using a Moser-type argument, we will show that these are exactly the weighted nonlinear flags for which the volumes of corresponding connected components in $N_i \setminus N_{i-1}$ and in $S_i \setminus \iota_{i-1}(S_{i-1})$ coincide. A more precise statement, using homologically weighted nonlinear flags, is formulated in Theorem 2.15. This description permits to show that $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M)$ is a splitting smooth submanifold of finite codimension in $\text{Flag}_S^{\text{wt}}(M)$ (see Theorem 2.11). Furthermore, the $\text{Diff}(M)$ action on $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M)$ admits local smooth sections. In particular, every $\text{Diff}(M)$ orbit is a union of connected components in $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M)$.

For a symplectic manifold M , we are interested in the orbits of the Hamiltonian group $\text{Ham}_c(M)$ acting on $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M)$. Clearly, the open subset of weighted symplectic nonlinear flags $\text{Flag}_{S, \iota, \mu}^{\text{wt symplectic}}(M)$ is invariant under this action. Although the Hamiltonian group acts locally transitive on unweighted symplectic nonlinear flags [9, Proposition 4.4], the action on $\text{Flag}_{S, \iota, \mu}^{\text{wt symplectic}}(M)$ is not locally transitive, not even in the Grassmannian case [7, Section 4]. Some orbits turn out to be nice submanifolds, e.g., when the weights are provided by the Liouville volume forms (cf. Remark 2.6), but the general orbit is considerably more singular.

In this paper, we will consider the action of the Hamiltonian group on the invariant set of weighted isotropic nonlinear flags $\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M)$. In view of the tubular neighborhood theorem for isotropic embeddings [25], this is a splitting smooth submanifold in $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M)$. If $H^1(S; \mathbb{R}) = 0$, the action is locally transitive. In general, the orbits of the Hamiltonian group provide a smooth foliation of codimension $\dim H^1(S; \mathbb{R})$ in $\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M)$, the analogue of Weinstein's isodrastic foliation in the Grassmannian case [26]. Each isodrastic leaf of weighted nonlinear flags comes equipped with a canonical symplectic form, and the map J in (1.1) restricts to an equivariant moment map for the $\text{Ham}_c(M)$ action, thus identifying the leaf with a coadjoint orbit of the Hamiltonian group (see Theorem 3.15). Moreover, this coadjoint

orbit is a splitting symplectic submanifold in a product of coadjoint orbits of weighted submanifolds of type S_i in M (see Remark 3.16). The lowest-dimensional examples are the coadjoint orbits of $\text{Ham}_c(\mathbb{R}^2)$ consisting of pointed weighted vortex loops, treated in [4]. We give more examples with nested spheres or tori, and provide explicit descriptions of the corresponding coadjoint orbits of the Hamiltonian group.

The results on weighted nonlinear flags sketched above generalize well-known results on weighted nonlinear Grassmannians, which correspond to $r = 1$. Many of the proofs presented in this article proceed by induction on the depth r of the nonlinear flags. Often a relative version of the corresponding result on nonlinear Grassmannians is required for the induction step. These relative versions are not readily available in the literature and often require new arguments. Lemma 2.12, for instance, is crucial in the proof of Proposition 2.10, and Lemma 3.7 is used in the proof of Proposition 3.14. Similar relative versions also play an important role in the study of unweighted nonlinear flags (see [9, Lemma 2.1] used, e.g., in the proof of [9, Proposition 2.9]). In Remark 2.14, we indicate the significance of a relative version of a well-known result on the group of volume-preserving diffeomorphisms [11] which appears naturally in this context. To the best of our knowledge, the latter relative version is still open.

2 Manifolds of weighted nonlinear flags

A nonlinear flag is a sequence of nested closed submanifolds $N_1 \subseteq \dots \subseteq N_r$ in a smooth manifold M . A weighted nonlinear flag is a nonlinear flag together with a volume density ν_i on each submanifold N_i . Integrating against test functions $f \in C^\infty(M)$, a weighted nonlinear flag provides a compactly supported distribution on M with mild singularities, $\sum_{i=1}^r \int_{N_i} f \nu_i$.

We will show that the space of all weighted nonlinear flags in M is a Fréchet manifold in a natural way. In fact, this is the total space of a locally trivial smooth bundle over the manifold of nonlinear flags discussed in [9]. The natural $\text{Diff}(M)$ action on the base of this bundle is locally transitive [9, Proposition 2.9(a)]. The main aim of this section is to describe the $\text{Diff}(M)$ orbits in the space of weighted nonlinear flags (see Theorem 2.11).

2.1 Weighted nonlinear Grassmannians

In this section, we recall some basic facts about the manifolds of weighted submanifolds that appear in [7, 15, 26]. These weighted nonlinear Grassmannians constitute a special case of the weighted nonlinear flags to be introduced in Section 2.2. We present them here in a manner that readily generalizes to the setting of nonlinear flags.

Let S be a closed manifold of dimension k , allowed to be nonconnected and nonorientable. For each manifold M , we let $\text{Gr}_S(M)$ denote the *nonlinear Grassmannian of type S in M* , i.e., the space of all smooth submanifolds in M that are diffeomorphic to S . Moreover, we let $\text{Emb}_S(M)$ denote the space of all parametrized submanifolds of type S in M , i.e., the space of all smooth embeddings of S into M . Both, $\text{Emb}_S(M)$ and $\text{Gr}_S(M)$, are Fréchet manifolds in a natural way. Furthermore, the $\text{Diff}(M)$ equivariant map

$$(2.1) \quad \text{Emb}_S(M) \rightarrow \text{Gr}_S(M), \quad \varphi \mapsto \varphi(S),$$

is a smooth principal bundle with structure group $\text{Diff}(S)$, a Fréchet Lie group (see [2, 6, 18, 19] and [14, Theorem 44.1]).

For each closed k -dimensional manifold S , let

$$\text{Den}(S) := \Gamma^\infty(|\Lambda|_S) = \Omega^k(S; \mathcal{O}_S)$$

denote the space of all smooth densities on S . Here, \mathcal{O}_S denotes the orientation bundle of S and $|\Lambda|_S = \Lambda^k T^*S \otimes \mathcal{O}_S$ (see, e.g., [16]). Densities on S are the geometric quantities which can be integrated over S in a coordinate independent way, without specifying an orientation or even assuming orientability. We denote by $\text{Den}_\times(S)$ the space of *volume densities*, i.e., the space of nowhere vanishing densities. Clearly, this is an open subset in the Fréchet space $\text{Den}(S)$.

We define the *weighted nonlinear Grassmannian of type S in M* by

$$(2.2) \quad \text{Gr}_S^{\text{wt}}(M) := \{(N, \nu) \mid N \in \text{Gr}_S(M), \nu \in \text{Den}_\times(N)\},$$

that is, the space of all submanifolds of type S in M , decorated with a nowhere zero density. We equip this space with the structure of a Fréchet manifold by declaring the natural bijection

$$(2.3) \quad \text{Gr}_S^{\text{wt}}(M) = \text{Emb}_S(M) \times_{\text{Diff}(S)} \text{Den}_\times(S), \quad (\varphi(S), \varphi_*\mu) \leftrightarrow [\varphi, \mu],$$

to be a diffeomorphism. Here, the right-hand side denotes the total space of the bundle associated with the nonlinear frame bundle in (2.1) and the natural $\text{Diff}(S)$ action on $\text{Den}_\times(S)$. In particular, the canonical forgetful map

$$(2.4) \quad \text{Gr}_S^{\text{wt}}(M) \rightarrow \text{Gr}_S(M), \quad (N, \nu) \mapsto N,$$

becomes a locally trivial smooth bundle with typical fiber $\text{Den}_\times(S)$. Indeed, it corresponds to the bundle projection of the associated bundle $\text{Emb}_S(M) \times_{\text{Diff}(S)} \text{Den}_\times(S) \rightarrow \text{Gr}_S(M)$ via the identification in (2.3).

There is a canonical $\text{Diff}(M)$ equivariant map

$$J: \text{Gr}_S^{\text{wt}}(M) \rightarrow C^\infty(M)^*, \quad \langle J(N, \nu), f \rangle := \int_N f \nu.$$

This map is injective, and its image consists of compactly supported distributions with mild singularities: $J(N, \nu)$ is supported on N , and its wave front set coincides with the conormal bundle of N .

Let $\mu \in \text{Den}_\times(S)$ be a volume density. The space

$$\text{Gr}_{S,\mu}^{\text{wt}}(M) := \{(N, \nu) \in \text{Gr}_S^{\text{wt}}(M) \mid (S, \mu) \cong (N, \nu)\}$$

is called the *nonlinear Grassmannian of weighted submanifolds of type (S, μ) in M* . It consists of all weighted submanifolds (N, ν) in M such that there exists a diffeomorphism $S \rightarrow N$ taking μ to ν . Denoting the $\text{Diff}(S)$ orbit of μ by $\text{Den}(S)_\mu$, the identification in (2.3) restricts to a canonical bijection

$$(2.5) \quad \text{Gr}_{S,\mu}^{\text{wt}}(M) = \text{Emb}_S(M) \times_{\text{Diff}(S)} \text{Den}(S)_\mu.$$

It is well known [22] that the $\text{Diff}(S)_0$ orbit of μ is a convex subset that consists of all volume densities on S that represent the same cohomology class as μ in $H^k(S; \mathcal{O}_S)$, the de Rham cohomology with coefficients in the orientation bundle. Hence, the $\text{Diff}(S)$

orbit of μ coincides with the set of all volume densities on S that are in the preimage of $H^k(S; \mathcal{O}_S)_{[\mu]}$, the (finite) $\text{Diff}(S)$ orbit of $[\mu]$ in $H^k(S; \mathcal{O}_S)$, under the $\text{Diff}(S)$ equivariant linear map

$$(2.6) \quad h_S : \text{Den}(S) \rightarrow H^k(S; \mathcal{O}_S), \quad h_S(\alpha) = [\alpha].$$

More succinctly,

$$(2.7) \quad \text{Den}(S)_\mu = \text{Den}_\times(S) \cap h_S^{-1}(H^k(S; \mathcal{O}_S)_{[\mu]}).$$

Hence, $\text{Den}(S)_\mu$ is an open subset in a finite union of parallel closed affine subspaces with finite codimension. In particular, $\text{Den}(S)_\mu$ is a splitting smooth submanifold in $\text{Den}_\times(S)$ with finite codimension $\dim H^k(S; \mathcal{O}_S)$ and with tangent spaces

$$(2.8) \quad T_\alpha \text{Den}(S)_\mu = \ker h_S = d\Omega^{k-1}(S; \mathcal{O}_S).$$

Using (2.5), we conclude that $\text{Gr}_{S,\mu}^{\text{wt}}(M)$ is a splitting smooth submanifold in $\text{Gr}_S^{\text{wt}}(M)$ with finite codimension $\dim H^k(S; \mathcal{O}_S)$. Moreover, the canonical forgetful map in (2.4) restricts to a locally trivial smooth fiber bundle $\text{Gr}_{S,\mu}^{\text{wt}}(M) \rightarrow \text{Gr}_S(M)$ with typical fiber $\text{Den}(S)_\mu$.

The space of *homologically weighted submanifolds of type S in M* is defined as

$$\text{Gr}_S^{\text{hwt}}(M) := \{(N, [\nu]) : N \in \text{Gr}_S(M), [\nu] \in H^k(N; \mathcal{O}_N)\}.$$

Using the canonical bijection

$$(2.9) \quad \text{Gr}_S^{\text{hwt}}(M) = \text{Emb}_S(M) \times_{\text{Diff}(S)} H^k(S; \mathcal{O}_S),$$

we turn $\text{Gr}_S^{\text{hwt}}(M)$ into a smooth vector bundle of finite rank $\dim H^k(S; \mathcal{O}_S)$ over $\text{Gr}_S(M)$. The canonical $\text{Diff}(M)$ equivariant map

$$h_{\text{Gr}_S(M)} : \text{Gr}_S^{\text{wt}}(M) \rightarrow \text{Gr}_S^{\text{hwt}}(M), \quad (N, \nu) \mapsto (N, [\nu]),$$

is a smooth bundle map over $\text{Gr}_S(M)$. Indeed, via the diffeomorphisms in (2.5) and (2.9), it corresponds to the map induced by (2.6).

The space of *homologically weighted submanifolds of type $(S, [\mu])$ in M* is defined by

$$\text{Gr}_{S,[\mu]}^{\text{hwt}}(M) := \{(N, [\nu]) \in \text{Gr}_S^{\text{hwt}}(M) : (N, [\nu]) \cong (S, [\mu])\}$$

and consists of all homologically weighted submanifolds $(N, [\nu])$ such that there exists a diffeomorphism $S \rightarrow N$ taking the cohomology class $[\mu]$ to $[\nu]$. As (2.9) restricts to a bijection

$$\text{Gr}_{S,[\mu]}^{\text{hwt}}(M) = \text{Emb}_S(M) \times_{\text{Diff}(S)} H^k(S; \mathcal{O}_S)_{[\mu]},$$

we see that $\text{Gr}_{S,[\mu]}^{\text{hwt}}(M)$ is a finite covering of $\text{Gr}_S(M)$. Using (2.7), we conclude

$$(2.10) \quad \text{Gr}_{S,\mu}^{\text{wt}}(M) = h_{\text{Gr}_S(M)}^{-1}(\text{Gr}_{S,[\mu]}^{\text{hwt}}(M)).$$

It is well known that the $\text{Diff}_c(M)$ action on $\text{Emb}_S(M)$ admits local smooth sections (see, for instance, [9, Lemma 2.1(c)]). Furthermore, the (transitive) $\text{Diff}(S)$ action on $\text{Den}(S)_\mu$ also admits local smooth sections. The latter can be shown using

Moser's method of proof in [22, Section 4] (see Lemma 2.12). Using Lemma A.1 in the Appendix, we conclude that the natural $\text{Diff}_c(M)$ action on $\text{Gr}_{S,\mu}^{\text{wt}}(M)$ admits local smooth sections. In particular, this action is locally transitive. Hence, each connected component of $\text{Gr}_{S,\mu}^{\text{wt}}(M)$ is a $\text{Diff}_c(M)_0$ orbit. Consequently, each $\text{Diff}_c(M)$ or $\text{Diff}(M)$ orbit in $\text{Gr}_{S,\mu}^{\text{wt}}(M)$ is a union of connected components.

Remark 2.1 Poincaré duality provides a canonical $\text{Diff}(S)$ equivariant isomorphism

$$H^k(S; \mathcal{O}_S) = H_0(S; \mathbb{R}).$$

Hence, specifying a cohomology class $[\mu] \in H^k(S; \mathcal{O}_S)$ amounts to specifying the total volume of μ on each connected component of S .

Example 2.2 If S is connected, then $H^k(S; \mathcal{O}_S) = \mathbb{R}$ and the $\text{Diff}(S)$ action is trivial on this cohomology. Hence, the orbit $H^k(S; \mathcal{O}_S)_{[\mu]}$ is a one-point set, and

$$(2.11) \quad \text{Den}(S)_\mu = \{ \alpha \in \text{Den}_\times(S) : \int_S \alpha = \int_S \mu \}$$

is connected. Correspondingly,

$$\text{Gr}_{S,\mu}^{\text{wt}}(M) = \{ (N, \nu) \in \text{Gr}_S^{\text{wt}}(M) : \int_N \nu = \int_S \mu \}.$$

This is the case considered in [7, 15, 26].

If S is built out of two diffeomorphic connected components, then $H^k(S; \mathcal{O}_S) \cong \mathbb{R}^2$ and any diffeomorphism swapping the two connected components acts nontrivially on this cohomology. If μ has equal total volume on the two connected components, then the orbit $H^k(S; \mathcal{O}_S)_{[\mu]}$ is a one-point set and $\text{Den}(S)_\mu$ is connected. Otherwise, $H^k(S; \mathcal{O}_S)_{[\mu]}$ consists of two points and, by (2.7), $\text{Den}(S)_\mu$ has two connected components.

Remark 2.3 Suppose $\mu \in \text{Den}_\times(S)$. It is well known that $\text{Diff}(S, \mu)$, the group of diffeomorphisms preserving μ , is a splitting Lie subgroup in $\text{Diff}(S)$ (see [11, Theorem III.2.5.3 on page 203]). Moreover, the map provided by the action, $\text{Diff}(S) \rightarrow \text{Den}(S)_\mu, f \mapsto f_*\mu$, is a smooth principal bundle with structure group $\text{Diff}(S, \mu)$. Via (2.5), this implies that the surjective and $\text{Diff}(M)$ equivariant map

$$\text{Emb}_S(M) \rightarrow \text{Gr}_{S,\mu}^{\text{wt}}(M), \quad \varphi \mapsto (\varphi(S), \varphi_*\mu),$$

is smooth principal bundle with structure group $\text{Diff}(S, \mu)$.

Remark 2.4 Suppose $(N, \nu) \in \text{Gr}_S^{\text{wt}}(M)$ and let $\text{Gr}_S^{\text{wt}}(M)_{(N,\nu)}$ denote its $\text{Diff}_c(M)$ orbit. Combining the preceding remark with the fact that the $\text{Diff}_c(M)$ action on $\text{Emb}_S(M)$ admits local smooth sections [9, Lemma 2.1(c)], we see that the map provided by the action,

$$\text{Diff}_c(M) \rightarrow \text{Gr}_S^{\text{wt}}(M)_{(N,\nu)}, \quad f \mapsto (f(N), f_*\nu),$$

is a smooth principal bundle with structure group $\text{Diff}_c(M, N, \nu)$, the group of diffeomorphisms preserving N and ν . The latter is a splitting Lie subgroup in $\text{Diff}_c(M)$, for it coincides with the preimage of $\text{Diff}(N, \nu)$ under the canonical bundle projection

$\text{Diff}_c(M, N) \rightarrow \text{Diff}(N)$ (see [9, Lemma 2.1(d)]). Hence, each orbit may be regarded as a homogeneous space,

$$\text{Gr}_S^{\text{wt}}(M)_{(N, \nu)} = \text{Diff}_c(M) / \text{Diff}_c(M, N, \nu).$$

2.2 Weighted nonlinear flag manifolds

Fix natural numbers k_i such that

$$(2.12) \quad 0 \leq k_1 < k_2 < \dots < k_r$$

and let $\mathcal{S} = (S_1, \dots, S_r)$ be a collection of closed smooth manifolds with $\dim S_i = k_i$.

For a smooth manifold M , we let

$$\text{Flag}_{\mathcal{S}}(M) := \left\{ (N_1, \dots, N_r) \in \prod_{i=1}^r \text{Gr}_{S_i}(M) \mid \forall i : N_i \subseteq N_{i+1} \right\}$$

denote the space of *nonlinear flags of type \mathcal{S} in M* , and we write

$$\text{Fr}_{\mathcal{S}}(M) := \left\{ (\varphi_1, \dots, \varphi_r) \in \prod_{i=1}^r \text{Emb}_{S_i}(M) \mid \forall i : \varphi_i(S_i) \subseteq \varphi_{i+1}(S_{i+1}) \right\}$$

for the space of *nonlinear frames of type \mathcal{S} in M* . In [9, Proposition 2.3], it has been shown that $\text{Flag}_{\mathcal{S}}(M)$ and $\text{Fr}_{\mathcal{S}}(M)$ are splitting smooth submanifolds of $\prod_{i=1}^r \text{Gr}_{S_i}(M)$ and $\prod_{i=1}^r \text{Emb}_{S_i}(M)$, respectively. Moreover, the canonical $\text{Diff}(M)$ equivariant map

$$(2.13) \quad \text{Fr}_{\mathcal{S}}(M) \rightarrow \text{Flag}_{\mathcal{S}}(M), \quad (\varphi_1, \dots, \varphi_r) \mapsto (\varphi_1(S_1), \dots, \varphi_r(S_r)),$$

is a smooth principal fiber bundle with structure group

$$\text{Diff}(\mathcal{S}) := \prod_{i=1}^r \text{Diff}(S_i).$$

We denote the space of *weighted nonlinear flags of type \mathcal{S} in M* by

$$(2.14) \quad \text{Flag}_{\mathcal{S}}^{\text{wt}}(M) := \left\{ ((N_1, \nu_1), \dots, (N_r, \nu_r)) \in \prod_{i=1}^r \text{Gr}_{S_i}^{\text{wt}}(M) \mid \forall i : N_i \subseteq N_{i+1} \right\}.$$

This is a splitting smooth submanifold in $\prod_{i=1}^r \text{Gr}_{S_i}^{\text{wt}}(M)$, for it coincides with the preimage of the splitting smooth submanifold $\text{Flag}_{\mathcal{S}}(M)$ under the bundle projection $\prod_{i=1}^r \text{Gr}_{S_i}^{\text{wt}}(M) \rightarrow \prod_{i=1}^r \text{Gr}_{S_i}(M)$. Moreover, the canonical $\text{Diff}(M)$ equivariant forgetful map

$$(2.15) \quad \text{Flag}_{\mathcal{S}}^{\text{wt}}(M) \rightarrow \text{Flag}_{\mathcal{S}}(M), \quad ((N_1, \nu_1), \dots, (N_r, \nu_r)) \mapsto (N_1, \dots, N_r),$$

is a smooth fiber bundle with typical fiber

$$(2.16) \quad \text{Den}_{\times}(\mathcal{S}) := \prod_{i=1}^r \text{Den}_{\times}(S_i).$$

The latter is a $\text{Diff}(\mathcal{S})$ invariant open subset in the Fréchet space $\text{Den}(\mathcal{S}) := \prod_{i=1}^r \text{Den}(S_i)$. Furthermore, the canonical $\text{Diff}(M)$ equivariant bijection

$$(2.17) \quad \text{Flag}_{\mathcal{S}}^{\text{wt}}(M) = \text{Fr}_{\mathcal{S}}(M) \times_{\text{Diff}(\mathcal{S})} \text{Den}_{\times}(\mathcal{S}),$$

$$\left((\varphi_1(S_1), (\varphi_1)_* \mu_1), \dots, (\varphi_r(S_r), (\varphi_r)_* \mu_r) \right) \leftrightarrow [(\varphi_1, \dots, \varphi_r), (\mu_1, \dots, \mu_r)],$$

is a diffeomorphism between $\text{Flag}_{\mathcal{S}}^{\text{wt}}(M)$ and the bundle associated with the nonlinear frame bundle in (2.13) and the canonical $\text{Diff}(\mathcal{S})$ action on $\text{Den}_{\times}(\mathcal{S})$. Indeed, this is just the bundle diffeomorphism $\prod_{i=1}^r \text{Gr}_{S_i}^{\text{wt}}(M) = \prod_{i=1}^r \text{Emb}_{S_i}(M) \times_{\text{Diff}(S_i)} \text{Den}_{\times}(S_i)$ obtained by taking the product of the diffeomorphisms in (2.3), restricted over the submanifold $\text{Flag}_{\mathcal{S}}(M)$ in its base $\prod_{i=1}^r \text{Gr}_{S_i}(M)$.

We have a canonical $\text{Diff}(M)$ equivariant map

$$(2.18) \quad J : \text{Flag}_{\mathcal{S}}^{\text{wt}}(M) \rightarrow C^{\infty}(M)^*, \quad \langle J((N_1, \nu_1), \dots, (N_r, \nu_r)), f \rangle := \sum_{i=1}^r \int_{N_i} f \nu_i.$$

The image of J consists of compactly supported distributions on M with mild singularities. More precisely, the wave front set of $J((N_1, \nu_1), \dots, (N_r, \nu_r))$ coincides with the union of the conormal bundles of N_1, \dots, N_r .

Lemma 2.5 *The map in (2.18) is injective.*

Proof Suppose $J((N_1, \nu_1), \dots, (N_r, \nu_r)) = J((N'_1, \nu'_1), \dots, (N'_r, \nu'_r))$. Proceeding by induction on r , it suffices to show $N_r = N'_r$ and $\nu_r = \nu'_r$.

To show $N_r = N'_r$, we assume by contradiction that there exists $x \in N_r$ with $x \notin N'_r$. Using (2.12), we see that $N_r \setminus N_{r-1}$ is dense in N_r . Thus, we may w.l.o.g. assume $x \notin N_{r-1}$. Moreover, $\nu_r(x) \neq 0$ as ν_r does not vanish on N_r . Hence, if f is a smooth bump function supported on a sufficiently small neighborhood of x , then $\langle J((N_1, \nu_1), \dots, (N_r, \nu_r)), f \rangle = \int_{N_r} f \nu_r \neq 0$ and $\langle J((N'_1, \nu'_1), \dots, (N'_r, \nu'_r)), f \rangle = 0$. Since this contradicts our assumption, we conclude $N_r = N'_r$.

To show $\nu_r = \nu'_r$, we assume by contradiction that there exists $x \in N_r = N'_r$ with $\nu_r(x) \neq \nu'_r(x)$. As before, we may w.l.o.g. assume $x \notin N_{r-1}$ and $x \notin N'_{r-1}$. Hence, if f is a smooth bump function supported in a sufficiently small neighborhood of x , then $\langle J((N_1, \nu_1), \dots, (N_r, \nu_r)), f \rangle = \int_{N_r} f \nu_r \neq \int_{N'_r} f \nu'_r = \langle J((N'_1, \nu'_1), \dots, (N'_r, \nu'_r)), f \rangle$. Since this contradicts our assumption, we conclude $\nu_r = \nu'_r$. ■

Remark 2.6 Suppose ω is a symplectic form on M , and let $\text{Flag}_{\mathcal{S}}^{\text{symp}}(M)$ denote the manifold of symplectic nonlinear flags of type \mathcal{S} (cf. [9, Section 4.2]). Recall that this is the open subset consisting of all flags $(N_1, \dots, N_r) \in \text{Flag}_{\mathcal{S}}(M)$ such that ω restricts to a symplectic form on each N_i . Hence, k_i must be even and $\omega^{k_i/2}$ pulls back to a volume form on N_i which in turn gives rise to a volume density $\nu_i = |i_{N_i}^* \omega^{k_i/2}|$ on N_i . Consequently, the symplectic form ω provides a $\text{Symp}(M, \omega)$ equivariant injective smooth map (section)

$$(2.19) \quad \text{Flag}_{\mathcal{S}}^{\text{symp}}(M) \rightarrow \text{Flag}_{\mathcal{S}}^{\text{wt}}(M)$$

which is right inverse to the restriction of the canonical bundle projection in (2.15). Composing the map in (2.19) with J in (2.18), we obtain the moment map considered in [9, equation (38)].

Remark 2.7 A Riemannian metric g on M induces a volume density on every submanifold of M . Hence, g provides a smooth section $\text{Flag}_S(M) \rightarrow \text{Flag}_S^{\text{wt}}(M)$ of the canonical bundle projection in (2.15), which is $\text{Isom}(M, g)$ equivariant (cf. [26, Section 6]).

2.3 Reduction of structure group

It will be convenient to use a reduction of the structure group for the principal frame bundle in (2.13). To this end, we fix embeddings $\iota_i: S_i \rightarrow S_{i+1}$ and put $\iota = (\iota_1, \dots, \iota_{r-1})$.

We begin by recalling some facts from [9, Proposition 2.10]. The space of *nonlinear flags of type (S, ι)* in M ,

$$\text{Flag}_{S,\iota}(M) := \left\{ (N_1, \dots, N_r) \in \text{Flag}_S(M) \mid \begin{array}{l} (S_1 \xrightarrow{\iota_1} S_2 \xrightarrow{\iota_2} \dots \xrightarrow{\iota_{r-1}} S_r) \\ \cong (N_1 \subseteq N_2 \subseteq \dots \subseteq N_r) \end{array} \right\},$$

consists of all nonlinear flags (N_1, \dots, N_r) in M such that there exist diffeomorphisms $S_i \rightarrow N_i$, $1 \leq i \leq r$ intertwining ι_i with the canonical inclusion $N_i \subseteq N_{i+1}$. This is a $\text{Diff}(M)$ invariant open and closed subset in $\text{Flag}_S(M)$. The space of *parametrized nonlinear flags (nonlinear frames) of type (S, ι)* in M ,

$$\text{Fr}_{S,\iota}(M) := \{ (\varphi_1, \dots, \varphi_r) \in \text{Fr}_S(M) \mid \forall i: \varphi_i = \varphi_{i+1} \circ \iota_i \},$$

is a splitting smooth submanifold of $\text{Fr}_S(M)$. Moreover, the map $\text{Fr}_{S,\iota}(M) \rightarrow \text{Flag}_{S,\iota}(M)$ obtained by restriction of (2.13) is a smooth principal bundle with structure group

$$(2.20) \quad \text{Diff}(S; \iota) := \left\{ (g_1, \dots, g_r) \in \prod_{i=1}^r \text{Diff}(S_i) \mid \forall i: g_{i+1} \circ \iota_i = \iota_i \circ g_i \right\}.$$

The latter is a splitting Lie subgroup in $\text{Diff}(S)$ with Lie algebra

$$(2.21) \quad \mathfrak{X}(S; \iota) = \left\{ (Z_1, \dots, Z_r) \in \prod_{i=1}^r \mathfrak{X}(S_i) \mid \forall i: Z_{i+1} \circ \iota_i = T\iota_i \circ Z_i \right\}.$$

We obtain a $\text{Diff}(M)$ equivariant commutative diagram

$$(2.22) \quad \begin{array}{ccc} \text{Fr}_{S,\iota}(M) & \hookrightarrow & \text{Fr}_S(M) \\ \text{Diff}(S; \iota) \downarrow & & \downarrow \text{Diff}(S) \\ \text{Flag}_{S,\iota}(M) & \hookrightarrow & \text{Flag}_S(M), \end{array}$$

which may be regarded as a reduction of the structure group for (2.13) along the inclusion $\text{Diff}(S; \iota) \subseteq \text{Diff}(S)$ over $\text{Flag}_{S,\iota}(M)$ (see [9, Proposition 2.10] for more details).

Remark 2.8 The $\text{Diff}(M)$ equivariant bijection

$$(2.23) \quad \text{Fr}_{S_r, \iota}(M) = \text{Emb}_{S_r}(M), \quad (\varphi_1, \dots, \varphi_r) \mapsto \varphi_r,$$

is a diffeomorphism [9, Proposition 2.10(b)]. Correspondingly, we have a group isomorphism

$$(2.24) \quad \text{Diff}(\mathcal{S}; \iota) = \text{Diff}(S_r; \Sigma), \quad (g_1, \dots, g_r) \mapsto g_r,$$

where $\text{Diff}(S_r; \Sigma)$ denotes the subgroup of all diffeomorphisms of S_r preserving the nonlinear flag $\Sigma = (\Sigma_1, \dots, \Sigma_{r-1})$ in S_r , where $\Sigma_i := (\iota_{r-1} \circ \dots \circ \iota_i)(S_i)$. The latter is a splitting Lie subgroup of $\text{Diff}(S_r)$ (see [9, Proposition 2.9(b)]), and (2.24) is a diffeomorphism of Lie groups [9, Proposition 2.10(a)]. The Lie algebra of $\text{Diff}(\mathcal{S}; \iota)$ can be identified in a similar way with $\mathfrak{X}(S_r; \Sigma)$, the Lie algebra of vector fields on S_r that are tangent to $\Sigma_1, \dots, \Sigma_{r-1}$.

We are interested in the reduction of structure group (2.22) because the $\text{Diff}_c(M)$ action on $\text{Fr}_{S_r, \iota}(M)$ admits local smooth sections. This follows from [9, Lemma 2.1(c)] and the diffeomorphism in (2.23).

Let $\text{Flag}_{S_r, \iota}^{\text{wt}}(M)$ denote the preimage of $\text{Flag}_{S_r, \iota}(M)$ under the bundle projection in (2.15). Restricting the diffeomorphism in (2.17) over $\text{Flag}_{S_r, \iota}(M)$ and combining this with the diffeomorphism in (2.23), we obtain a $\text{Diff}(M)$ equivariant diffeomorphism of bundles over $\text{Flag}_{S_r, \iota}(M)$,

$$(2.25) \quad \text{Flag}_{S_r, \iota}^{\text{wt}}(M) = \text{Emb}_{S_r}(M) \times_{\text{Diff}(S_r, \iota)} \text{Den}_\times(S).$$

2.4 The $\text{Diff}(M)$ action on the space of weighted nonlinear flags

In this section, we aim at describing the $\text{Diff}(M)$ orbits in $\text{Flag}_{S_r}^{\text{wt}}(M)$ (see Theorem 2.11).

Let $\iota = (\iota_1, \dots, \iota_{r-1})$ be a collection of embeddings $\iota_i : S_i \rightarrow S_{i+1}$ and suppose $\mu = (\mu_1, \dots, \mu_r) \in \text{Den}_\times(S)$. We define the space of *weighted flags of type (S, ι, μ)* in M by

$$\begin{aligned} & \text{Flag}_{S, \iota, \mu}^{\text{wt}}(M) \\ & := \left\{ \left((N_1, \nu_1), \dots, (N_r, \nu_r) \right) \in \text{Flag}_S^{\text{wt}}(M) \left| \begin{array}{l} (S_1 \xrightarrow{\iota_1} \dots \rightarrow S_r, \mu_1, \dots, \mu_r) \\ \cong (N_1 \subseteq \dots \subseteq N_r, \nu_1, \dots, \nu_r) \end{array} \right. \right\}, \end{aligned}$$

that is, the space of all weighted flags $((N_1, \nu_1), \dots, (N_r, \nu_r))$ in M such that there exist diffeomorphisms $S_i \rightarrow N_i$, $1 \leq i \leq r$, intertwining ι_i with the canonical inclusion $N_i \subseteq N_{i+1}$, and taking μ_i to ν_i .

Denoting the $\text{Diff}(S, \iota)$ orbit of μ by $\text{Den}(S)_{\iota, \mu}$, the diffeomorphism in (2.25) restricts to a $\text{Diff}(M)$ equivariant bijection

$$(2.26) \quad \text{Flag}_{S, \iota, \mu}^{\text{wt}}(M) = \text{Emb}_{S_r}(M) \times_{\text{Diff}(S, \iota)} \text{Den}(S)_{\iota, \mu}.$$

Consider the finite-dimensional vector space

$$(2.27) \quad H(S, \iota) := \prod_{i=1}^r H^{k_i}(S_i, \iota_{i-1}(S_{i-1}); \mathcal{O}_{S_i}) = \prod_{i=1}^r H_0(S_i \setminus \iota_{i-1}(S_{i-1}); \mathbb{R}).$$

Here, the left-hand side denotes relative de Rham cohomology with coefficients in the orientation bundle, and we are using the convention $S_0 = \emptyset$. The $\text{Diff}(\mathcal{S}, \iota)$ equivariant identification on the right-hand side indicates Poincaré–Lefschetz duality. We have a $\text{Diff}(\mathcal{S}, \iota)$ equivariant linear map

$$(2.28) \quad h_{\mathcal{S}, \iota}: \text{Den}(\mathcal{S}) \rightarrow H(\mathcal{S}, \iota), \quad h_{\mathcal{S}, \iota}(\mu_1, \dots, \mu_r) := ([\mu_1], \dots, [\mu_r]).$$

Pinning down the class $[\mu] := h_{\mathcal{S}, \iota}(\mu)$ thus amounts to specifying the integrals of μ_i over each connected component of $S_i \setminus \iota_{i-1}(S_{i-1})$ for $i = 1, \dots, r$.

Remark 2.9 (Large codimensions) If the codimensions $\dim(S_i) - \dim(S_{i-1})$ are all strictly larger than one, then $H^{k_i}(S_i, \iota_{i-1}(S_{i-1}); \mathcal{O}_{S_i}) = H^{k_i}(S_i; \mathcal{O}_{S_i}) = H_0(S_i; \mathbb{R})$, and

$$H(\mathcal{S}, \iota) = \prod_{i=1}^r H^{k_i}(S_i; \mathcal{O}_{S_i}) = \prod_{i=1}^r H_0(S_i; \mathbb{R}).$$

Hence, in this case, the cohomology space $H(\mathcal{S}, \iota)$ does not depend on the embeddings ι .

Proposition 2.10 *In this situation, the following hold true:*

- (a) *The $\text{Diff}(\mathcal{S}, \iota)_0$ orbit of μ coincides with the convex set $\text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S}, \iota}^{-1}([\mu])$. In particular, this orbit is a splitting smooth submanifold in $\text{Den}_\times(\mathcal{S})$ with finite codimension $\dim H(\mathcal{S}, \iota)$.*
- (b) *The $\text{Diff}(\mathcal{S}, \iota)_0$ action on $\text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S}, \iota}^{-1}([\mu])$ admits local smooth sections.*
- (c) *Denoting the (finite) $\text{Diff}(\mathcal{S}; \iota)$ orbit of $[\mu]$ by $H(\mathcal{S}, \iota)_{[\mu]}$, the $\text{Diff}(\mathcal{S}; \iota)$ orbit of μ is*

$$(2.29) \quad \text{Den}(\mathcal{S})_{\iota, \mu} = \text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S}, \iota}^{-1}(H(\mathcal{S}, \iota)_{[\mu]}).$$

In particular, $\text{Den}(\mathcal{S})_{\iota, \mu}$ is a splitting smooth submanifold in $\text{Den}_\times(\mathcal{S})$ with finite codimension $\dim H(\mathcal{S}, \iota)$ and with tangent spaces

$$(2.30) \quad T_\alpha \text{Den}(\mathcal{S})_{\iota, \mu} = \ker h_{\mathcal{S}, \iota} = \{(d\gamma_1, \dots, d\gamma_r) : \gamma_i \in \Omega^{k_i-1}(S_i; \mathcal{O}_{S_i}), \iota_{i-1}^* \gamma_i = 0\}.$$

Moreover, the $\text{Diff}(\mathcal{S}, \iota)$ action on $\text{Den}(\mathcal{S})_{\iota, \mu}$ admits local smooth sections.

- (d) *The canonical inclusion $\text{Den}(\mathcal{S})_{\iota, \mu} \subseteq \prod_{i=1}^r \text{Den}(S_i)_{\mu_i}$ is a splitting smooth submanifold of finite codimension.*

We postpone the proof of this proposition and proceed with the main result in this section.

Theorem 2.11 *In this situation, the following hold true:*

- (a) *The space $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt}}(M)$ is a splitting smooth submanifold in $\text{Flag}_{\mathcal{S}, \iota}^{\text{wt}}(M)$ with finite codimension $\dim H(\mathcal{S}, \iota)$.*
- (b) *The canonical $\text{Diff}(M)$ equivariant forgetful map $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt}}(M) \rightarrow \text{Flag}_{\mathcal{S}, \iota}^{\text{wt}}(M)$ is a locally trivial smooth fiber bundle with typical fiber $\text{Den}(\mathcal{S})_{\iota, \mu}$.*
- (c) *The canonical inclusion $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt}}(M) \subseteq \prod_{i=1}^r \text{Gr}_{S_i, \mu_i}^{\text{wt}}(M)$ is a splitting smooth submanifold.*

(d) *The $\text{Diff}_c(M)$ action on $\text{Flag}_{S,t,\mu}^{\text{wt}}(M)$ admits local smooth sections. In particular, each connected component of $\text{Flag}_{S,t,\mu}^{\text{wt}}(M)$ is a $\text{Diff}_c(M)_0$ orbit. Furthermore, every $\text{Diff}(M)$ or $\text{Diff}_c(M)$ orbit in $\text{Flag}_{S,t,\mu}^{\text{wt}}(M)$ is a union of connected components.*

Proof Parts (a) and (b) follow by combining (2.25) and (2.26) with Proposition 2.10(c).

Part (c) follows from Proposition 2.10(d) and the reduction of structure groups in (2.22) via the diffeomorphisms in (2.5) and (2.26) (see also (2.23)).

Let us finally turn to part (d). By Proposition 2.10(c), the (transitive) $\text{Diff}(S; \iota)$ action on $\text{Den}(S)_{i,\mu}$ admits local smooth sections. The $\text{Diff}_c(M)$ action on $\text{Emb}_{S_r}(M)$ admits local smooth sections too (cf. [9, Lemma 2.1(c)]). Using Lemma A.1, we conclude that the $\text{Diff}_c(M)$ action on $\text{Flag}_{S,t,\mu}^{\text{wt}}(M)$ admits local smooth sections (cf. (2.25)). ■

We will prove Proposition 2.10 by induction on the depth of the flags, using the following crucial lemma whose proof we postpone.

Lemma 2.12 *Let S be a closed submanifold of N such that $\dim(S) < \dim(N) = n$, and consider the $\text{Diff}(N, S)$ equivariant linear map*

$$h : \text{Den}(N) \rightarrow H^n(N, S; \mathcal{O}_N), \quad h(\mu) = [\mu].$$

Then, for each $\kappa \in H^n(N, S; \mathcal{O}_N)$, the natural $\text{Diff}(N, S)_0$ action on

$$(2.31) \quad \text{Diff}(S) \times (\text{Den}_\times(N) \cap h^{-1}(\kappa))$$

admits local smooth sections.

Proof of Proposition 2.10 We proceed by induction on r using Lemma 2.12. Let us denote the truncated sequences by $S' := (S_1, \dots, S_{r-1})$, $\mu' := (\mu_1, \dots, \mu_{r-1})$, and $\iota' := (\iota_1, \dots, \iota_{r-2})$. By induction, the $\text{Diff}(S', \iota')_0$ action on $\text{Den}_\times(S') \cap h_{S',\iota'}^{-1}([\mu'])$ admits local smooth sections. Hence, there exist an open neighborhood U' of μ' in $\text{Den}_\times(S') \cap h_{S',\iota'}^{-1}([\mu'])$ and a smooth map

$$\text{Den}_\times(S') \cap h_{S',\iota'}^{-1}([\mu']) \supseteq U' \xrightarrow{\sigma'} \text{Diff}(S'; \iota')_0,$$

such that for all $\tilde{\mu}' \in U'$, we have

$$(2.32) \quad (\sigma'(\tilde{\mu}'))_* \mu' = \tilde{\mu}' \quad \text{and} \quad \sigma'(\mu') = \text{id}.$$

Recall that $\text{Diff}(S', \iota')$ is a splitting Lie subgroup in $\text{Diff}(S_{r-1})$ (cf. [9, Propositions 2.9(b) and 2.10(a)]). Using [9, Lemma 2.1(d)], this implies that $\text{Diff}(S, \iota)$ is a splitting Lie subgroup in $\text{Diff}(S_r, \iota_{r-1}(S_{r-1}))$. Hence, restricting a local smooth section as in Lemma 2.12, we see that the $\text{Diff}(S, \iota)_0$ action on $\text{Diff}(S', \iota') \times (\text{Den}_\times(S_r) \cap h^{-1}([\mu_r]))$ admits local smooth sections. In other words, there exist an open neighborhood V of the identity in $\text{Diff}(S'; \iota')$, an open neighborhood U'' of μ_r in $\text{Den}_\times(S_r) \cap h^{-1}([\mu_r])$, and a smooth map

$$\text{Diff}(S'; \iota') \times (\text{Den}_\times(S_r) \cap h^{-1}([\mu_r])) \supseteq V \times U'' \xrightarrow{\sigma''} \text{Diff}(S; \iota)_0,$$

such that for all $g \in V$ and $\tilde{\mu}_r \in U''$, we have

$$(2.33) \quad \sigma''(g, \tilde{\mu}_r) \cdot (\text{id}, \mu_r) = (g, \tilde{\mu}_r) \quad \text{and} \quad \sigma''(\text{id}, \mu_r) = \text{id}.$$

Hence, $U := (\sigma')^{-1}(V) \times U''$ is an open neighborhood of μ in $\text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S},t}^{-1}([\mu])$, and

$$\begin{aligned} \text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S},t}^{-1}([\mu]) &\supseteq U \xrightarrow{\sigma} \text{Diff}(\mathcal{S}; \iota)_0, \\ \sigma(\tilde{\mu}_1, \dots, \tilde{\mu}_r) &:= \sigma''(\sigma'(\tilde{\mu}_1, \dots, \tilde{\mu}_{r-1}), \tilde{\mu}_r) \end{aligned}$$

is a local smooth section for the $\text{Diff}(\mathcal{S}; \iota)_0$ action on $\text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S},t}^{-1}([\mu])$, i.e.,

$$\sigma(\mu) = \text{id} \quad \text{and} \quad \sigma(\tilde{\mu})_* \mu = \tilde{\mu},$$

for all $\tilde{\mu} \in U$. By convexity, $\text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S},t}^{-1}([\mu])$ is connected. Therefore, the $\text{Diff}(\mathcal{S}; \iota)_0$ action is transitive on $\text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S},t}^{-1}([\mu])$. This shows (a) and (b). Part (c) follows immediately.

To see (d), let A denote the preimage of $\prod_{i=1}^r H^{k_i}(S_i; \mathcal{O}_{S_i})_{[\mu_i]}$ under the canonical linear surjection $H(\mathcal{S}, \iota) \rightarrow \prod_{i=1}^r H^{k_i}(S_i; \mathcal{O}_{S_i})$. Hence, A is a finite union of affine subspaces in $H(\mathcal{S}, \iota)$. In view of (2.7), we have $\prod_{i=1}^r \text{Den}(S_i)_{\mu_i} = \text{Den}_\times(\mathcal{S}) \cap h_{\mathcal{S},t}^{-1}(A)$. Combining this with (2.29), we conclude that $\text{Den}(\mathcal{S})_{i,\mu}$ is a splitting smooth submanifold in $\prod_{i=1}^r \text{Den}(S_i)_{\mu_i}$ with finite codimension $\dim A$. ■

Let us next establish the following infinitesimal version of Lemma 2.12.

Lemma 2.13 *Let S be a closed submanifold of N such that $\dim(S) < \dim(N) = n$. Suppose $\mu \in \text{Den}_\times(N)$, $\gamma \in \Omega^{n-1}(N; \mathcal{O}_N) := \{\alpha \in \Omega(N; \mathcal{O}_N) : \iota_S^* \alpha = 0\}$, and $Z \in \mathfrak{X}(S)$. Then there exists a vector field $X \in \mathfrak{X}(N)$ such that $L_X \mu = d\gamma$ and $X|_S = Z$.*

Proof Let $\tilde{Z} \in \mathfrak{X}(N)$ be any extension of Z , i.e., $\tilde{Z}|_S = Z$. Note that $i_{\tilde{Z}} \mu \in \Omega^{n-1}(N; \mathcal{O}_N)$ vanishes when pulled back to S ; hence, the same holds for $\beta := \gamma - i_{\tilde{Z}} \mu \in \Omega^{n-1}(N; \mathcal{O}_N)$.

Let us first construct $\tilde{\gamma} \in \Omega^{n-1}(N; \mathcal{O}_N)$ such that $d\tilde{\gamma} = d\beta$ and $\tilde{\gamma}|_S = 0$. To this end, we fix a smooth homotopy $h: N \times [0, 1] \rightarrow N$ such that $h_1 = \text{id}_N$, $h_t|_S = \text{id}_S$, and such that h_0 maps a neighborhood of S into S . Consider the corresponding chain homotopy $\phi: \Omega^*(N; \mathcal{O}_N) \rightarrow \Omega^{*-1}(N; \mathcal{O}_N)$ defined by $\phi(\alpha) := \int_0^1 \iota_t^* i_{\partial_t} h^* \alpha \, dt$, where $\iota_t: N \rightarrow N \times [0, 1]$ denotes the inclusion at t , that is, $\iota_t(x) := (x, t)$. Then $\phi(\alpha)|_S = 0$ and $d(\phi(\alpha)) + \phi(d\alpha) = h_1^* \alpha - h_0^* \alpha$, for all forms $\alpha \in \Omega^*(N; \mathcal{O}_N)$. In particular, $d\alpha = d(h_0^* \alpha + \phi(d\alpha))$. Defining $\tilde{\gamma} := h_0^* \beta + \phi(d\beta)$, we obtain $d\tilde{\gamma} = d\beta$. Moreover, $h_0^* \beta|_S = 0$ because β vanishes when pulled back to S , and hence $\tilde{\gamma}|_S = 0$, as desired.

Defining a vector field $Y \in \mathfrak{X}(N)$ by $i_Y \mu := \tilde{\gamma}$, we obtain $Y|_S = 0$ and $L_Y \mu = d i_Y \mu = d\tilde{\gamma} = d(\gamma - i_{\tilde{Z}} \mu) = d\gamma - L_{\tilde{Z}} \mu$. Hence, the vector field $X := \tilde{Z} + Y$ has the desired properties. ■

Proof of Lemma 2.12 Recall that the infinitesimal $\text{Diff}(N, S)$ action on $\text{Diff}(S) \times \text{Den}(N)$ is

$$\zeta_X(f, \mu) = (R_{X|_S}(f), -L_X \mu),$$

where $X \in \mathfrak{X}(N, S)$, $f \in \text{Diff}(S)$, and $\mu \in \text{Den}(S)$. Here, for $Z \in T_{\text{id}} \text{Diff}(S) = \mathfrak{X}(S)$, we let $R_Z(f)$ denote the right invariant vector field on $\text{Diff}(S)$ such that $R_Z(\text{id}) = Z$.

Note that the vector field \tilde{Z} in the proof of Lemma 2.13 can be chosen to depend smoothly (and linearly) on Z . Hence, the proof of said lemma actually provides a smooth map

$$\tilde{\sigma} : T_{\text{id}} \text{Diff}(S) \times T(\text{Den}_\times(N) \cap h^{-1}(\kappa)) \rightarrow \mathfrak{X}(N, S)$$

such that $\zeta_{\tilde{\sigma}(Z, d\gamma)}(\text{id}, \mu) = (Z, d\gamma)$ for all $Z \in \mathfrak{X}(S) = T_{\text{id}} \text{Diff}(S)$, $\mu \in \text{Den}_\times(N) \cap h^{-1}(\kappa)$, and $d\gamma \in d\Omega^{n-1}(N, S; \mathcal{O}_N) = T_\mu(\text{Den}_\times(N) \cap h^{-1}(\kappa))$. Combining this with the right trivialization of $T \text{Diff}(S)$, we obtain a smooth map

$$\sigma : T(\text{Diff}(S) \times (\text{Den}_\times(N) \cap h^{-1}(\kappa))) \rightarrow \mathfrak{X}(N, S)$$

such that $\zeta_{\sigma(Z, \xi)}(f, \mu) = (Z, \xi)$ for all $f \in \text{Diff}(S)$, $Z \in T_f \text{Diff}(S)$, $\mu \in \text{Den}_\times(N) \cap h^{-1}(\kappa)$, and $\xi \in T_\mu(\text{Den}_\times(N) \cap h^{-1}(\kappa))$. As $\text{Diff}(N, S)$ is a regular Lie group, we may apply Lemma A.2 to conclude that the $\text{Diff}(N, S)_0$ action on (2.31) admits local smooth sections. ■

This completes the proof of Theorem 2.11.

Remark 2.14 In view of Remark 2.3, we expect that the isotropy subgroup

$$(2.34) \quad \text{Diff}(\mathcal{S}; \iota, \mu) := \{(g_1, \dots, g_r) \in \text{Diff}(\mathcal{S}, \iota) \mid \forall i : g_i^* \mu_i = \mu_i\}$$

is a splitting Lie subgroup of $\text{Diff}(\mathcal{S}; \iota)$ with Lie algebra

$$(2.35) \quad \mathfrak{X}(\mathcal{S}, \iota, \mu) = \{(Z_1, \dots, Z_r) \in \mathfrak{X}(\mathcal{S}, \iota) \mid \forall i : L_{Z_i} \mu_i = 0\},$$

and the surjective map provided by the action, $\text{Diff}(\mathcal{S}, \iota) \rightarrow \text{Den}(\mathcal{S})_{\iota, \mu}$, is a locally trivial smooth principal fiber bundle with structure group $\text{Diff}(\mathcal{S}; \iota, \mu)$. This would follow in a rather straightforward manner, via induction on the depth of the flags, if one could show that the isotropy group $\{f \in \text{Diff}(N, S) : f|_S = \text{id}, f^* \mu = \mu\}$ is a splitting Lie subgroup in $\text{Diff}(N, S)$, whenever S is a closed submanifold of N and μ is a volume density on N . The proof in [11, Theorem III.2.5.3 on page 203] covers the case $S = \emptyset$. However, the adaptation of said proof to nontrivial S is not entirely straightforward, and we will not attempt to prove this here. Note that via the diffeomorphism in (2.24), the group $\text{Diff}(\mathcal{S}; \iota, \mu)$ corresponds to the subgroup of $\text{Diff}(S_r; \Sigma)$ consisting of all diffeomorphisms that preserve μ_r and whose restriction to Σ_i preserves $(\iota_{r-1} \circ \dots \circ \iota_i)_* \mu_i$, for $1 \leq i \leq r-1$. Similarly, the Lie algebra $\mathfrak{X}(\mathcal{S}; \iota, \mu)$ can be identified to the corresponding subalgebra of $\mathfrak{X}(S_r; \Sigma)$.

If the expectation formulated in the preceding paragraph were indeed true, then the surjective and $\text{Diff}(M)$ equivariant map

$$(2.36) \quad \text{Emb}_{S_r}(M) = \text{Fr}_{S, \iota}(M) \rightarrow \text{Flag}_{S, \iota, \mu}^{\text{wt}}(M),$$

$$(2.37) \quad (\varphi_1, \dots, \varphi_r) \mapsto \left((\varphi_1(S_1), (\varphi_1)_* \mu_1), \dots, (\varphi_r(S_r), (\varphi_r)_* \mu_r) \right),$$

would be a locally trivial smooth principal fiber bundle with structure group $\text{Diff}(\mathcal{S}; \iota, \mu)$ (see (2.26)). Moreover, generalizing Remark 2.4, the isotropy group of a weighted flag (\mathcal{N}, ν) ,

$$\text{Diff}_c(M; \mathcal{N}, \nu) := \{g \in \text{Diff}_c(M) \mid \forall i : g(N_i) = N_i, g|_{N_i}^* \nu_i = \nu_i\},$$

would be a splitting Lie subgroup of $\text{Diff}_c(M)$, for it coincides with the preimage of $\text{Diff}(S; \iota, \mu)$ under the bundle projection $\text{Diff}_c(M; \mathcal{N}) \rightarrow \text{Diff}(\mathcal{N}, \iota_{\mathcal{N}})$ (cf. [9, Lemma 2.1(d) and Propositions 2.9(b) and 2.10(a)]). Furthermore, the orbit map $\text{Diff}_c(M) \rightarrow \text{Flag}_S^{\text{wt}}(M)_{(\mathcal{N}, \nu)}$ provided by the action would be a locally trivial smooth principal bundle with structure group $\text{Diff}_c(M; \mathcal{N}, \nu)$. Hence, the $\text{Diff}_c(M)$ orbit of (\mathcal{N}, ν) could be regarded as a homogeneous space

$$\text{Flag}_S^{\text{wt}}(M)_{(\mathcal{N}, \nu)} = \text{Diff}_c(M) / \text{Diff}_c(M; \mathcal{N}, \nu).$$

2.5 A homological description

In this section, we give a more explicit description of the manifold $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M)$.

If $\mathcal{N} = (N_1, \dots, N_r)$ is a nonlinear flag of type S in M , we put

$$H(\mathcal{N}) := \prod_{i=1}^r H^{k_i}(N_i, N_{i-1}; \mathcal{O}_{N_i}) = \prod_{i=1}^r H_0(N_i \setminus N_{i-1}; \mathbb{R}),$$

with the convention that $N_0 = \emptyset$.

We define the space of *homologically weighted flags of type S in M* by

$$\text{Flag}_S^{\text{hwt}}(M) := \{(\mathcal{N}, [\nu]) : \mathcal{N} \in \text{Flag}_S(M), [\nu] \in H(\mathcal{N})\}.$$

Note that we have a $\text{Diff}(M)$ equivariant forgetful map

$$(2.38) \quad \text{Flag}_S^{\text{hwt}}(M) \rightarrow \text{Flag}_S(M),$$

as well as a $\text{Diff}(M)$ equivariant map

$$(2.39) \quad h_{\text{Flag}_S(M)} : \text{Flag}_S^{\text{wt}}(M) \rightarrow \text{Flag}_S^{\text{hwt}}(M), \quad (\mathcal{N}, \nu) \mapsto (\mathcal{N}, [\nu]).$$

Let $\text{Flag}_{S, \iota}^{\text{hwt}}(M)$ denote the preimage of the open subset $\text{Flag}_{S, \iota}(M)$ under the projection in (2.38). Using the canonical $\text{Diff}(M)$ equivariant identifications

$$(2.40) \quad \text{Flag}_{S, \iota}^{\text{hwt}}(M) = \text{Emb}_{S, r}(M) \times_{\text{Diff}(S, \iota)} H(S, \iota),$$

we equip $\text{Flag}_S^{\text{hwt}}(M)$ with the structure of a smooth vector bundle of finite (possibly nonconstant) rank over $\text{Flag}_S(M)$ and with projection (2.38). The map in (2.39) is a smooth bundle map over $\text{Flag}_S(M)$. Indeed, via the identifications in (2.25) and (2.40), the map $h_{S, \iota}$ in (2.28) induces a bundle map $\text{Flag}_{S, \iota}^{\text{wt}}(M) \rightarrow \text{Flag}_{S, \iota}^{\text{hwt}}(M)$ which coincides with the restriction of (2.39).

We define the space of *homologically weighted flags of type $(S, \iota, [\mu])$ in M* by

$$\text{Flag}_{S, \iota, [\mu]}^{\text{hwt}}(M) := \left\{ (\mathcal{N}, [\nu]) \in \text{Flag}_S^{\text{hwt}}(M) \mid \left(S_1 \xrightarrow{\iota_1} S_2 \xrightarrow{\iota_2} \dots \rightarrow S_r, [\mu] \right) \right. \\ \left. \cong (N_1 \subseteq N_2 \subseteq \dots \subseteq N_r, [\nu]) \right\},$$

that is, the space of all homologically weighted flags $(\mathcal{N}, [\nu])$ in M such that there exist diffeomorphisms $S_i \rightarrow N_i, 1 \leq i \leq r$, intertwining ι_i with the canonical inclusion $N_i \subseteq N_{i+1}$ and taking $[\mu] \in H(S, \iota)$ to $[\nu] \in H(\mathcal{N})$. The diffeomorphism in (2.40) restricts to a bijection

$$(2.41) \quad \text{Flag}_{S, \iota, [\mu]}^{\text{hwt}}(M) = \text{Emb}_{S, r}(M) \times_{\text{Diff}(S, \iota)} H(S, \iota)_{[\mu]}.$$

Hence, since the orbit $H(\mathcal{S}, \iota)_{[\mu]}$ is finite, $\text{Flag}_{\mathcal{S}, \iota, [\mu]}^{\text{hwt}}(M)$ is a $\text{Diff}(M)$ invariant closed submanifold in $\text{Flag}_{\mathcal{S}, \iota}^{\text{hwt}}(M)$ of codimension $\dim H(\mathcal{S}, \iota)$ by (2.40). Moreover, the forgetful map

$$(2.42) \quad \text{Flag}_{\mathcal{S}, \iota, [\mu]}^{\text{hwt}}(M) \rightarrow \text{Flag}_{\mathcal{S}, \iota}(M)$$

is a $\text{Diff}(M)$ equivariant covering map with finite fibers.

We complement Theorem 2.11 with the following.

Theorem 2.15 *In this situation, we have*

$$\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt}}(M) = h_{\text{Flag}_{\mathcal{S}}(M)}^{-1} \left(\text{Flag}_{\mathcal{S}, \iota, [\mu]}^{\text{hwt}}(M) \right).$$

Proof This follows from the identity (2.29) in Proposition 2.10(c), using (2.26) and (2.41). ■

Remark 2.16 If the action of $\text{Diff}(\mathcal{S}, \iota)$ on $H(\mathcal{S}, \iota)$ is trivial, then the $\text{Diff}(\mathcal{S}, \iota)_0$ and $\text{Diff}(\mathcal{S}, \iota)$ orbits in $\text{Den}_\times(\mathcal{S})$ coincide in view of Proposition 2.10, and the forgetful (covering) map in (2.42) is a diffeomorphism, $\text{Flag}_{\mathcal{S}, \iota, [\mu]}^{\text{hwt}}(M) = \text{Flag}_{\mathcal{S}, \iota}(M)$. This happens in particular when all $S_i \setminus \iota(S_{i-1})$ are connected, as in Examples 2.17 and 2.18. Under the latter connectedness assumption, $H(\mathcal{S}, \iota) = \mathbb{R}^r$ via the isomorphism $[\mu] \mapsto (\int_{S_1} \mu_1, \dots, \int_{S_r} \mu_r)$ and

$$\text{Den}(\mathcal{S})_{\iota, \mu} = \left\{ \alpha \in \text{Den}_\times(\mathcal{S}) : \int_{S_i} \alpha_i = \int_{S_i} \mu_i \right\}.$$

Moreover, Theorem 2.15 ensures that

$$(2.43) \quad \text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt}}(M) = \left\{ (\mathcal{N}, \nu) \in \text{Flag}_{\mathcal{S}, \iota}^{\text{wt}}(M) : \int_{N_i} \nu_i = \int_{S_i} \mu_i \right\}.$$

This also applies in the situation of Remark 2.9, provided each model manifold S_i is connected. A description for nested spheres in the same vein can be found in Example 2.19.

Example 2.17 (Nested tori) If (\mathcal{S}, ι) denotes the standard (meridional) embeddings between tori, $\mathbb{T}^0 \subseteq \mathbb{T}^1 \subseteq \dots \subseteq \mathbb{T}^r$, then $H(\mathcal{S}, \iota) = \mathbb{R}^{r+1}$ via the isomorphism $[\mu] \mapsto (\int_{\mathbb{T}^i} \mu_i)$.

Example 2.18 (Nested projective spaces) If (\mathcal{S}, ι) denotes the standard embeddings between projective spaces, $\mathbb{P}^0 \subseteq \mathbb{P}^1 \subseteq \dots \subseteq \mathbb{P}^r$, then $H(\mathcal{S}, \iota) = \mathbb{R}^{r+1}$ via the isomorphism $[\mu] \mapsto (\int_{\mathbb{P}^i} \mu_i)$.

Example 2.19 (Nested spheres [13]) If (\mathcal{S}, ι) denotes the standard equatorial embeddings between spheres, $S^0 \subseteq S^1 \subseteq \dots \subseteq S^r$, then $H(\mathcal{S}, \iota) = \mathbb{R}^{2(r+1)}$. The $2(r+1)$ numbers assigned to $[\mu] \in H(\mathcal{S}, \iota)$ by this isomorphism are

$$(a_0^+, a_0^-, a_1^+, a_1^-, \dots, a_r^+, a_r^-) = \left(\int_{S_+^0} \mu_0, \int_{S_-^0} \mu_0, \int_{S_+^1} \mu_1, \int_{S_-^1} \mu_1, \dots, \int_{S_+^r} \mu_r, \int_{S_-^r} \mu_r \right),$$

where S_+^i and S_-^i denote the northern and southern hemispheres of S^i , respectively. Considering reflections on hyperplanes, we see that for each $0 \leq i \leq r$ there exists a diffeomorphism in $\text{Diff}(\mathcal{S}, \iota)$ swapping S_+^i with S_-^i , but leaving all other hemispheres S_\pm^k invariant. Such a diffeomorphism interchanges a_i^+ with a_i^- , but leaves all other numbers a_k^\pm unchanged. Hence, the $\text{Diff}(\mathcal{S}, \iota)$ orbit $H(\mathcal{S}, \iota)_{[\mu]}$ has 2^s elements,

where s is the number of $0 \leq i \leq r$ with $a_i^+ \neq a_i^-$. Actually, the $\text{Diff}(S, \iota)$ action on $H(S, \iota) \cong \mathbb{R}^{2(r+1)}$ factorizes through a $(\mathbb{Z}_2)^{2(r+1)}$ action by switching or not the numbers a_i^+ and a_i^- . We obtain a description similar to the one in (2.43):

$$(2.44) \quad \text{Flag}_{S, \iota, \mu}^{\text{wt}}(M) = \left\{ (\mathcal{N}, \nu) \in \text{Flag}_{S, \iota}^{\text{wt}}(M) \left| \begin{array}{l} \{\int_{N_i^+} \nu_i, \int_{N_i^-} \nu_i\} = \{a_i^+, a_i^-\}, \\ \text{where } N_i^\pm \text{ denote the connected} \\ \text{components of } N_i \setminus N_{i-1} \end{array} \right. \right\}.$$

3 Coadjoint orbits of the Hamiltonian group

Throughout this section, (M, ω) denotes a symplectic manifold. The nonlinear Grassmannian of all isotropic submanifolds of type S_r in M , denoted by $\text{Gr}_{S_r}^{\text{iso}}(M)$, is a splitting smooth submanifold in $\text{Gr}_{S_r}(M)$ which is invariant under the Hamiltonian group. In fact, the $\text{Ham}(M)$ orbits provide a smooth foliation of finite codimension in $\text{Gr}_{S_r}^{\text{iso}}(M)$ which is called the isodrastic foliation [15, 26].

Suppose \mathcal{L} is an isodrastic leaf in $\text{Gr}_{S_r}^{\text{iso}}(M)$, and let $\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ denote the preimage of \mathcal{L} under the canonical bundle projection $\text{Flag}_{S, \iota, \mu}^{\text{wt}}(M) \rightarrow \text{Gr}_{S_r}(M)$. We will show that the natural $\text{Ham}_c(M)$ action on $\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ admits local smooth sections. In particular, each connected component of the latter space is an orbit of $\text{Ham}_c(M)$.

The space $\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ comes equipped with a canonical weakly non-degenerate symplectic form, and the restriction of (2.18) provides a $\text{Ham}(M)$ equivariant injective moment map for the $\text{Ham}_c(M)$ action,

$$J: \text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}} \hookrightarrow \mathfrak{ham}_c(M)^*, \quad \langle J(\mathcal{N}, \nu), X_f \rangle = \sum_{i=1}^r \int_{N_i} f \nu_i.$$

This moment map J maps each connected component of $\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ one-to-one onto the corresponding coadjoint orbit (see Theorem 3.15). Thereby, we identify coadjoint orbits of the Hamiltonian group $\text{Ham}_c(M)$ that can be modeled on weighted nonlinear flags.

The material in this section is inspired by the results in [7, 15, 26] on weighted isotropic nonlinear Grassmannians.

3.1 Isodrasts as $\text{Ham}(M)$ orbits

In view of the tubular neighborhood theorem for isotropic embeddings [24, 25], the space $\text{Gr}_S^{\text{iso}}(M)$ of all isotropic submanifolds of type S in M is a splitting smooth submanifold of $\text{Gr}_S(M)$ (see, for instance, [15, Section 8]). The tangent space at an isotropic submanifold N is

$$T_N \text{Gr}_S^{\text{iso}}(M) = \{u_N \in \Gamma(TN^\perp) | \iota_N^* i_{u_N} \omega \in \Omega^1(N) \text{ closed}\},$$

where $TN^\perp := TM|_N/TN$ denotes the normal bundle to N .

Weinstein's [26] isodrastic distribution $\underline{\mathcal{D}}$ on $\text{Gr}_S^{\text{iso}}(M)$ is given by

$$(3.1) \quad \underline{\mathcal{D}}_N := \{u_N \in \Gamma(TN^\perp) | \iota_N^* i_{u_N} \omega \in dC^\infty(N)\}$$

and has finite codimension $\dim H^1(S; \mathbb{R})$. This is an integrable distribution [15, 26] whose leaves are orbits of $\text{Ham}(M)$. It gives rise to a smooth foliation of $\text{Gr}_S^{\text{iso}}(M)$ called the *isodrastic foliation*. In particular, the $\text{Ham}(M)$ orbits in $\text{Gr}_S^{\text{iso}}(M)$ are splitting smooth submanifolds.

Restricting the fundamental frame bundle in (2.1), we obtain a principal $\text{Diff}(S)$ bundle $\text{Emb}_S^{\text{iso}}(M) \rightarrow \text{Gr}_S^{\text{iso}}(M)$ with total space $\text{Emb}_S^{\text{iso}}(M)$, the splitting smooth submanifold of $\text{Emb}_S(M)$ consisting of all isotropic embeddings. On $\text{Emb}_S^{\text{iso}}(M)$, we consider the pullback of the isodrastic distribution $\underline{\mathcal{D}}$:

$$(3.2) \quad \mathcal{D}_\varphi := \{u_\varphi \in \Gamma(\varphi^* TM) : \varphi^* i_{u_\varphi} \omega \in dC^\infty(S)\}.$$

This is an integrable distribution with the same codimension, $\dim H^1(S; \mathbb{R})$, and the leaves of \mathcal{D} are connected components in the preimage of a leaf of $\underline{\mathcal{D}}$. According to [15, 26], the group $\text{Ham}(M)$ acts transitively on the leaves of \mathcal{D} . We need the subsequent slightly stronger statement in Proposition 3.1.

The Lie algebra of compactly supported Hamiltonian vector fields will be denoted by $\mathfrak{ham}_c(M) = \{X_f : f \in C_c^\infty(M)\}$. We let $\text{Ham}_c(M)$ denote the group of diffeomorphisms obtained by integrating time dependent vector fields in $\mathfrak{ham}_c(M)$. For our purpose, it will not be necessary to consider $\text{Ham}_c(M)$ as an infinite-dimensional Lie group (cf. [14, Section 43.13]). By a smooth map (section) into $\text{Ham}_c(M)$, we will simply mean a smooth map into $\text{Diff}_c(M)$ that takes values in $\text{Ham}_c(M)$.

Proposition 3.1 *The $\text{Ham}_c(M)$ action on each leaf $\mathcal{E} \subseteq \text{Emb}_S^{\text{iso}}(M)$ of the isodrastic foliation \mathcal{D} admits local smooth sections.*

Proof To show infinitesimal transitivity, suppose $\varphi \in \text{Emb}_S^{\text{iso}}(M)$ and $u_\varphi \in \mathcal{D}_\varphi$ (cf. (3.2)). Hence, there exists $\tilde{f} \in C^\infty(S)$ such that $d\tilde{f} = \varphi^* i_{u_\varphi} \omega$. We extend \tilde{f} to a function $f_1 \in C_c^\infty(M)$ such that $\tilde{f} = f_1 \circ \varphi$. The 1-form β on M along S defined by

$$(3.3) \quad \beta = df_1 \circ \varphi - i_{u_\varphi} \omega \in \Gamma(\varphi^* T^* M)$$

vanishes on vectors tangent to $\varphi(S) \subseteq M$. Hence, β can be seen as a fiberwise linear function on the normal bundle $T\varphi(S)^\perp$ whose differential along the zero section $\varphi(S)$ coincides with β itself. Thus, with the help of a tubular neighborhood of $\varphi(S)$ in M and a suitable bump function, we get $f_2 \in C_c^\infty(M)$ such that $\beta = df_2 \circ \varphi$. It follows from (3.3) that $df \circ \varphi = i_{u_\varphi} \omega$ for $f = f_1 - f_2$. We conclude that $u_\varphi = X_f \circ \varphi$ is the infinitesimal generator at φ for the Hamiltonian vector field $X_f \in \mathfrak{ham}_c(M)$.

Using tubular neighborhoods constructed with the help of a Riemannian metric, say, we see that the function f may be chosen to depend smoothly on φ and u_φ , for φ in a sufficiently small open neighborhood of a fixed isotropic embedding $\varphi_0 \in \mathcal{E}$. Hence, we may apply Lemma A.2 and conclude that the $\text{Ham}_c(M)$ action admits local smooth sections. ■

Corollary 3.2 *The $\text{Ham}_c(M)$ action on each leaf $\mathcal{L} \subseteq \text{Gr}_S^{\text{iso}}(M)$ of the isodrastic foliation $\underline{\mathcal{D}}$ admits local smooth sections.*

Example 3.3 Every embedded closed curve in the plane is a Lagrangian submanifold of (\mathbb{R}^2, ω) , where ω is the canonical area form, thus an element of $\text{Gr}_{S^1}^{\text{iso}}(\mathbb{R}^2)$. The isodrastic distribution $\underline{\mathcal{D}}$ has codimension one. The enclosed area a singles out one isodrast $\mathcal{L}_a \subseteq \text{Gr}_{S^1}^{\text{iso}}(\mathbb{R}^2)$, i.e., one orbit of $\text{Ham}_c(\mathbb{R}^2)$.

A similar phenomena happens for Lagrangian k -tori in \mathbb{R}^{2k} , i.e., elements of $\text{Gr}_{\mathbb{T}^k}^{\text{iso}}(\mathbb{R}^{2k})$, where $\mathbb{T}^k := (S^1)^k$. To any $\varphi \in \text{Emb}_{\mathbb{T}^k}^{\text{iso}}(\mathbb{R}^{2k})$, we assign the symplectic area a_i of the surface in \mathbb{R}^{2k} enclosed by the i th meridian $\varphi_i(\theta) = \varphi(1, \dots, \theta, \dots, 1)$ of the embedded k -torus. These numbers are independent of the choice of the meridian in its homotopy class and of the surface having the meridian as boundary. The k -tuple (a_1, \dots, a_k) is an invariant under isodrastic deformations. Actually, a_i is the action integral of the i th meridian, as defined in [26].

Let $\mathcal{E}_{a_1, \dots, a_k} \subseteq \text{Emb}_{\mathbb{T}^k}^{\text{iso}}(\mathbb{R}^{2k})$ be the space of all isotropic embeddings having symplectic areas a_1, \dots, a_k . It is a union of isodrastic leaves of Lagrangian embeddings, but it is not necessarily $\text{Diff}(\mathbb{T}^k)$ saturated. The $\text{Diff}(\mathbb{T}^k)$ action on the k -tuples (a_1, \dots, a_k) factorizes through a $\text{GL}(k, \mathbb{Z})$ action. Let $[a_1, \dots, a_k]$ denote the orbit of (a_1, \dots, a_k) . We define $\mathcal{L}_{[a_1, \dots, a_k]} \subseteq \text{Gr}_{\mathbb{T}^k}^{\text{iso}}(\mathbb{R}^{2k})$ to be the image of $\mathcal{E}_{a_1, \dots, a_k}$ under the principal $\text{Diff}(\mathbb{T}^k)$ bundle projection (2.1). Thus, $\mathcal{L}_{[a_1, \dots, a_k]}$ is a union of isodrastic leaves of Lagrangian k -tori in \mathbb{R}^{2k} .

There is also a direct description of $\mathcal{L}_{[a_1, \dots, a_k]}$. Given a Lagrangian torus $N \in \text{Gr}_{\mathbb{T}^k}^S(\mathbb{R}^{2k})$, we choose a base $\{[\gamma_1], \dots, [\gamma_k]\}$ of $H_1(N, \mathbb{Z})$, where γ_i are loops in N with action integrals a_i . We observe that the $\text{GL}(k, \mathbb{Z})$ orbit $[a_1, \dots, a_k]$ is independent of the choices and $\mathcal{L}_{[a_1, \dots, a_k]}$ is the space of all Lagrangian tori in $\text{Gr}_{\mathbb{T}^k}^S(\mathbb{R}^{2k})$ such that the orbit of these action integrals is $[a_1, \dots, a_k]$.

We will also need the following observation.

Lemma 3.4 *If L is an isotropic submanifold in M , then the canonical inclusion $\text{Gr}_S(L) \subseteq \text{Gr}_S^{\text{iso}}(M)$ is a splitting smooth submanifold. Moreover, each connected component of $\text{Gr}_S(L)$ is a splitting smooth submanifold in an isodrastic leaf in $\text{Gr}_S^{\text{iso}}(M)$.*

Proof Suppose $N \in \text{Gr}_S(L)$, i.e., $N \cong S$ is a closed submanifold in L . By the tubular neighborhood theorem, we may w.l.o.g. assume that L is the total space of a vector bundle $p: L \rightarrow N$, the normal bundle of N in L , and identify N with the zero section in L . We have a canonical short exact sequence $0 \rightarrow p^*L \rightarrow TL \rightarrow p^*TN \rightarrow 0$ of vector bundles over L . Choosing a linear connection on L , we obtain a splitting of this sequence and thus an isomorphism $TL \cong p^*TN \oplus p^*L$ of vector bundles over L . Dualizing, we obtain an isomorphism $T^*L \cong p^*T^*N \oplus p^*L^*$ of vector bundles over L . We regard this as a diffeomorphism

$$T^*L \cong T^*N \oplus L \oplus L^*$$

that maps the zero section $L \subseteq T^*L$ identically onto the summand L on the right-hand side. Via this isomorphism, we have

$$\theta_L = \pi_1^* \theta_N + \pi_2^* \kappa,$$

where π_1 and π_2 denote the projections from $T^*N \oplus L \oplus L^*$ onto T^*N and $L \oplus L^*$, respectively, $\theta_L \in \Omega^1(T^*L)$ and $\theta_N \in \Omega^1(T^*N)$ denote the tautological 1-forms, and $\kappa \in \Omega^1(L \oplus L^*)$. Indeed, a straightforward computation yields $\kappa(\xi) = \ell'(C(Tq \cdot \xi))$ where $x \in N, \ell \in L_x, \ell' \in L_x^*, \xi \in T_{(\ell, \ell')}(L \oplus L^*), q: L \oplus L^* \rightarrow L$ denotes the projection, and C denotes the linear connection on L , viewed as a fiberwise linear map $C: TL \rightarrow p^*L$ over L .

By the tubular neighborhood theorem for isotropic embeddings [24, 25], we may assume $M = T^*L \oplus p^*E$ and $\omega = \tilde{\pi}_1^*d\theta_L + \tilde{\pi}_2^*\rho$, where E is a vector bundle over N , $\tilde{\pi}_1$ and $\tilde{\pi}_2$ denote the projections from $T^*L \oplus p^*E$ onto T^*L and p^*E , respectively, and ρ is a closed 2-form on the total space of p^*E . Combining this with the diffeomorphism in the previous paragraph, we obtain a diffeomorphism

$$(3.4) \quad M = T^*N \oplus L \oplus L^* \oplus E,$$

mapping $L \subseteq M$ identically onto the summand L on the right-hand side, and such that

$$\omega = \pi_1^*d\theta_N + \pi_2^*\sigma,$$

where π_1 and π_2 denote the projections from $T^*N \oplus L \oplus L^* \oplus E$ onto T^*N and $L \oplus L^* \oplus E$, respectively, and σ is a closed 2-form on the total space of $L \oplus L^* \oplus E$.

The diffeomorphism in (3.4) provides a (standard) chart for the smooth structure on $\text{Gr}_S(M)$ centered at N ,

$$\Gamma(T^*N \oplus L \oplus L^* \oplus E) \rightarrow \text{Gr}_S(M), \quad \phi \mapsto \phi(N).$$

In this chart, the inclusions $\text{Gr}_S(L) \subseteq \text{Gr}_S^{\text{iso}}(M) \subseteq \text{Gr}_S(M)$ become

$$(3.5) \quad \Gamma(L) \subseteq \{(\alpha, \xi) \in \Omega^1(N) \times \Gamma(L \oplus L^* \oplus E) : d\alpha + \xi^*\sigma = 0\} \subseteq \Omega^1(N) \times \Gamma(L \oplus L^* \oplus E).$$

As L is isotropic, σ vanishes when pulled back to L . Hence, by the Poincaré lemma, $\sigma = d\beta$ for a 1-form β on the total space of $L \oplus L^* \oplus E$ which vanishes along L . Via the diffeomorphism of $\Omega^1(N) \times \Gamma(L \oplus L^* \oplus E)$ given by $(\alpha, \xi) \mapsto (\alpha - \xi^*\beta, \xi)$, the inclusions in (3.5) become linear inclusions,

$$\Gamma(L) \subseteq Z^1(N) \times \Gamma(L \oplus L^* \oplus E) \subseteq \Omega^1(N) \times \Gamma(L \oplus L^* \oplus E),$$

where $Z^1(N)$ denotes the space of closed 1-forms on N . Clearly, both inclusions admit complementary subspaces. In particular, $\text{Gr}_S(L)$ is a splitting smooth submanifold in $\text{Gr}_S^{\text{iso}}(M)$. The second assertion follows from the fact that the isodrastic leaf through N corresponds to the subspace $B^1(N) \times \Gamma(L \oplus L^* \oplus E)$ (see [15, Section 8]). ■

3.2 Weighted isotropic nonlinear Grassmannians as coadjoint orbits

In this section, we recall the results in [7, 15, 26] about coadjoint orbits of the Hamiltonian group $\text{Ham}_c(M)$ modeled on weighted isotropic nonlinear Grassmannians of type S in M , and extend them to a possibly nonconnected model manifold S . Here, we present them in a manner that readily generalizes to manifolds of weighted nonlinear flags.

Let S be a closed k -dimensional manifold. The preimage of $\text{Gr}_S^{\text{iso}}(M)$ under the canonical bundle projection $\text{Gr}_S^{\text{wt}}(M) \rightarrow \text{Gr}_S(M)$ is a splitting smooth submanifold in $\text{Gr}_S^{\text{wt}}(M)$ which will be denoted by $\text{Gr}_S^{\text{wt iso}}(M)$. The diffeomorphism in (2.3) restricts to a diffeomorphism of bundles over $\text{Gr}_S^{\text{iso}}(M)$,

$$\text{Gr}_S^{\text{wt iso}}(M) = \text{Emb}_S^{\text{iso}}(M) \times_{\text{Diff}(S)} \text{Den}_\times(S).$$

The preimage of $\text{Gr}_S^{\text{iso}}(M)$ under the canonical bundle projection $\text{Gr}_{S,\mu}^{\text{wt}}(M) \rightarrow \text{Gr}_S(M)$ is a splitting smooth submanifold in $\text{Gr}_{S,\mu}^{\text{wt}}(M)$ which will be denoted by $\text{Gr}_{S,\mu}^{\text{wt iso}}(M)$ and will be referred to as the *nonlinear Grassmannian of weighted isotropic submanifolds of type (S, μ) in M* . The diffeomorphism in (2.5) restricts to a diffeomorphism of bundles over $\text{Gr}_S^{\text{iso}}(M)$,

$$\text{Gr}_{S,\mu}^{\text{wt iso}}(M) = \text{Emb}_S^{\text{iso}}(M) \times_{\text{Diff}(S)} \text{Den}(S)_\mu.$$

We recall the $\text{Diff}(S)$ equivariant linear map $h_S : \text{Den}(S) \rightarrow H^k(S, \mathcal{O}_S)$, $h_S(\alpha) = [\alpha]$ in (2.6), with kernel $d\Omega^{k-1}(S, \mathcal{O}_S)$, which we restrict to $\text{Den}_\times(S)$. The product of integrable distributions $\mathcal{D} \times \ker Th_S$ on $\text{Emb}_S^{\text{iso}}(M) \times \text{Den}_\times(S)$ descends to an integrable distribution

$$\tilde{\mathcal{D}} := \mathcal{D} \times_{\text{Diff}(S)} \ker Th_S$$

on $\text{Gr}_S^{\text{wt iso}}(M)$, of codimension $\dim H^1(S; \mathbb{R}) + \dim H^k(S; \mathcal{O}_S)$. The image of $\tilde{\mathcal{D}}$ under the forgetful map $\text{Gr}_S^{\text{wt iso}}(M) \rightarrow \text{Gr}_S^{\text{iso}}(M)$ is the isodrastic distribution $\underline{\mathcal{D}}$ in (3.1). In view of (2.7), each leaf \mathcal{G} of $\tilde{\mathcal{D}}$ is a connected component in the preimage of an isodrast \mathcal{L} in $\text{Gr}_S^{\text{iso}}(M)$ under the bundle projection $\text{Gr}_{S,\mu}^{\text{wt iso}}(M) \rightarrow \text{Gr}_S^{\text{iso}}(M)$, for some volume density μ on S . Hence, each leaf \mathcal{G} of $\tilde{\mathcal{D}}$ is a splitting smooth submanifold of codimension $\dim H^1(S; \mathbb{R})$ in $\text{Gr}_{S,\mu}^{\text{wt iso}}(M)$.

According to [15, 26], the group $\text{Ham}(M)$ acts transitively on the leaves of $\tilde{\mathcal{D}}$. We need the following slightly stronger statement.

Proposition 3.5 *The $\text{Ham}_c(M)$ action on each leaf $\mathcal{G} \subseteq \text{Gr}_S^{\text{wt iso}}(M)$ of the isodrastic foliation $\tilde{\mathcal{D}}$ admits local smooth sections.*

Proof Suppose \mathcal{L} is an isodrastic leaf in $\text{Gr}_S^{\text{iso}}(M)$, and let $\text{Gr}_{S,\mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ denote its preimage under the bundle projection $\text{Gr}_{S,\mu}^{\text{wt iso}}(M) \rightarrow \text{Gr}_S^{\text{iso}}(M)$. It suffices to show that the $\text{Ham}_c(M)$ action on $\text{Gr}_{S,\mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ admits local smooth sections. The diffeomorphism in (2.5) restricts to a diffeomorphism

$$(3.6) \quad \text{Gr}_{S,\mu}^{\text{wt iso}}(M)|_{\mathcal{L}} = \text{Emb}_S^{\text{iso}}(M)|_{\mathcal{L}} \times_{\text{Diff}(S)} \text{Den}(S)_\mu.$$

By Proposition 3.1, the $\text{Ham}_c(M)$ action on $\text{Emb}_S^{\text{iso}}(M)|_{\mathcal{L}}$ admits local smooth sections. According to Proposition 2.10(b), the $\text{Diff}(S)$ action on $\text{Den}(S)_\mu$ admits local smooth sections too. Using Lemma A.1 in the Appendix, we conclude that the $\text{Ham}_c(M)$ action on the associated bundle in (3.6) admits local smooth sections. ■

Lemma 3.6 [15] *The leafwise differential 2-form on $(\text{Emb}_S^{\text{iso}}(M) \times \text{Den}_\times(S), \mathcal{D} \times \ker Th_S)$, given by*

$$(3.7) \quad \Omega_{(\varphi, \alpha)}((u_\varphi, d\lambda), (v_\varphi, d\gamma)) := \int_S (\omega(u_\varphi, v_\varphi)\alpha - \varphi^* i_{u_\varphi} \omega \wedge \gamma + \varphi^* i_{v_\varphi} \omega \wedge \lambda),$$

is closed and $\text{Diff}(S)$ invariant. Moreover, its kernel is spanned by the infinitesimal generators of the $\text{Diff}(S)$ action.

In Section 3.4, we will need the following relative version of part of Lemma 3.6.

Lemma 3.7 *Let Σ be a closed submanifold in S of codimension at least one. Suppose $(\varphi, \alpha) \in \text{Emb}_S^{\text{iso}}(M) \times \text{Den}_\times(S)$ and $(u_\varphi, d\lambda) \in \mathcal{D}_\varphi \times \ker T_\alpha h_S$. If the leafwise differential 2-form (3.7) vanishes for all $(v_\varphi, d\gamma) \in \mathcal{D}_\varphi \times \ker T_\alpha h_S$ with $i_\Sigma^* \gamma = 0$ and $v_\varphi|_\Sigma = 0$, then $(u_\varphi, d\lambda)$ is the infinitesimal generator of some $Z \in \mathfrak{X}(S)$ at (φ, α) , i.e., $(u_\varphi, d\lambda) = (T\varphi \circ Z, L_Z \alpha)$.*

Proof Choosing $v_\varphi = 0$ in (3.7), we obtain $\int_S (\varphi^* i_{u_\varphi} \omega) \wedge \gamma = 0$ for all $\gamma \in \Omega^{k-1}(S, \mathcal{O}_S)$ with $i_\Sigma^* \gamma = 0$. Since the codimension of Σ in S is at least one, this yields $\varphi^* i_{u_\varphi} \omega = 0$; hence, u_φ takes values in the symplectic orthogonal to $\varphi(S)$ in M . In particular, (3.7) yields

$$(3.8) \quad \int_S \omega(u_\varphi, v_\varphi) \alpha + \int_S \varphi^* i_{v_\varphi} \omega \wedge \lambda = 0$$

for all v_φ with $v_\varphi|_\Sigma = 0$.

To show that u_φ is tangential to $\varphi(S)$, we assume by contradiction that there is a point $x \in S$ such that $u_\varphi(x)$ is not tangent to $\varphi(S)$. Since the codimension of Σ is at least one, we may assume $x \notin \Sigma$. We now consider tangent vectors $v_\varphi \in \mathcal{D}_\varphi$ taking values in the symplectic orthogonal to $\varphi(S)$, i.e., $\varphi^* i_{v_\varphi} \omega = 0$, supported in $S \setminus \Sigma$. The isotropic embedding theorem of Weinstein [25] allows to choose v_φ such that in addition $\omega(u_\varphi, v_\varphi) \in C^\infty(S)$ is a nonzero nowhere negative function. Plugging v_φ in (3.8), we obtain $\int_S \omega(u_\varphi, v_\varphi) \alpha > 0$, leading to a contradiction. Thus, there exists $Z \in \mathfrak{X}(S)$ with $u_\varphi = T\varphi \circ Z$.

For each $h \in C^\infty(S)$ with $dh|_\Sigma = 0$, there exists $v_\varphi \in \mathcal{D}_\varphi$ with $v_\varphi|_\Sigma = 0$ such that $\varphi^* i_{v_\varphi} \omega = dh$. Plugging this into the identity in (3.8) and using Stokes' theorem yields $\int_S h d(\lambda - i_Z \alpha) = 0$. In particular, this holds for all h with $\text{supp}(h) \cap \Sigma = \emptyset$. As the codimension of Σ is at least one, this implies $d(\lambda - i_Z \alpha) = 0$, and hence $d\lambda = L_Z \alpha$. ■

By Lemma 3.6, the leafwise differential 2-form Ω in (3.7) descends to a leafwise symplectic form $\tilde{\Omega}$ on $(\text{Gr}_S^{\text{wt iso}}(M), \tilde{\mathcal{D}})$. Thus, every leaf \mathcal{G} of the isodrastic distribution $\tilde{\mathcal{D}}$ in $\text{Gr}_S^{\text{wt iso}}(M)$ is endowed with a symplectic form, the restriction of $\tilde{\Omega}$, which we denote by the same letter. By Proposition 3.5, \mathcal{G} is an orbit for the $\text{Ham}_c(M)$ action on the weighted isotropic nonlinear Grassmannian $\text{Gr}_S^{\text{wt iso}}(M)$. This action is Hamiltonian with injective and $\text{Symp}(M)$ equivariant moment map [15]

$$(3.9) \quad J : \mathcal{G} \subseteq \text{Gr}_S^{\text{wt iso}}(M) \rightarrow \mathfrak{ham}_c(M)^*, \quad \langle J(N, v), X_f \rangle = \int_N f v.$$

Here, we use the $\text{Symp}(M)$ equivariant isomorphism $\mathfrak{ham}_c(M) = C_0^\infty(M)$ where the latter denotes the Lie algebra of all compactly supported functions on M for which the integral with respect to the Liouville form vanishes on all closed connected components of M .

For connected S , the subsequent theorem is due to Weinstein [26] in the Lagrangian case and due to Lee [15] for isotropic submanifolds.

Theorem 3.8 [15, 26] *The moment map $J : (\mathcal{G}, \tilde{\Omega}) \rightarrow \mathfrak{ham}_c(M)^*$ in (3.9) is one-to-one onto a coadjoint orbit of the Hamiltonian group $\text{Ham}_c(M)$. The Kostant–Kirillov–Souriau symplectic form ω_{KKS} on the coadjoint orbit satisfies $J^* \omega_{\text{KKS}} = \tilde{\Omega}$.*

In this generality, the theorem follows from the discussion above and the following folklore result (see, for instance, the Appendix in [9]):

Proposition 3.9 *Suppose the action of G on (\mathcal{M}, Ω) is transitive with injective equivariant moment map $J: \mathcal{M} \rightarrow \mathfrak{g}^*$. Then J is one-to-one onto a coadjoint orbit of G . Moreover, it pulls back the Kostant–Kirillov–Souriau symplectic form ω_{KKS} on the coadjoint orbit to the symplectic form Ω .*

The same coadjoint orbits of the Hamiltonian group, under the additional restriction $H^1(S; \mathbb{R}) = 0$ on the closed connected manifold S , can be obtained via symplectic reduction in the Marsden–Weinstein ideal fluid dual pair [5, 17], as shown in [7]. With a choice of an ambient Riemannian metric, the reduced symplectic form can be expressed as a sum of three terms (see [7, Theorem 4.1]). These are analogous to the three summands appearing in our natural approach via the associated bundle construction (see (3.7)), which avoids the auxiliary Riemannian metric.

3.3 Manifolds of isotropic nonlinear flags

In this section, we extend the constructions of the previous section to nonlinear flags.

As in Section 2.3, we let $\iota = (\iota_1, \dots, \iota_{r-1})$ denote a collection of fixed embeddings $\iota_i: S_i \rightarrow S_{i+1}$. We let $\text{Flag}_{S,\iota}^{\text{iso}}(M)$ denote the preimage of $\text{Gr}_{S_r}^{\text{iso}}(M)$ under the canonical bundle projection $\text{Flag}_{S,\iota}(M) \rightarrow \text{Gr}_{S_r}(M)$ (cf. [9, Remark 2.11]). This is a splitting smooth submanifold in $\text{Flag}_{S,\iota}(M)$ called the manifold of *isotropic nonlinear flags of type (S, ι) in M* . Using Lemma 3.4, and proceeding by induction on the depth of the flag, one readily shows that the canonical inclusion

$$(3.10) \quad \text{Flag}_{S,\iota}^{\text{iso}}(M) \subseteq \prod_{i=1}^r \text{Gr}_{S_i}^{\text{iso}}(M)$$

is a splitting smooth submanifold.

Let $\text{Flag}_{S,\iota}^{\text{wt iso}}(M)$ denote the preimage of $\text{Flag}_{S,\iota}^{\text{iso}}(M)$ under the canonical bundle projection $\text{Flag}_{S,\iota}^{\text{wt}}(M) \rightarrow \text{Flag}_{S,\iota}(M)$. This is a splitting smooth submanifold in $\text{Flag}_{S,\iota}^{\text{wt}}(M)$ called the manifold of *weighted isotropic nonlinear flags of type (S, ι) in M* . The diffeomorphism in (2.25) restricts to a diffeomorphism of bundles over $\text{Flag}_{S,\iota}^{\text{iso}}(M)$,

$$\text{Flag}_{S,\iota}^{\text{wt iso}}(M) = \text{Emb}_{S_r}^{\text{iso}}(M) \times_{\text{Diff}(S;\iota)} \text{Den}_{\times}(S).$$

The canonical inclusion $\text{Flag}_{S,\iota}^{\text{wt iso}}(M) \subseteq \prod_{i=1}^r \text{Gr}_{S_i}^{\text{wt iso}}(M)$ is a splitting smooth submanifold in view of (3.10).

As in Section 2.4, we fix $\mu = (\mu_1, \dots, \mu_r) \in \text{Den}_{\times}(S)$, and let $\text{Flag}_{S,\iota,\mu}^{\text{wt iso}}(M)$ denote the preimage of $\text{Flag}_{S,\iota}^{\text{iso}}(M)$ under the canonical bundle projection $\text{Flag}_{S,\iota,\mu}^{\text{wt}}(M) \rightarrow \text{Flag}_{S,\iota}(M)$. This is a splitting smooth submanifold in $\text{Flag}_{S,\iota,\mu}^{\text{wt}}(M)$ called the manifold of *weighted isotropic nonlinear flags of type (S, ι, μ) in M* . The diffeomorphism in (2.26) restricts to a diffeomorphism

$$(3.11) \quad \text{Flag}_{S,\iota,\mu}^{\text{wt iso}}(M) = \text{Emb}_{S_r}^{\text{iso}}(M) \times_{\text{Diff}(S;\iota)} \text{Den}(S)_{\iota,\mu}.$$

Using (3.10) and proceeding as in the proof of Theorem 2.11(c), one readily shows that the canonical inclusion $\text{Flag}_{S,\iota,\mu}^{\text{wt iso}}(M) \subseteq \prod_{i=1}^r \text{Gr}_{S_i}^{\text{wt iso}}(M)$ is a splitting smooth submanifold.

Recall the $\text{Diff}(\mathcal{S}; \iota)$ equivariant linear map $h_{\mathcal{S}, \iota} : \text{Den}(\mathcal{S}) \rightarrow H(\mathcal{S}, \iota)$ in (2.28) with kernel

$$\ker Th_{\mathcal{S}, \iota} = \{(d\gamma_1, \dots, d\gamma_r) : \gamma_i \in \Omega^{k_i-1}(\mathcal{S}_i; \mathcal{O}_{\mathcal{S}_i}), \iota_{i-1}^* \gamma_i = 0\},$$

whose restriction to $\text{Den}_\times(\mathcal{S})$ we also denote by $h_{\mathcal{S}, \iota}$. The product with the isodrastic distribution gives the integrable distribution $\mathcal{D} \times \ker Th_{\mathcal{S}, \iota}$ on $\text{Emb}_{\mathcal{S}_r}^{\text{iso}}(M) \times \text{Den}_\times(\mathcal{S})$, of finite codimension $\dim H^1(\mathcal{S}_r; \mathbb{R}) + \dim H(\mathcal{S}, \iota)$. This $\text{Diff}(\mathcal{S}; \iota)$ invariant distribution descends to an integrable distribution

$$\bar{\mathcal{D}} = \mathcal{D} \times_{\text{Diff}(\mathcal{S}; \iota)} \ker Th_{\mathcal{S}, \iota}$$

of the same codimension on $\text{Flag}_{\mathcal{S}, \iota}^{\text{wt iso}}(M)$. The image of $\bar{\mathcal{D}}$ under the forgetful map is an integrable distribution on $\text{Flag}_{\mathcal{S}, \iota}^{\text{iso}}(M)$ of codimension $\dim H^1(\mathcal{S}_r; \mathbb{R})$, which coincides with the distribution that descends from the $\text{Diff}(\mathcal{S}; \iota)$ invariant isodrastic distribution \mathcal{D} on $\text{Emb}_{\mathcal{S}_r}^{\text{iso}}(M)$ by the principal bundle projection. Using Proposition 2.10(c) and (3.11), we see that each leaf \mathcal{F} of $\bar{\mathcal{D}}$ is a connected component in the preimage of an isodrast \mathcal{L} in $\text{Gr}_{\mathcal{S}_r}^{\text{iso}}(M)$ under the bundle projection $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt iso}}(M) \rightarrow \text{Gr}_{\mathcal{S}_r}^{\text{iso}}(M)$, for some volume density μ on \mathcal{S} . Hence, each leaf \mathcal{F} of $\bar{\mathcal{D}}$ is a splitting smooth submanifold of codimension $\dim H^1(\mathcal{S}_r; \mathbb{R})$ in $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt iso}}(M)$.

Remark 3.10 If $H^1(\mathcal{S}_r; \mathbb{R}) = 0$, then the leaves of the isodrastic foliation $\bar{\mathcal{D}}$ are the connected components of $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt iso}}(M)$.

Proposition 3.11 *The $\text{Ham}_c(M)$ action on each leaf $\mathcal{F} \subseteq \text{Flag}_{\mathcal{S}, \iota}^{\text{wt iso}}(M)$ of the isodrastic foliation $\bar{\mathcal{D}}$ admits local smooth sections.*

Proof Suppose \mathcal{L} is an isodrastic leaf in $\text{Gr}_{\mathcal{S}_r}^{\text{iso}}(M)$, and let $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ denote its preimage under the canonical projection $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt iso}}(M) \rightarrow \text{Gr}_{\mathcal{S}_r}(M)$. It suffices to show that the $\text{Ham}_c(M)$ action on $\text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}}$ admits local smooth sections. The diffeomorphism in (3.11) restricts to a diffeomorphism

$$(3.12) \quad \text{Flag}_{\mathcal{S}, \iota, \mu}^{\text{wt iso}}(M)|_{\mathcal{L}} = \text{Emb}_{\mathcal{S}_r}^{\text{iso}}(M)|_{\mathcal{L}} \times_{\text{Diff}(\mathcal{S}; \iota)} \text{Den}(\mathcal{S})_{\iota, \mu}.$$

By Proposition 3.1, the $\text{Ham}_c(M)$ action on $\text{Emb}_{\mathcal{S}_r}^{\text{iso}}(M)|_{\mathcal{L}}$ admits local smooth sections. According to Proposition 2.10(c), the $\text{Diff}(\mathcal{S}; \iota)$ action on $\text{Den}(\mathcal{S})_{\iota, \mu}$ admits local smooth sections too. With the help of Lemma A.1, we conclude that the $\text{Ham}_c(M)$ action on the associated bundle (3.12) admits local smooth sections. ■

Corollary 3.12 *The $\text{Ham}_c(M)$ action on each leaf of the isodrastic foliation on $\text{Flag}_{\mathcal{S}, \iota}^{\text{iso}}(M)$ admits local smooth sections.*

3.4 Weighted isotropic nonlinear flag manifolds as coadjoint orbits

In this section, we describe coadjoint orbits of the Hamiltonian group consisting of weighted isotropic nonlinear flags.

We aim at defining a leafwise symplectic form $\tilde{\Omega}$ on $(\text{Flag}_{\mathcal{S},\iota}^{\text{wt iso}}(M), \tilde{\mathcal{D}})$. We start with leafwise differential 2-forms Ω_i on $(\text{Emb}_{S_i}^{\text{iso}}(M) \times \text{Den}_{\times}(S_i), \mathcal{D}_i \times \ker Th_{S_i})$, for $i = 1, \dots, r$, defined as in Lemma 3.6. Let $j_i := \iota_{r-1} \circ \dots \circ \iota_i \in \text{Emb}(S_i, S_r)$. We consider

$$q_i := j_i^* \times \text{pr}_i : \text{Emb}_{S_r}^{\text{iso}}(M) \times \text{Den}_{\times}(S) \rightarrow \text{Emb}_{S_i}^{\text{iso}}(M) \times \text{Den}_{\times}(S_i),$$

which maps the distribution $\mathcal{D} \times \ker Th_{\mathcal{S},\iota}$ to $\mathcal{D}_i \times \ker Th_{S_i}$. Then

$$(3.13) \quad \Omega := \sum_{i=1}^r q_i^* \Omega_i$$

is a closed leafwise differential 2-form on $(\text{Emb}_{S_r}^{\text{iso}}(M) \times \text{Den}_{\times}(S), \mathcal{D} \times \ker Th_{\mathcal{S},\iota})$.

Remark 3.13 The image under Tq_i of the distribution $\mathcal{D} \times \ker Th_{\mathcal{S},\iota}$ is, in general, strictly included in the distribution $\mathcal{D}_i \times \ker Th_{S_i}$. The reason is that the projection on the i th factor $\text{pr}_i : \ker Th_{\mathcal{S},\iota} \rightarrow \ker Th_{S_i}$ is not surjective in general. This happens, for instance, in the setting of Example 3.19, where $\iota : \{1, \dots, k\} \rightarrow S^1$ with $k \geq 2$: the projection on the second factor is $d\{\gamma \in C^\infty(S^1) : \gamma \circ \iota = 0\}$, strictly included in $\ker Th_{S^1} = dC^\infty(S^1)$.

The $\text{Diff}(\mathcal{S}; \iota)$ action on $\text{Emb}_{S_r}^{\text{iso}}(M) \times \text{Den}_{\times}(S)$,

$$(3.14) \quad g \cdot (\varphi, (\alpha_i)) = (\varphi \circ g_r, (g_i^* \alpha_i)), \text{ for } g = (g_1, \dots, g_r) \in \text{Diff}(\mathcal{S}; \iota),$$

has infinitesimal generators of the form

$$(3.15) \quad \zeta_Z(\varphi, (\alpha_i)) = (T\varphi \circ Z_r, (L_{Z_i} \alpha_i)), \text{ for } Z = (Z_1, \dots, Z_r) \in \mathfrak{X}(\mathcal{S}; \iota).$$

They belong to the integrable distribution $\mathcal{D} \times \ker Th_{\mathcal{S},\iota}$. Notice that $q_i = j_i^* \times \text{pr}_i$ is equivariant over the homomorphism $g \in \text{Diff}(\mathcal{S}; \iota) \mapsto g_i \in \text{Diff}(S_i)$, because $j_i \circ g_i = g_r \circ j_i$ for all i . Now, from the $\text{Diff}(S_i)$ invariance of Ω_i and (3.13), we deduce the $\text{Diff}(\mathcal{S}; \iota)$ invariance of Ω .

Proposition 3.14 *The kernel of the leafwise differential 2-form Ω on $(\text{Emb}_{S_r}^{\text{iso}}(M) \times \text{Den}_{\times}(S), \mathcal{D} \times \ker Th_{\mathcal{S},\iota})$ in (3.13) is spanned by the infinitesimal generators of the $\text{Diff}(\mathcal{S}; \iota)$ action (3.14) on $\text{Emb}_{S_r}^{\text{iso}}(M) \times \text{Den}_{\times}(S)$.*

Proof The contraction of Ω with an infinitesimal generator ζ_Z , given in (3.15), vanishes for all $Z \in \mathfrak{X}(\mathcal{S}; \iota)$. This follows from the analogous statement for Ω_i and the infinitesimal generators ζ_{Z_i} on $\text{Emb}_{S_i}^{\text{iso}}(M) \times \text{Den}_{\times}(S_i)$ in Lemma 3.6, together with the fact that the infinitesimal generators ζ_Z and ζ_{Z_i} are q_i related (cf. (3.13)).

To prove the converse implication, we will proceed by induction on r , the length of \mathcal{S} , using Lemma 3.7. The formula for the leafwise 2-form in Lemma 3.6 provides a similar formula for Ω :

$$(3.16) \quad \begin{aligned} \Omega_{(\varphi, (\alpha_i))}((u_\varphi, (d\lambda_i)), (v_\varphi, (d\gamma_i))) &= \sum_{i=1}^r \int_{S_i} j_i^* \omega(u_\varphi, v_\varphi) \alpha_i \\ &- \sum_{i=1}^r \int_{S_i} j_i^* (\varphi^* i_{u_\varphi} \omega) \wedge \gamma_i + \sum_{i=1}^r \int_{S_i} j_i^* (\varphi^* i_{v_\varphi} \omega) \wedge \lambda_i. \end{aligned}$$

Suppose $(u_\varphi, (d\lambda_i))$ is in the kernel of $\Omega_{(\varphi, (\alpha_i))}$. Considering $\gamma_1 = \dots = \gamma_{r-1} = 0$, we obtain

$$(3.17) \quad \int_{S_r} \omega(u_\varphi, v_\varphi) \alpha_r - \int_{S_r} \varphi^* i_{u_\varphi} \omega \wedge \gamma_r + \int_{S_r} \varphi^* i_{v_\varphi} \omega \wedge \lambda_r = 0,$$

for all γ_r with $\iota_{r-1}^* \gamma_r = 0$, and all v_φ with $v_\varphi|_{\iota_{r-1}(S_{r-1})} = 0$. By Lemma 3.7, there exists $Z_r \in \mathfrak{X}(S_r)$ such that $T\varphi \circ Z_r = u_\varphi$ and $L_{Z_r} \alpha_r = d\lambda_r$. Using Lemma 3.6, we conclude that (3.17) holds for all γ_r and v_φ . Combining this with (3.16), we see that $(u_\varphi \circ \iota_{r-1}, (d\lambda_i))$ is in the kernel of the leafwise differential 2-form $\Omega'_{(\varphi \circ \iota_{r-1}, (\alpha_i))}$ on $(\text{Emb}_{S_{r-1}}^{\text{iso}}(M) \times \text{Den}_\times(S'), \mathcal{D}' \times \ker Th_{S', \iota'})$ where $S' = (S_1, \dots, S_{r-1})$ and $\iota' = (\iota_1, \dots, \iota_{r-2})$. By induction, there exists $Z' = (Z_1, \dots, Z_{r-1}) \in \mathfrak{X}(S', \iota')$ such that $T(\varphi \circ \iota_{r-1}) \circ Z_{r-1} = u_\varphi \circ \iota_{r-1}$ and $L_{Z_i} \alpha_i = d\lambda_i$ for $i = 1, \dots, r-1$. Combining this with $T\varphi \circ Z_r = u_\varphi$ and using the injectivity of $T\varphi$, we obtain $T\iota_{r-1} \circ Z_{r-1} = Z_r \circ \iota_{r-1}$. Hence, $Z = (Z_1, \dots, Z_r) \in \mathfrak{X}(S, \iota)$ and $(u_\varphi, (d\lambda_i))$ is the infinitesimal generator at $(\varphi, (\alpha_i))$. This ensures that the kernel of Ω is spanned by the infinitesimal generators of the $\text{Diff}(S; \iota)$ action. ■

As a consequence, Ω descends to a leafwise symplectic form $\tilde{\Omega}$ on the associated bundle

$$(\text{Flag}_{S, \iota}^{\text{wt iso}}(M), \tilde{\mathcal{D}}) = (\text{Emb}_{S_r}^{\text{iso}}(M) \times_{\text{Diff}(S, \iota)} \text{Den}_\times(S), \mathcal{D} \times_{\text{Diff}(S, \iota)} \ker Th_{S, \iota}).$$

The restriction of $\tilde{\Omega}$ to $(\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M), \tilde{\mathcal{D}})$ is a leafwise symplectic form as well. Thus, every leaf of $\tilde{\mathcal{D}}$ is symplectic.

Let \mathcal{F} denote a leaf of the isodrastic distribution $\tilde{\mathcal{D}}$ on $\text{Flag}_{S, \iota}^{\text{wt iso}}(M)$, equipped with the symplectic form $\tilde{\Omega}$. Restricting (2.18), we obtain a $\text{Symp}(M)$ equivariant smooth map

$$(3.18) \quad J: \mathcal{F} \rightarrow \mathfrak{ham}_c(M)^*, \quad \langle J(\mathcal{N}, \nu), X_f \rangle = \sum_{i=1}^r \int_{N_i} f \nu_i,$$

where we identify $\mathfrak{ham}_c(M) = C_0^\infty(M)$ as in (3.9). This is a moment map for the (Hamiltonian) action of $\text{Ham}_c(M)$ on $(\mathcal{F}, \tilde{\Omega})$. Indeed, this follows readily by combining (3.13) with the expression for the moment map in (3.9). Moreover, J is injective according to Proposition 2.5. By Proposition 3.11, the $\text{Ham}_c(M)$ action on \mathcal{F} is transitive. Using Proposition 3.9, we thus obtain the following generalization of Theorem 3.8.

Theorem 3.15 *The moment map $J: (\mathcal{F}, \tilde{\Omega}) \rightarrow \mathfrak{ham}_c(M)^*$ in (3.18) is one-to-one onto a coadjoint orbit of the Hamiltonian group $\text{Ham}_c(M)$. The Kostant–Kirillov–Souriau symplectic form ω_{KKS} on the coadjoint orbit satisfies $J^* \omega_{\text{KKS}} = \tilde{\Omega}$.*

Remark 3.16 In Section 3.3, we have seen that the inclusion

$$\text{Flag}_{S, \iota, \mu}^{\text{wt iso}}(M) \subseteq \prod_{i=1}^r \text{Gr}_{S_i, \mu_i}^{\text{wt iso}}(M)$$

is a splitting smooth submanifold. The symplectic leaf \mathcal{F} described in Theorem 3.15 is a splitting symplectic submanifold of a product $\prod_{i=1}^r \mathcal{G}_i$ of symplectic leaves described in Theorem 3.8. To show that this is indeed a splitting smooth submanifold, one can

proceed as in the proof of Theorem 2.11(c), using the fact that each isodrastic leaf in $\text{Flag}_{S,t}^{\text{iso}}(M)$ is a splitting smooth submanifold in a product of isodrastic leaves in $\prod_{i=1}^r \text{Gr}_{S_i}^{\text{iso}}(M)$. The latter can be show by induction on the depth of the flag using Lemma 3.4.

Remark 3.17 The leafwise symplectic form $\tilde{\Omega}$ on $\text{Flag}_{S,t}^{\text{wt iso}}(M)$ can be seen as a Poisson structure whose symplectic leaves are the isodrasts \mathcal{F} from Theorem 3.15 (see Remark 3.4 in [26] about isodrasts in the nonlinear Grassmannian of weighted Lagrangian submanifolds). Restricting (2.18), we obtain a Poisson moment map

$$J : \text{Flag}_{S,t}^{\text{wt iso}}(M) \rightarrow \mathfrak{ham}_c(M)^*, \quad \langle J(\mathcal{N}, \nu), X_f \rangle = \sum_{i=1}^r \int_{N_i} f \nu_i,$$

for the Poisson action of $\text{Ham}_c(M)$ on weighted isotropic nonlinear flags, where we identify $\mathfrak{ham}_c(M) = C_0^\infty(M)$ as before.

Example 3.18 Let M be a symplectic manifold that possesses isotropic submanifolds diffeomorphic to the sphere S^r with $r > 1$. We use the setting of Example 2.19 to describe some coadjoint orbits of $\text{Ham}_c(M)$ consisting of nested weighted spheres. Since $H^1(S^r; \mathbb{R}) = 0$, by Remark 3.10, each connected component of $\text{Flag}_{S,t,\mu}^{\text{wt iso}}(M)$ is a coadjoint orbit of $\text{Ham}_c(M)$. Similarly to (2.44), we get

$$\text{Flag}_{S,t,\mu}^{\text{wt iso}}(M) = \left\{ (\mathcal{N}, \nu) \in \text{Flag}_{S,t}^{\text{wt iso}}(M) \left| \begin{array}{l} \{ \int_{N_i^+} \nu_i, \int_{N_i^-} \nu_i \} = \{ a_i^+, a_i^- \}, \\ \text{where } N_i^\pm \text{ denote the connected} \\ \text{components of } N_i \setminus N_{i-1} \end{array} \right. \right\}.$$

Example 3.19 The coadjoint orbits of $\text{Ham}_c(\mathbb{R}^2)$ consisting of pointed vortex loops, studied in [4], are the lowest-dimensional examples of coadjoint orbits of weighted nonlinear flags as described in Theorem 3.15.

Their type is (S, ι) , with $\iota : \{1, \dots, k\} \rightarrow S^1, \iota(i) = t_i$ consecutive points on the circle. We get the cohomology space

$$H(S, \iota) \cong \mathbb{R}^k \times \mathbb{R}^k,$$

where the class $[\mu] \in H(S, \iota)$ is identified with its integrals over connected components of $\{1, \dots, k\}$ resp. $S^1 \setminus \{t_1, \dots, t_k\}$. These are $\Gamma_i = \int_{\{i\}} \mu_0$ and $w_i = \int_{t_i}^{t_{i+1}} \mu_1$, for $i = 1, \dots, k$. The $\text{Diff}(S, \iota)$ action on $H(S, \iota)$ factorizes through an action of the dihedral group D_{2k} on $\mathbb{R}^k \times \mathbb{R}^k$. More precisely, one let the dihedral group act on a regular k -gon, with Γ_i assigned to the vertex i and w_i assigned to the edge $[i, i + 1]$. For generic density $\mu \in \text{Den}_\times(S)$, the orbit $H(S, \iota)_{[\mu]}$ consists of $2k$ elements. The orbit is a one-point set if and only if $\Gamma_1 = \dots = \Gamma_k$ and $w_1 = \dots = w_k$.

Let us denote the weighted flags of type (S, ι) in \mathbb{R}^2 by $((\{x_1, \dots, x_k\}, \nu_0), (C, \nu_1))$. The area a enclosed by the curve C singles out one isodrast $\mathcal{L}_a \subset \text{Gr}_{S^1}^{\text{iso}}(\mathbb{R}^2)$, as in Example 3.3. Theorem 3.15 implies that each connected component of $\text{Flag}_{S,t,\mu}^{\text{wt iso}}(\mathbb{R}^2)|_{\mathcal{L}_a}$ is a coadjoint orbit of $\text{Ham}_c(\mathbb{R}^2)$. Using the above description of $\text{Diff}(S, \iota)$ action on $H(S, \iota)$, we get

$$(3.19) \quad \text{Flag}_{S,t,\mu}^{\text{wt iso}}(\mathbb{R}^2)|_{\mathcal{L}_a} = \left\{ \left((\{x_1, \dots, x_k\}, v_0), (C, v_1) \right) \left| \begin{array}{l} x_i \text{ consecutive points in } C \in \mathcal{L}_a, \\ \exists \sigma \in D_{2k} \text{ such that} \\ \int_{\{x_i\}} v_0 = \sigma(\Gamma_i), \int_{x_i}^{x_{i+1}} v_1 = \sigma(w_i) \end{array} \right. \right\}.$$

In particular, the invariants are the area a enclosed by the loop C , the point vorticities Γ_i , and the net vorticities $w_i = \int_{x_i}^{x_{i+1}} v_1$ between two consecutive points on the loop.

Example 3.20 Let $\mathcal{L}_{[a_1, a_2]}$ be a union of isodrastic leaves of Lagrangian 2-tori in \mathbb{R}^4 , as in Example 3.3, with $[a_1, a_2]$ the $\text{GL}(2, \mathbb{Z})$ orbit of the pair of action integrals $(a_1, a_2) \in \mathbb{R}^2$. Let S_1 be a disjoint union of k circles, $S_2 = \mathbb{T}^2$ the 2-torus, and $\iota : S_1 \rightarrow S_2$ the embedding that maps the i th circle to the circle $\{t_i\} \times S^1 \subseteq \mathbb{T}^2$, with consecutive points $t_1, \dots, t_k \in S^1$. For fixed density $\mu \in \text{Den}_\times(S)$, we aim at describing the coadjoint orbits of $\text{Ham}_c(\mathbb{R}^4)$ that are connected components of $\text{Flag}_{S,t,\mu}^{\text{wt iso}}(\mathbb{R}^4)|_{\mathcal{L}_{[a_1, a_2]}}$. To this end, we need the isomorphism

$$H(S, \iota) \cong \mathbb{R}^k \times \mathbb{R}^k$$

given by integration of μ_1 over the components of S_1 and of μ_2 over the torus surface between two successive embedded circles. As in the previous example, the $\text{Diff}(S, \iota)$ action on $H(S, \iota)$ factorizes through the dihedral group. One can now express these coadjoint orbits of $\text{Ham}_c(\mathbb{R}^4)$ as in (3.19), with the help of the dihedral group. Thus, the invariants are: the $\text{GL}(2, \mathbb{Z})$ orbit of the two action integrals for the embedded 2-torus, the total weights of the k isotopic loops on the torus, and the partial weights between two such consecutive loops on the torus.

A Transitive actions on associated bundles

The manifolds and Lie groups in this appendix may be infinite-dimensional and are assumed to be modeled on convenient vector spaces as in [14].

Recall that a smooth G action on M is said to admit local smooth sections if every point x_0 in M admits an open neighborhood U and a smooth map $\sigma : U \rightarrow G$ such that $\sigma(x)x_0 = x$, for all $x \in U$. Clearly, such an action is locally and infinitesimally transitive. Due to the lack of a general implicit function theorem, one cannot expect the converse implication to hold for general Fréchet manifolds.

Lemma A.1 Let $P \rightarrow B$ be a principal G -bundle endowed with the action of a Lie group H on P that commutes with the principal G action. Suppose the structure group G acts on another manifold Q , and consider the canonically induced H action on the associated bundle $P \times_G Q$. If the H action on P and the G action on Q both admit local smooth sections, then the H action on $P \times_G Q$ admits local smooth sections too.

Proof Suppose $\xi_0 \in P \times_G Q$. As the canonical projection $P \times Q \rightarrow P \times_G Q$ is a locally trivial smooth bundle [14, Theorem 37.12], there exist an open neighborhood U of ξ_0 and smooth maps $\pi : U \rightarrow P$ and $\rho : U \rightarrow Q$ such that for all $\xi \in U$ we have

$$[\pi(\xi), \rho(\xi)] = \xi.$$

Put $p_0 := \pi(\xi_0) \in P$ and $q_0 := \rho(\xi_0) \in Q$. As the H action on P admits local sections, there exist an open neighborhood V of p_0 in P and a smooth map $\sigma' : V \rightarrow H$ such

that

$$\sigma'(p_0) = e_H \quad \text{and} \quad \sigma'(p)p_0 = p,$$

for all $p \in V$. As the G action on Q admits local sections, there exist an open neighborhood W of q_0 in Q and a smooth map $\sigma'' : W \rightarrow G$ such that

$$\sigma''(q_0) = e_G \quad \text{and} \quad \sigma''(q)q_0 = q,$$

for all $q \in W$. Possibly replacing U with a smaller neighborhood of ξ_0 , we may assume that for all $\xi \in U$ we have $\rho(\xi) \in W$ and $\pi(\xi)\sigma''(\rho(\xi)) \in V$. Hence, $\sigma : U \rightarrow H$,

$$\sigma(\xi) := \sigma'(\pi(\xi)\sigma''(\rho(\xi)))$$

is a well-defined smooth map. For $\xi \in U$ we obtain

$$\begin{aligned} \sigma(\xi)\xi_0 &= \sigma(\xi)[p_0, q_0] = [\sigma(\xi)p_0, q_0] = [\sigma'(\pi(\xi)\sigma''(\rho(\xi)))p_0, q_0] \\ &= [\pi(\xi)\sigma''(\rho(\xi)), q_0] = [\pi(\xi), \sigma''(\rho(\xi))q_0] = [\pi(\xi), \rho(\xi)] = \xi. \end{aligned}$$

Hence, σ is the desired local smooth section of the H action on $P \times_G Q$. ■

We will denote the fundamental vector fields of a smooth G action on M by

$$\zeta_X(x) := \left. \frac{\partial}{\partial t} \right|_{t=0} \exp(tX)x,$$

where $X \in \mathfrak{g}$ and $x \in M$.

Lemma A.2 *Let G be a regular Lie group acting on a smooth manifold M . Suppose every point x_0 in M admits an open neighborhood U and a smooth map $\sigma : TM|_U \rightarrow \mathfrak{g}$ such that*

$$(A.1) \quad \zeta_{\sigma(X)}(x) = X,$$

for all $x \in U$ and $X \in T_x M$. Then the G action on M admits local smooth sections.

Proof We will construct a local section using an argument due to Moser [22, Section 4]. Suppose $c : [0, 1] \rightarrow U$ is a smooth curve. We seek a smooth curve $g : [0, 1] \rightarrow G$ such that

$$(A.2) \quad c(t) = g(t)c(0).$$

Differentiating, we obtain

$$(A.3) \quad c'(t) = \zeta_{\dot{g}(t)}(c(t)),$$

where $\dot{g}(t) := \left. \frac{\partial}{\partial h} \right|_{h=t} g(h)g(t)^{-1}$ denotes the right logarithmic derivative of g .

Since G is regular [14, Definition 38.4], there exists a unique smooth curve $g = \text{Evol}^r(\sigma \circ c')$ in G such that $\dot{g}(t) = \sigma(c'(t))$ and $g(0) = e$. Using (A.1), we see that (A.3) and, thus, (A.2) hold true. Evaluating at $t = 1$, we obtain a smooth map

$$s : C^\infty([0, 1], U) \rightarrow G, \quad s(c) := g(1) = \text{evol}^r(\sigma \circ c').$$

By construction, $c(1) = s(c)c(0)$, for all smooth curves $c : [0, 1] \rightarrow U$.

To obtain a local smooth section for the G action on M , it suffices to compose s with a smooth map $U \rightarrow C^\infty([0, 1], U)$, $x \mapsto c_x$ satisfying $c_x(0) = x_0$ and $c_x(1) = x$. The latter can readily be constructed using a chart for M centered at $x_0 = 0$. Indeed,

shrinking U so that it becomes star-shaped with center $x_0 = 0$ in such a chart, we may use $c_x(t) := tx$. ■

Acknowledgment The first author would like to thank the West University of Timișoara for the warm hospitality. The second author would like to thank the University of Vienna for the warm hospitality. We thank an anonymous referee for very helpful suggestions.

References

- [1] M. Bauer, M. Bruveris, and P. W. Michor, *Overview of the geometries of shape spaces and diffeomorphism groups*. J. Math. Imaging Vision 50(2014), 60–97.
- [2] E. Binz and H. R. Fischer, *The manifold of embeddings of a closed manifold. With an appendix by P. Michor*. In: Differential geometric methods in mathematical physics (proc. internat. conf., tech. Univ. Clausthal, Clausthal-Zellerfeld, 1978), Lecture Notes in Physics, 139, Springer, Berlin–New York, 1981, pp. 310–329.
- [3] I. Ciuclea, A. B. Tumpach, and C. Vizman, *Shape spaces of nonlinear flags*. In: F. Nielsen and F. Barbaresco (eds.), Geometric science of information. GSI 2023, Lecture Notes in Computer Science, 14071, Springer, Cham, 2023.
- [4] I. Ciuclea and C. Vizman, *Pointed weighted vortex loops in 2D ideal fluids*. J. Phys. A Math. Theor. 56(2023), 245201.
- [5] F. Gay-Balmaz and C. Vizman, *Dual pairs in fluid dynamics*. Ann. Glob. Anal. Geom. 41(2012), 1–24.
- [6] F. Gay-Balmaz and C. Vizman, *Principal bundles of embeddings*. Ann. Glob. Anal. Geom. 46(2014), 293–312.
- [7] F. Gay-Balmaz and C. Vizman, *Isotropic submanifolds and coadjoint orbits of the Hamiltonian group*. J. Symp. Geom. 17(2019), 663–702.
- [8] S. Haller and C. Vizman, *Non-linear Grassmannians as coadjoint orbits*. Math. Ann. 329(2004), 771–785.
- [9] S. Haller and C. Vizman, *Non-linear flag manifolds as coadjoint orbits*. Ann. Glob. Anal. Geom. 58(2020), 385–413.
- [10] S. Haller and C. Vizman, *A dual pair for the contact group*. Math. Z. 301(2022), 2937–2973.
- [11] R. S. Hamilton, *The inverse function theorem of Nash and Moser*. Bull. Amer. Math. Soc. 7(1982), 65–222.
- [12] I. H. Jermyn, S. Kurtek, H. Laga, and A. Srivastava, *Elastic shape analysis of three-dimensional objects*. Synth. Lect. Comput. Vision 12(2017), 1–185.
- [13] S. Jung, I. Dryden, and J. S. Marron, *Analysis of principal nested spheres*. Biometrika 99(2012), 551–568.
- [14] A. Kriegel and P. W. Michor, *The convenient setting of global analysis*, Mathematical Surveys and Monographs, 53, American Mathematical Society, Providence, RI, 1997.
- [15] B. Lee, *Geometric structures on spaces of weighted submanifolds*. SIGMA 5(2009), Article no. 099, 46 pp.
- [16] J. M. Lee, *Introduction to smooth manifolds*, Graduate Texts in Mathematics, 218, Springer, New York, 2003.
- [17] J. E. Marsden and A. Weinstein, *Coadjoint orbits, vortices, and Clebsch variables for incompressible fluids*. Phys. D 7(1983), 305–323.
- [18] P. W. Michor, *Manifolds of smooth maps. III. The principal bundle of embeddings of a noncompact smooth manifold*. Cah. Topol. Géom. Diff. 21(1980), 325–337.
- [19] P. W. Michor, *Manifolds of differentiable mappings*, Shiva Mathematics Series, 3, Shiva Publishing Ltd., Nantwich, 1980.
- [20] P. W. Michor, *Manifolds of mappings and shapes. In the legacy of Bernhard Riemann after one hundred and fifty years. Vol. II*, Advanced Lectures in Mathematics, 35.2, International Press, Somerville, MA, 2016, pp. 459–486.
- [21] P. W. Michor, *Manifolds of mappings for continuum mechanics*. In: R. Segev and M. Epstein (eds.), Geometric continuum mechanics, Advances in Mechanics and Mathematics, 42, Birkhäuser, Basel, 2020, pp. 3–75.
- [22] J. Moser, *On the volume elements on a manifold*. Trans. Amer. Math. Soc. 120(1965), 286–294.

- [23] J. Strait, S. Kurtek, E. Bartha, and S. N. MacEachern, *Landmark-constrained elastic shape analysis of planar curves*. J. Amer. Stat. Assoc. 112(2017), 521–533.
- [24] A. Weinstein, *Lectures on symplectic manifolds. Expository lectures from the CBMS regional conference held at the University of North Carolina, March 8–12, 1976*, Regional Conference Series in Mathematics, 29, American Mathematical Society, Providence, RI, 1977.
- [25] A. Weinstein, *Neighborhood classification of isotropic embeddings*. J. Diff. Geom. 16(1981), 125–128.
- [26] A. Weinstein, *Connections of Berry and Hannay type for moving Lagrangian submanifolds*. Adv. Math. 82(1990), 133–159.

Department of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria
e-mail: stefan.haller@univie.ac.at

Department of Mathematics, West University of Timișoara, Bulevardul Vasile Parvan 4, 300223 Timișoara, Romania
e-mail: cornelia.vizman@e-uvt.ro