

ON THE EXISTENCE OF PIECEWISE CONTINUOUS OPTIMAL CONTROLS

E. S. NOUSSAIR

(Received 5 May 1976)

(Revised 18 November 1976)

Abstract

Existence of piecewise optimal control is proved when the cost function includes one or both of (a) a cost of sudden switching (discontinuity) of control variables, and (b) a cost associated with the maximum rate of variation of the control over segments of the path for which the control is continuous.

1. Introduction

Consider a system characterized by the equation

$$\dot{x}(t) = f(t, x(t), u(t)), \quad x(0) = x_0, \tag{1}$$

together with the associated cost functional

$$I(u) = J(x, u, \Phi) = \int_0^T f_0(t, x(t), u(t)) dt + \Phi(u), \tag{2}$$

where $T \in (0, \infty)$ is prespecified. The vectors x and f will be n -dimensional. An admissible control u will be a piecewise-continuous r -vector-valued function on $[0, T]$ which may have values at time t in a non-empty compact set $M(t)$, with the set-valued function M piecewise-continuous in the Hausdorff topology [1]. We will assume the vector function f is continuous in all argument, is continuously differentiable with respect to x , and that the inner product inequality

$$(x, f(t, x, u)) \leq C [1 + |x|^2] \tag{3}$$

holds for some constant C and all $t \in [0, T]$, x , and $u \in M(t)$, where (\cdot, \cdot) and $|\cdot|$ are the usual inner product and norm in E_n , respectively. This condition prevents finite escape time of trajectories and could be replaced by any condition which allows us to restrict attention to a compact set of the (t, x) space. Using condition (3) we can assume that

$$|x(t)| \leq (1 + |x_0|^2) \exp(2CT) = C_1 \tag{4}$$

for any trajectory x of the system (1). The function $f_0(t, x, u)$ is assumed to be non-negative and continuous in all arguments for $t \in [0, T]$, $|x| \leq C_1$, $u \in M(t)$. The

functional $\Phi(u)$ is defined by

$$\Phi(u) = \gamma_1(N(u) + \gamma_2 S(u), \tag{5}$$

where $\gamma_1 \geq 0, \gamma_2 \geq 0, N(u)$ is the number of discontinuities of u on $[0, T]$, and $S(u)$ is given by

$$S(u) = \sup \left\{ \frac{|u(t) - u(t')|}{|t - t'|} : t \neq t' \text{ and } u \text{ is continuous on } [t, t'] \right\}. \tag{6}$$

We shall denote by Ω the set of admissible controls with $S(u) < \infty$.

An important problem in optimal control theory is that of determining whether or not there exists a control function $u^* \in \Omega$ such that $I(u^*)$ equals the infimum of $I(u)$ as $u(\cdot)$ varies over all admissible control functions. In the special case $\gamma_1 = 0 = \gamma_2$ we have the Pontryagin Policy for which existence proofs are available only under restrictive convexity assumptions imposed on the functions f, f_0 . A sample of these results can be found in [2], [3]. Indeed, it is easy to construct quite simple examples for which one can prove by explicit calculations that no optimal control exists in Ω , [3].

In [5, pp. 276–281, Theorem 7] E. B. Lee and L. Markus have considered the case when the set of admissible controls consists of functions u of bounded variations such that $\|Du\| = |u(0) - u(0-)| + \text{var } u \leq E$ for a prescribed uniform bound $E \geq 0$. Moreover, the cost functional contains a non-integral cost term of the form $\gamma(\sup |X(t)|, \|Du\|)$, where γ is a continuous monotone nonincreasing in each of its two arguments. In this paper no uniform bound is prescribed on $\|Du\|$ and the functional $\Phi(u)$ has been chosen to satisfy two requirements:

- (1) It should be possible to prove an existence theorem, to the effect that at least one piecewise continuous optimal control path exists; and
 - (2) It should be possible to find usable necessary conditions for the control path.
- This paper is devoted to point 1. Point 2 is covered in a joint paper by the author and J. M. Blatt [4].

2. Main results

LEMMA 1. *Let $\{u_n\}_{n=1}^\infty$ be a sequence in Ω such that $\Phi(u_n) \leq K, n = 1, 2, \dots$. Then a subsequence $\{u_{n_i}\}$ and a function $u \in \Omega$ can be chosen with the following properties:*

- (1) $\Phi(u_{n_i})$ converges and $\Phi(u) \leq \lim_{i \rightarrow \infty} \Phi(u_{n_i})$.
- (2) If $t_i, i = 1, 2, \dots, N(u), 0 < t_1 < t_2 \dots < t_{N(u)}$, are the points of discontinuities of u in $(0, T)$, then for any sufficiently small $\epsilon > 0$ the sequence $\{u_{n_i}\}$ converges uniformly to u on the intervals $[0, t_1 - \epsilon], [t_1 + \epsilon, t_2 - \epsilon], \dots, [t_{N(u)} + \epsilon, T]$.

PROOF. By choosing a subsequence if necessary and using the hypothesis $\Phi(u_n) \leq K, n = 1, 2, \dots$, we can assume that $\Phi(u_n)$ converges as $n \rightarrow \infty$ and $N(u_n) = N_K$ for some constant N_K independent of n . For simplicity of notations we shall make the following assumptions:

- (1) The functions $u_n, n = 1, 2, \dots$, are scalar valued.
- (2) $N_K = 1$.

Both these assumptions will impose no loss of generality as will be seen from the nature of the proof. Let $0 < t_n < T$ denote the point of discontinuity of u_n . By selecting a subsequence if necessary we can assume that t_n converges to \hat{t} with $0 \leq \hat{t} \leq T$. Assume $0 < \hat{t} < T$.

For any given N let I_N denote the set $I_N = [0, \hat{t} - (1/N)] \cup [\hat{t} + (1/N), T]$. Choose $\xi > 0$ large enough so that $\hat{t} - (1/\xi) > 0$ and $\hat{t} + (1/\xi) < T$. We choose first a subsequence $\{u_{\tilde{n}}^\xi\}$ of $\{u_n\}$ and a function u^ξ which is defined and continuous on I_ξ .

Since $\lim_{n \rightarrow \infty} t_n = \hat{t}$, we can select a subsequence $\{u_{\tilde{n}}\}$ of $\{u_n\}$ such that $t_{\tilde{n}} \in (\hat{t} - (1/\xi), \hat{t} + (1/\xi))$ for all \tilde{n} . By the hypotheses on the set Ω , the sequence $\{u_{\tilde{n}}\} \subset C(I_\xi) =$ the set of continuous functions on I_ξ and $u_n(t) \in M(t)$ a compact subset of R . Furthermore, by the hypotheses $\Phi(u_n) \leq K, n = 1, 2, \dots$, we have that $|u_n(t) - u_n(t')| \leq K|t - t'|$ for all t, t' in I_ξ . Using Ascoli Arzella's Theorem we can select a subsequence $\{u_{\tilde{n}}\}$ of $\{u_{\tilde{n}}\}$ which converges uniformly to a continuous function u^ξ on I_ξ .

We now proceed by induction to select subsequences $\{u_{\tilde{n}}^{\xi+j}\}$ of $\{u_{\tilde{n}}^\xi\}$ and continuous functions $u^{\xi+j}$ on $I_{\xi+j}, j = 1, 2, \dots$, as follows. Suppose $\{u_{\tilde{n}}^{\xi+j}\}$ and $u^{\xi+j}$ has been chosen. To construct $\{u_{\tilde{n}}^{\xi+j+1}\}$ we choose first a subsequence $\{u_{\tilde{n}}^{\xi+j}\}$ of $\{u_{\tilde{n}}^{\xi+j}\}$ such that $t_{\tilde{n}}^{\xi+j} \in (\hat{t} - (\xi+j+1)^{-1}, \hat{t} + (\xi+j+1)^{-1})$ for all \tilde{n} where $t_{\tilde{n}}^{\xi+j}$ is the point of discontinuity of $u_{\tilde{n}}^{\xi+j}$. Applying Ascoli Arzella's Theorem as before, we can select a subsequence $\{u_{\tilde{n}}^{\xi+j+1}\} \subset \{u_{\tilde{n}}^{\xi+j}\} \subset \{u_{\tilde{n}}^\xi\}$ which converges to a continuous function $u^{\xi+j+1}$ on $I_{\xi+j+1} \supset I_{\xi+j}$.

From the above construction we notice that

- (1) The subsequences $\{u_{\tilde{n}}^{\xi+j}\}, j = 0, 1, 2, \dots$, are subsequences of $\{u_n\}$, and $\{u_{\tilde{n}}^{\xi+j+1}\} \subset \{u_{\tilde{n}}^{\xi+j}\}$.
- (2) $u^{\xi+j+1}(t) = u^{\xi+j}(t)$ for $t \in I_{\xi+j} \subset I_{\xi+j+1}$.

Define $u(t) = u^{\xi+j}(t)$ if $t \in I_{\xi+j}, j = 0, 1, 2, \dots$. Since $\bigcup_{j=1}^\infty I_{\xi+j} = [0, \hat{t}] \cup (\hat{t}, T]$, the function $u(t)$ is defined and continuous on $[0, T] - \{\hat{t}\}$. Let $\{u_n\}$ denote the subsequence defined by

$$u_{n_i} = u_{\hat{t}^i}^{\xi+i}, \quad i = 1, 2, \dots$$

We show now that the sequence $\{u_n\}$ and the function u defined above satisfy the required properties (1) and (2) of Lemma 1.

First notice that \hat{t} is the only possible point of discontinuity of u . Hence $N(u) \leq N_K$. For any given $\epsilon > 0$ there is $\eta = \eta(\epsilon)$ such that $I_{\xi+\eta} \supset [0, \hat{t} - \epsilon] \cup [\hat{t} + \epsilon, T]$ and $\{u_n, j \geq \eta\}$ converges uniformly to u on $I_{\xi+\eta}$. This proves property (2).

We notice that $\lim_{t \rightarrow \hat{t}^-} u(t)$ and $\lim_{t \rightarrow \hat{t}^+} u(t)$ both exist. This follows from the inequality

$$\begin{aligned} |u(t) - u(t')| &= \lim |u_{n_i}(t) - u_{n_i}(t')| \\ &\leq K|t - t'| \end{aligned}$$

for $t, t' \in [0, \hat{t})$ or $t, t' \in (\hat{t}, T]$. If \hat{t} is a point of continuity of u then $\{u_{n_i}(t)\}$ converges to $u(t)$. Hence $u(t) \in M(t)$.

To prove property (1) we notice first that $\Phi(u) = \gamma_1 N(u) + \gamma_2 S(u) \leq \gamma_1 N_K + \gamma_2 S(u)$ and $\lim_{i \rightarrow \infty} \Phi(u_{n_i}) = \gamma_1 N_K + \gamma_2 \lim S(u_{n_i})$. Hence it is enough to show that

$$S(u) \leq \lim_{i \rightarrow \infty} S(u_{n_i}) = S^* \tag{*}$$

To prove (*) we consider the following two cases:

(I) ℓ is a point of discontinuity of $u(t)$: Let $t \neq t'$ and t, t' belong to either $[0, \ell)$ or $(\ell, T]$. Then

$$\begin{aligned} |u(t) - u(t')| &\leq |u(t) - u_{n_i}(t)| + |u_{n_i}(t) - u_{n_i}(t')| + |u_{n_i}(t') - u(t')| \\ &\leq \lim_{i \rightarrow \infty} S(u_{n_i}) |t - t'|. \end{aligned} \tag{7}$$

(II) $u(t)$ is continuous on $[0, T]$: Assume $t < \ell < t'$ and $t, t' \in [0, T]$.

Let $\epsilon > 0$ and choose $\delta > 0$ such that

$$|u(\ell - \delta) - u(\ell + \delta)| < \epsilon. \tag{8}$$

Using (7) and (8) we obtain

$$\begin{aligned} |u(t) - u(t')| &\leq |u(t) - u(\ell - \delta)| + \epsilon + |u(\ell + \delta) - u(t')| \\ &\leq S^*(t - t') + \epsilon \end{aligned} \tag{9}$$

for any $\epsilon > 0$. Hence

$$|u(t) - u(t')| \leq S^*(t - t'). \tag{10}$$

Using (7) and (10) we obtain (*). This completes the proof of Lemma 1.

DEFINITION. Let

$$\Omega_K = \{u \in \Omega : \Phi(u) \leq K\}. \tag{11}$$

LEMMA 2. Let $\{u_n\}_{n=1}^\infty$ be a sequence in Ω_K . Then there exists a subsequence $\{u_{n_i}\} \subset \{u_n\}$ and a function $u \in \Omega_K$ such that $\Phi(u_{n_i})$ converges as $i \rightarrow \infty$, and

$$\lim_{i \rightarrow \infty} I(u_{n_i}) = \int_0^T f_0(t, x(t), u(t)) dt + \lim_{i \rightarrow \infty} \Phi(u_{n_i}). \tag{12}$$

PROOF. By Lemma 1 we can choose a subsequence $\{u_{n_i}\}$ and an element $u \in \Omega_K$ satisfying properties (1) and (2) of Lemma 1.

Let $x_{n_i}(t), x(t)$ denote the trajectories of the initial value problem (1) corresponding to u_{n_i}, u , respectively. Since $M(t)$ is continuous in the Hausdorff topology for $t \in [0, T]$ there is a constant L such that $|u_{n_i}(t)| \leq L, |u(t)| \leq L, t \in [0, T]$. Also, f is assumed continuously differentiable with respect to x . Let C be the maximum of the absolute value of all partial derivatives $(\partial/\partial x_j)(f_i(t, x, u))$ for $t \in [0, T], |x| \leq C_1$,

$|u| \leq L$. Then

$$\begin{aligned}
 |x_{n_i}(t) - x(t)| &\leq \int_0^t |f(x_{n_i}(t), u_{n_i}(t), t) - f(x(t), u(t), t)| dt \\
 &\leq \int_0^t |f(x_{n_i}(t), u_{n_i}(t), t) - f(x(t), u_{n_i}(t), t)| dt \\
 &\quad + \int_0^T |f(x(t), u_{n_i}(t), t) - f(x(t), u(t), t)| dt \\
 &\leq C \int_0^t |x_{n_i}(t) - x(t)| dt + \alpha_i,
 \end{aligned}
 \tag{13}$$

where

$$\alpha_i = \int_0^T |f(x(t), u_{n_i}(t), t) - f(x(t), u(t), t)| dt.$$

An application of Gronwall inequality [1] to (13) gives

$$|x_{n_i}(t) - x(t)| \leq \alpha_i e^{CT}.
 \tag{14}$$

Let $g_i(t) = |f(t, x(t), u_{n_i}(t)) - f(t, x(t), u(t))|$. Since $u_{n_i}(t)$ converges to $u(t)$ for each $t \neq t_i$, $i = 1, 2, \dots, N(u)$, and $f(t, x, u)$ is assumed continuous in all arguments for $t \in [0, T]$, $|x| \leq C_1$, $|u| \leq L$, $g_i(t)$ converges to 0 for $t \neq t_i$, $i = 1, 2, \dots, N(u)$. Also, $g_i(t) \leq 2L$. Hence the Dominated Convergence Theorem implies that $\lim_{i \rightarrow \infty} \alpha_i = 0$. We then conclude from (14) that x_{n_i} converges uniformly to x on $[0, T]$. We obtain by a similar application of the Dominated Convergence Theorem that

$$\lim_{i \rightarrow \infty} \int_0^T f_0(t, x_{n_i}(t), u_{n_i}(t)) dt = \int_0^T f_0(t, x(t), u(t)) dt$$

and the desired conclusion (12) follows.

THEOREM 3 (Existence of an optimal control): *There exists a function $u^* \in \Omega$ such that $I(u^*) = \inf \{I(u) : u \in \Omega\}$.*

PROOF: Let $K = I(0)$. Then we claim that

$$\inf \{I(u) : u \in \Omega\} = \inf \{I(u) : u \in \Omega, \Phi(u) \leq K\}.
 \tag{15}$$

To prove (15) it is enough to show that

$$\inf \{I(u) : u \in \Omega\} \geq \inf \{I(u) : u \in \Omega, \Phi(u) \leq K\}.
 \tag{16}$$

So assume to the contrary that there is $u \in \Omega$ with $\Phi(u) > K$ such that

$$I(u) < \inf \{I(u) : u \in \Omega, \Phi(u) \leq K\}
 \tag{17}$$

But $I(u) \geq \Phi(u) > K$. From (17) we then obtain

$$K < \Phi(u) \leq I(u) < I(0) = K$$

which is a contradiction. This proves (15).

Let $\alpha_K = \inf\{I(u) : u \in \Omega, \Phi(u) \leq K\}$. Choose $\{u_n\} \subset \Omega$ with $\Phi(u_n) \leq K$ such that

$$\alpha_K = \lim_{n \rightarrow \infty} I(u_n). \tag{18}$$

By choosing a subsequence if necessary we can assume that $\Phi(u_n)$ converges. Let

$$\Phi_K^* = \lim_{n \rightarrow \infty} \Phi(u_n). \tag{19}$$

By Lemma 1 we can select a subsequence $\{u_{n_i}\} \subset \{u_n\}$ and a function $u^* \subset \Omega$ satisfying properties (1) and (2) of Lemma 1. By Lemma 2 we have

$$\lim_{i \rightarrow \infty} I(u_{n_i}) = \int_0^T f_0(t, x(t), u^*(t)) dt + \lim_{i \rightarrow \infty} \Phi(u_{n_i}). \tag{20}$$

By (18), (19) and (20) we have

$$\alpha_K = \lim_{i \rightarrow \infty} I(u_{n_i}) = \int_0^T f_0(t, x(t), u^*(t)) dt + \Phi_K^*. \tag{21}$$

But from Lemma 1 we have

$$\Phi(u^*) \leq \Phi_K^*. \tag{22}$$

Using (21) and (22) we obtain

$$I(u^*) = \int_0^T f_0(t, x(t), u^*(t)) dt + \Phi(u^*) \leq \alpha_K. \tag{23}$$

Using (15), (18), (22) and (23) we obtain

$$I(u^*) = \inf\{I(u) : u \in \Omega\}.$$

This proves Theorem 3.

REMARK 1. The hypotheses on the function f can be relaxed. Instead of requiring continuous differentiability of f with respect to x , it is enough to require the existence of a non-negative, measurable and locally Lebesgue integrable function L defined on $[0, T]$ so that

$$|f(t, x, u) - f(t, y, u)| \leq L(t)|x - y| \quad \text{almost everywhere on } [0, T] \text{ for all } u \in \Omega$$

REMARK 2. Other control problems [Lagrange problems, free problems ($m = u$, $f = u$)] can be considered. These will appear elsewhere.

REMARK 3. Other choices of the term $\Phi(u)$, representing the cost of ‘‘variation’’ in the control u , are possible and lead to interesting problems. Let us write Ω_Φ to denote the class of admissible controls associated with the cost functional (2). Then the following problems admit existence of optimal controls:

- I. $\Phi = \gamma_1 N(u)$, $\gamma_1 > 0$, $\Omega_\Phi = \{u \in \Omega : S(u) \leq K_0\}$, where K_0 is some prescribed constant. In this case it is clear that Theorem 3 still holds. An optimal control will depend on K_0 .

- II. $\Phi = \gamma_2 S(u)$, $\Omega_\Phi = \{u \in C[0, T]: u(t) \in M(t)\}$. In this case one can easily prove that an optimal control exists by a simple application of Ascoli Arzella's Theorem.
- III. $\Phi = \gamma T[u]$, where $T[u]$ is the total variation of u on $[0, T]$.
 $\Omega_\Phi = \{u: u(t) \text{ is of bounded variation on } [0, T] \text{ and } u(t) \in M \text{ a compact subset of } E_m\}$.

Obviously there are several other choices of Φ and Ω_Φ . However, one has to make sure that Ω_Φ is not too restrictive to allow for usable necessary conditions. Necessary conditions for the control path and a discussion of problems I and II are given in [4].

The author wishes to thank Professor J. M. Blatt for valuable discussions.

References

- [1] H. Hermes and J. P. La Salle, *Functional Analysis and Time Optimal Control*, Mathematics in Science and Engineering, Vol. 56, Academic Press, New York and London (1969).
- [2] L. S. Pontryagin, V. G. Boltyanskii, R. V. Gamkrelidze and E. F. Mishchenko, *The Mathematical Theory of Optimal Processes*, Gosudarst, Moscow, 1961; English transl. Interscience, New York (1962), Pergamon Press, New York (1964).
- [3] L. Cesari, "An existence theorem in problems of optimal control", *J. SIAM Control*, Ser. A, Vol. 3, No. 1 (1965).
- [4] E. S. Noussair and J. M. Blatt, "Optimal control with costs of rapid variation of control" (submitted).
- [5] E. B. Lee and L. Markus, *Foundations of Optimal Control Theory*, The SIAM Series in Applied Mathematics, John Wiley and Sons, New York, London and Sydney (1967).

School of Mathematics
 University of New South Wales
 Sydney, N.S.W., 2033
 Australia