

---

*Seventh Meeting, 14th May 1897.*

---

Professor GIBSON in the Chair.

---

**The Bessel Functions and their Zeros.**

By Dr PEDDIE.

---

**A Geometrical Theorem with application to the Proof of the Collinearity of the mid-points of the Diagonals of the Complete Quadrilateral.**

By R. F. MUIRHEAD, M.A., B.Sc.

---

**Geometrical Note.**

By R. TUCKER, M.A.

On the sides BC, CA, AB of the triangle ABC are described two sets of equilateral triangles,  
 the set Ba'C, Cb'A, Ac'B externally, and  
 the set BaC, CbA, AcB internally.

The lines Aa', Bb', Cc' conintersect in Q, the centre of Perspective of the triangles ABC, a'b'c',

and the lines Aa, Bb, Cc in P, the centre of Perspective of ABC, abc.

Since a, a', b, b', c, c' are on the perpendicular bisectors of BC, CA, AB, their joins conintersect in the circumcentre, O, which is the centre of Perspective of abc, a'b'c'.

Now 
$$\begin{aligned} Oa' &= 2R\cos(60^\circ - A), \\ Oa &= -2R\cos(60^\circ + A), \end{aligned}$$

hence 
$$aa' = a\sqrt{3}, \quad bb' = b\sqrt{3}, \quad cc' = c\sqrt{3};$$

and also 
$$\Sigma(aa')^2 = 3 \quad \Sigma(a^2) = 3k.$$

Using trilinear coordinates,

Q is 
$$a\sin(60^\circ + A) = \beta\sin(60^\circ + B) = \gamma\sin(60^\circ + C);$$

P is 
$$a\sin(60^\circ - A) = \beta\sin(60^\circ - B) = \gamma\sin(60^\circ - C).$$