



Weeds

VOLUME IV
JULY 1956
NUMBER 3

Journal of the Weed Society of America

Weeds

Issued Quarterly by the Weed Society of America

K. P. BUCHHOLTZ, *Editor*, Dept. of Agronomy, Univ. of Wisconsin,
Madison, Wisconsin.

W. C. JACOB, *Business Manager*, Dept. of Agronomy, Univ. of
Illinois, Urbana, Illinois.

EDITORIAL COMMITTEE

R. J. ALDRICH, Weed Investigations Section, USDA, Rutgers Univ.,
New Brunswick, New Jersey

O. C. LEE, Dept. of Botany and Plant Pathology, Purdue Univ.,
Lafayette, Indiana.

W. C. ROBOCKER, Weed Investigations Section, USDA, Univ. of
Nevada, Reno, Nevada.

W. K. PORTER, JR., Dept. of Plant Pathology, Univ. of Louisiana,
Baton Rouge, Louisiana.

WEEDS is a quarterly journal published by the Weed Society of America. Editorial offices are located at the University of Wisconsin, Madison, Wisconsin. Printing is by the W. F. Humphrey Press Inc., Geneva, New York. Subscription price is \$6.00 yearly for four issues; single copies \$1.50. Address all communications regarding subscriptions, advertising and reprints to W. C. Jacob, Department of Agronomy, University of Illinois, Urbana, Illinois. Inquiries concerning information on manuscripts and other material for publication should be addressed to the Editorial offices. All checks, money orders and other remittances should be made payable to the Weed Society of America.

Entered as second-class matter at the post office at Urbana,
Illinois with additional entry at Geneva, New York.

Table of Contents

	<i>Page</i>
Leaching and Decomposition of 2,2-Dichloropropionic Acid in Several Iowa Soils. J. T. Holstun, Jr. and W. E. Loomis.....	205
Translocation of Herbicides in Marabú. I. Translocation of 2,4,5-Trichlorophenoxyacetic Acid Following Application to the Bark or to Cut-surfaces of Stumps. J. R. Hay.....	218
Some Factors that Affect the Phytotoxicity of Water-soluble DNBP. D. E. Davis.....	227
Aquatic Weed Control Practice in Potable Water Supply Reservoirs. William K. Weight.....	235
Studies of Factors Affecting the Control of Chamise (<i>Adenostoma fasciculatum</i>) with Herbicides. O. A. Leonard.....	241
The Crop Toxicity Period of CMU in a Sandy Clay Loam Soil. L. L. Danielson.....	255
Studies on the Control of Crabgrass in Bluegrass Lawns. R. E. Nylund and Richard Stadtherr.....	264
A Note on the Sugars and Amino Acids of Leafy Spurge, <i>Euphorbia esula</i> . Duane Le Tourneau.....	275
Report of the Terminology Committee—Weed Society of America.....	278
News and Notes.....	288
Sustaining Members.....	290
Bibliography of Weed Investigations for 1955.....	291

Advertisers Index

du Pont de Nemours & Co.	ii
Standard Agricultural Chemicals Inc.	iii
Chipman Chemical Co.	iv
Diamond Alkali Co.	v
American Cyanamid Co.	vi
Spraying Systems Co.	vii
Carbide and Carbon Chemicals Co.	viii

For Industry and Agriculture . . .

Du Pont UREA HERBICIDES

***offer new economies
and efficiency in killing
weeds, grass and brush***

KARMEX® DL for pre-emergence weed control in cotton.

"KARMEX" W for weed control in asparagus, sugar cane, and pineapple. Also for irrigation and drainage ditch weed control.

TELVAR® for industrial weed and grass control. Also in certain areas, it is recommended for brush control. Available in two formulations: "Telvar" W and "Telvar" DW.

The urea herbicides, products of Du Pont research, kill vegetation through the roots. Their efficiency is demonstrated by the relatively low dosages required to do the job. They can be easily applied, are non-flammable, non-volatile, non-corrosive and extremely low in toxicity.



Better Things for Better Living . . . *Through Chemistry*

**E. I. DU PONT DE NEMOURS & CO. (INC.)
GRASELLI CHEMICALS DEPT.**

WILMINGTON 98, DELAWARE

depend upon **SINOX**

for

.....
DINITRO
.....
WEED
.....
CONTROL

Our field research technical experts—alert to all current developments—constantly strive toward improved chemical products and techniques. We are devoted to the idea of *progress* in weed control.

SINOX W For selective spraying of seeding alfalfa, onions, flax, peas and grain seeded to legumes.

SINOX PE For pre-emergence spraying of potatoes, beans, corn and peas; control of winter annuals in fall seeded alfalfa and clover.

SINOX GENERAL As a desiccant for Seed Alfalfa, Clovers, Sudan Grass, Milo, and Flax. Used for general weed control in orchards, vineyards, irrigation ditches and roadsides.
—Potato Vine Killing.

STANDARD AGRICULTURAL CHEMICALS, INC.



1301 Jefferson St., Hoboken, N. J.
429 Forum Building, Sacramento, Calif.



ATLACIDE: A chlorate weed killer...widely used for non-selective eradication of bindweed, Canada thistle, quack grass, Johnson grass and other tough perennials. Kills roots...discourages regrowth. Applied as spray or in original dry form.

ATLACIDE WITH 2, 4-D: A combination of Atlacide and 2,4-D acid. Particularly recommended for Canada thistle control.

CHLOREA: A uniform, non-separating combination of sodium chlorate, borate and CMU. Kills all types of weeds and grasses. Combines the proven effectiveness of chlorate on deep-rooted weeds with the soil-surface action of CMU on shallow-rooted grasses and annual seedling growth. Has lasting residual effect to inhibit new growth. Does not create a fire hazard when used as directed. Applied dry or as a water-mixed spray. For industrial, railroad and certain agricultural uses.

CHLORAX "40": A non-separating composition of sodium chlorate and borate...for weed and grass control. Has a lasting residual effect. Does not create a fire hazard. Applied dry or as a spray.

CHLORAX LIQUID: Similar to Chlorax "40"...in liquid form for easy mixing with water.

ATLAS "A": A 40% sodium arsenite solution (4 lbs. arsenic trioxide per gal.). Destroys certain submerged vegetation in ponds and lakes. Controls crabgrass, chickweed and clover in turf. Used as general weed killer and to kill trees and stumps. Also used to kill potato vines prior to harvesting.

SODIUM ARSENITE: A powder containing 75% arsenic trioxide. Used for the same purposes as Atlas "A". Applied dry or as a spray.

2, 4-D WEED KILLERS: A complete line—Available as 2,4-D Amine and 2,4-D Ester liquids; also 2,4-D Ester dusts.

METHOXONE: Contains 2 pounds of MCP sodium salt per gallon. Used for weed control in small grains, flax, rice and grass. Controls same weeds as 2,4-D; considered safer for selective spraying.

Low Volatile 2,4,5-T
 Low Volatile Brush Killer
 Chloro IPC • IPC 25% Liquid

Write for Weed Control Booklets

CHIPMAN CHEMICAL COMPANY, INC.

Chicago, Ill.

BOUND BROOK, N. J.

Portland, Ore.

Palo Alto, Calif.

Pasadena, Tex.

Bessemer, Ala.

Manufacturers of Weed Killers Since 1912

For Dependable Insecticides and Herbicides

think first of

DIAMOND

Technical and Formulations

Insecticides

DIAMOND dependable insecticides have long been favored because of their uniform high quality. DDT, BHC and Lindane Technicals are supplied in many types and forms.

Insecticides for Formulators and Producers

DDT—100% Technical	Lindane—100% Gamma Isomer of BHC
DDT—Wettable Powders	BHC—40% Technical
DDT—Dust Concentrates and Emulsions	BHC—12% Wettable Powders and Dusts
Lindane—Emulsions, Wettable Powders, Dusts	BHC—Emulsion Concentrates
	K-101—Miticide

Weed Killers and Brush Control Chemicals

DIAMOND weed and brush killers, based on the 2,4-D and 2,4-T chemical groups, meet every weed and brush control problem.

These dependable DIAMOND herbicides

are produced in two forms: *Technical* for use by formulators and processors in their own formulations and *ready-to-use formulations* for farmers, ranchers, country clubs and homeowners.

Technical for Formulators

WEED KILLERS

2,4-D Acid • Technical Butyl-D
Technical Isopropyl-D
Technical 2-Ethyl Hexyl (Iso-Octyl)-D (Low Volatile)
Technical BEP-D (Low Volatile)

BRUSH CONTROL

Technical Butyl-T • Technical Isopropyl-T
Technical BEP-T (Low Volatile)
Technical 2-Ethyl Hexyl (Iso-Octyl)-T (Low Volatile)

DIAMOND research is constantly seeking new and better insecticides and herbicides and working with formulators and agricultural chemists in the development of more efficient forms and application methods. We will be glad to work with you. For information on DIAMOND Chemicals and technical co-operation write, DIAMOND ALKALI COMPANY, 300 Union Commerce Bldg., Cleveland 14, O.

Ready-To-Use Formulations

WEED KILLERS

*4# Mixed Amine-D • 4# Dimethylamine-D
2.67# Butyl-D • 4# Butyl-D • 6# Butyl-D
4# 2-Ethyl Hexyl (Iso-Octyl)-D (Low Volatile)
4# BEP-D (Low Volatile) • 3.34# Isopropyl-D

BRUSH CONTROL

*4# Butyl-T • 4# BEP-T (Low Volatile)
4# 2-Ethyl Hexyl (Iso-Octyl)-T (Low Volatile)
2,4-D—2,4,5-T Mixtures • 2#—2# Butyl Brush Killer
2#—2# 2-Ethyl Hexyl Brush Killer (Low Volatile)
2#—2# BEP Brush Killer (Low Volatile)

*Numbers are pounds of 2,4,5-T or 2,4-D acid equivalent per gallon



Diamond Chemicals

A *Cyanamid* **PRODUCTS**

for effective, economical weed control

POTASSIUM CYANATE

for pre-emergence contact weedkilling, post-emergence selective contact weedkilling, top-killing and defoliation. Breaks down rapidly on contact with soil.

AERO[®] CYANAMID, Granular

Contains 20% nitrogen and 70% hydrated lime. For pre-emergence weed control in peas, corn, asparagus and other crops. For pre-seeding weed control in tobacco and other plant beds, for establishing or renovating weed-free turf. In granular form for easy handling and application.

AERO[®] CYANAMID, Special Grade

Contains 21% nitrogen and 70% hydrated lime. In dust form for pre-emergence residual and contact weed control. Defoliates cotton, field beans and other crops. For pre-harvest top-killing of tomatoes and potatoes.

AMINO TRIAZOLE (3-amino-1,2,4-triazole)

(LIMITED QUANTITIES AVAILABLE FOR COMMERCIAL AND EXPERIMENTAL USE)

Herbicide, Defoliant, Growth Regulator

Amino Triazole has demonstrated effectiveness in control of a number of troublesome weeds, including Canada thistle, sow thistle, poison ivy, poison oak, quack grass, nut grass and certain woody species. By virtue of a short residual life in the soil, Amino Triazole can be sprayed on weed infestations a short time before planting without injury to the crop.

Amino Triazole translocates readily through the plant and produces unusual systemic effects, manifested by albinism or chlorophyll inhibition in new growth. This typical effect has continued to show up as long as one year after spraying certain species.

At rates of $\frac{1}{2}$ to $1\frac{1}{2}$ pounds per acre, cotton has been defoliated and re-growth controlled for a sufficient time to permit harvest.

AMERICAN Cyanamid COMPANY


AGRICULTURAL CHEMICALS DIVISION

30 Rockefeller Plaza, New York 20, N. Y.




the precision nozzle for effective spraying


Supplied in a full range of interchangeable orifice tip and strainer sizes to meet every capacity requirement. TeeJet Spray Nozzles for Weed Control by spraying make it possible to take maximum advantage of the chemical and sprayer unit. TeeJet nozzles are precision built and provide a flat spray with uniform distribution. Atomization is properly controlled to give coverage with an absolute minimum of driftage. Patented tip design, with set-back orifice opening protects precision orifice from accidental damage. TeeJet spray nozzles are built for use on spray booms and portable sprayers.



TEEJET SPRAY NOZZLE male pipe connection



TEEJET SPRAY NOZZLE female pipe connection



INTERCHANGEABLE ORIFICE TIPS flat and cone spray types

OFF-CENTER SPRAY NOZZLES

Spraying Systems Spray Nozzles with TeeJet tips are supplied in a variety of special body types to meet any unusual spraying requirement. For example, one type of off-center spray nozzle with swivel body provides a flat spray up to 35 feet wide for spraying areas with a single nozzle, that are not accessible with a boom.

SUPPLEMENTARY EQUIPMENT

Complete accessories relating to nozzle use are supplied. These include strainers, special nozzle fittings, and hand valve equipment.

TeeJet Spray Nozzles are supplied for Weed Control... as well as all other types of agricultural spraying. For complete information and reference data write for Catalog 30.

SPRAYING SYSTEMS CO.
Engineers and Manufacturers

3275 RANDOLPH STREET

BELLWOOD, ILLINOIS

New Herbicides FOR **WEED CONTROL**

**IN... Tomatoes
Sugar Beets
Cucurbits**

**Peanuts
and
Hybrid Seed Corn**

TOMATOES and TRANSPLANTED CROPS

*Experimental Herbicide Natrin** (sodium 2,4,5-trichlorophenoxy-ethyl sulfate). Apply after clean cultivation to get a weed-free harvest without crop injury.

SUGAR BEETS and CUCURBITS

*Experimental Herbicide DCU** (dichloral urea) mixed with the top layer of soil before planting gives annual grass control for as long as three months.

PEANUTS and SEED CORN BREEDING STOCK

Crag Herbicide-1 (SES) cuts peanut production costs. Bigger yields result from increased weed control and less Southern Blight. In corn it is safe on sensitive inbreds, single and double crosses.

*For experimental use only by or under the supervision of Federal or State agencies authorized by law to conduct research in the field of economic poisons. Treated crops should not be used for food or feed.

CRAG AGRICULTURAL CHEMICALS

CARBIDE AND CARBON CHEMICALS COMPANY

A Division of Union Carbide and Carbon Corporation
30 East 42nd Street, New York 17, New York

Gentlemen:

Please send additional information and your application blank for free experimental material for Experimental Herbicide Natrin....., Experimental Herbicide DCU....., and Crag Herbicide-1.....

Name.....

Address.....

City..... State.....

"Crag" is a registered trade-mark of Union Carbide and Carbon Corporation.